

cases	doc_1		doc_2			decision	id		
			authors	<ul style="list-style-type: none"><li>Paul L. Butzer</li><li>Gerhard Schmeisser</li><li>Rudolf L. Stens</li></ul>			DUPLICATES	1101	
	authors	<ul style="list-style-type: none"><li>Butzer, P.</li><li>Schmeisser, G.</li><li>Stens, R.</li></ul>	title	Sobolev Spaces of Fractional Order, Lipschitz Spaces, Readapted Modulation Spaces and Their Interrelations; Applications					
	publication_date	2016-01-01 00:00:00	publication_date	2016-05-04 15:09:07+00:00					
	source	SupportedSources.CROSSREF	source	SupportedSources.ARXIV					
	journal		journal	None					
	volume		volume						
	doi	10.1016/j.jat.2016.08.001	doi						
	urls	<ul style="list-style-type: none"><li>https://api.elsevier.com/content/article/PII:S002190451630065X?httpAccept=text/xml</li><li>https://api.elsevier.com/content/article/PII:S002190451630065X?httpAccept=text/plain</li><li>http://dx.doi.org/10.1016/j.jat.2016.08.001</li></ul>	urls	<ul style="list-style-type: none"><li>http://arxiv.org/pdf/1605.02777v1</li><li>http://arxiv.org/abs/1605.02777v1</li><li>http://arxiv.org/pdf/1605.02777v1</li></ul>					
	id	id-6032634324185341168	id	id-2473659246203455862					
	abstract		abstract	The purpose of this investigation is to extend basic equations and inequalities which hold for functions $f$ in a Bernstein space $B_{\sigma^2}$ to larger spaces by adding a remainder term which involves the distance of $f$ from $B_{\sigma^2}$ . First we present a modification of the classical modulation space $M^{2,1}(\mathbb{R})$ , the so-called readapted modulation space $M^{2,1}_{\text{a}}(\mathbb{R})$ . Our approach to the latter space and its role in functional analysis is novel. In fact, we establish several chains of inclusion relations between $M^{2,1}_{\text{a}}(\mathbb{R})$ and the more common Lipschitz and Sobolev spaces, including Sobolev spaces of fractional order. Next we introduce an appropriate metric for describing the distance of a function belonging to one of the latter spaces from a Bernstein space. It will be used for estimating remainders and studying rates of convergence. In the main part, we present the desired extensions. Our applications include the classical Whittaker-Kotel'nikov-Shannon sampling formula, the reproducing kernel formula, the Parseval decomposition formula, Bernstein's inequality for derivatives, and Nikol'skiĭ's inequality estimating the $L^p(\mathbb{Z})$ norm in terms of the $L^p(\mathbb{R})$ norm.					
	versions		versions						