

cases	doc_1		doc_2		decision	id
					DUPLICATES	636
	authors	<ul style="list-style-type: none">Bogfjellmo, G.Dahmen, R.Schmeding, A.	authors	<ul style="list-style-type: none">Geir BogfjellmoRafael DahmenAlexander Schmeding		
	title	Character groups of Hopf algebras as infinite-dimensional Lie groups	title	Character groups of Hopf algebras as infinite-dimensional Lie groups		
	publication_date	2016-01-01 00:00:00	publication_date	2015-01-21 16:38:13+00:00		
	source	SupportedSources.CROSSREF	source	SupportedSources.ARXIV		
	journal		journal	Ann. Inst. Fourier (Grenoble), 66 no. 5 (2016), p. 2101-2155		
	volume		volume			
	doi	10.5802/aif.3059	doi			
	urls	<ul style="list-style-type: none">http://www.cedram.org/cgi-bin/textserve/AIF_2016__66_5_2101_0http://dx.doi.org/10.5802/aif.3059	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1501.05221v3http://arxiv.org/abs/1501.05221v3http://arxiv.org/pdf/1501.05221v3		
	id	id-8543682470854848039	id	id1046935923355746699		
	abstract		abstract	In this article character groups of Hopf algebras are studied from the perspective of infinite-dimensional Lie theory. For a graded and connected Hopf algebra we construct an infinite-dimensional Lie group structure on the character group with values in a locally convex algebra. This structure turns the character group into a Baker--Campbell--Hausdorff--Lie group which is regular in the sense of Milnor. Furthermore, we show that certain subgroups associated to Hopf ideals become closed Lie subgroups of the character group. If the Hopf algebra is not graded, its character group will in general not be a Lie group. However, we show that for any Hopf algebra the character group with values in a weakly complete algebra is a pro-Lie group in the sense of Hofmann and Morris.		
	versions		versions			