		doc_1		doc_2	decision	id
cases			authors	Tu T. C Nguyen Vu A. Le		
	authors	Nguyen, T.Le, V.	title	Representation of Real Solvable Lie Algebras Having 2-dimensional Derived Ideal and Geometry of Coadjoint Orbits of Corresponding Lie Groups		
			publication_date	2021-05-02 19:16:33+00:00		
	title	Representation of real solvable Lie algebras having 2-dimensional derived ideal and geometry of coadjoint orbits of corresponding Lie groups	source	SupportedSources.ARXIV		
			journal	None		
			volume			
	source	SupportedSources.CROSSREF	doi		DUPLICATES 46	
	journal		urls	• http://arxiv.org/pdf/2105.00536v4		ES 460
	volume			• http://arxiv.org/abs/2105.00536v4		
	doi	10.1142/s1793557122501935		• http://arxiv.org/pdf/2105.00536v4		
	urls	 https://www.worldscientific.com/doi/pdf/10.1142/S1793557122501935 http://dx.doi.org/10.1142/s1793557122501935 	id	id-9012397468683714886		
		7		Let {\Lnk} be the class of all \$n\$-dimensional real solvable Lie algebras having \$k\$-dimensional derived ideals. In 2020 the authors	s III	
	id	id-3659438126554080711		et al. gave a classification of all non 2-step nilpotent Lie algebras of {\Li}. We propose in this paper to study representations of these Lie algebras as well as their corresponding connected and simply connected Lie groups. That is, for each algebra, we give an upper		
	abstract		abstract	bound of the minimal degree of a faithful representation. Then, we give a geometrical description of coadjoint orbits of		
	versions			corresponding groups. Moreover, we show that the characteristic property of the family of maximal dimensional coadjoint orbits of a MD-group studied by K. P. Shum and the second author et al. is still true for the Lie groups considered here. Namely, we prove that,		
				for each considered group, the family of the maximal dimensional coadjoint orbits forms a measurable foliation in the sense of Connes. The topological classification of these foliations is also provided.		
			versions]	