

cases	doc_1		doc_2		decision	id
					DUPLICATES	193
	authors	<ul style="list-style-type: none">Daniela GrassoEmanuele TassiMichael Kraus	authors	<ul style="list-style-type: none">Michael KrausEmanuele TassiDaniela Grasso		
	title	Variational Integrators for Reduced Magnetohydrodynamics	title	Variational Integrators for Reduced Magnetohydrodynamics		
	publication_date	2015-11-30 00:00:00	publication_date	2015-11-30 14:09:55+00:00		
	source	SupportedSources.PAPERS_WITH_CODE	source	SupportedSources.ARXIV		
	journal		journal	Journal of Computational Physics, Volume 321, Pages 435-458, 2016		
	volume		volume			
	doi		doi	10.1016/j.jcp.2016.05.047		
	urls	<ul style="list-style-type: none">https://arxiv.org/pdf/1511.09314v2.pdfhttps://github.com/DDMGNI/viRMHD2D	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1511.09314v2http://dx.doi.org/10.1016/j.jcp.2016.05.047http://arxiv.org/abs/1511.09314v2http://arxiv.org/pdf/1511.09314v2		
	id	id930606907037439324	id	id2264357767703093180		
	abstract	Reduced magnetohydrodynamics is a simplified set of magnetohydrodynamics equations with applications to both fusion and astrophysical plasmas, possessing a noncanonical Hamiltonian structure and consequently a number of conserved functionals. We propose a new discretisation strategy for these equations based on a discrete variational principle applied to a formal Lagrangian. The resulting integrator preserves important quantities like the total energy, magnetic helicity and cross helicity exactly (up to machine precision). As the integrator is free of numerical resistivity, spurious reconnection along current sheets is absent in the ideal case. If effects of electron inertia are added, reconnection of magnetic field lines is allowed, although the resulting model still possesses a noncanonical Hamiltonian structure. After reviewing the conservation laws of the model equations, the adopted variational principle with the related conservation laws are described both at the continuous and discrete level. We verify the favourable properties of the variational integrator in particular with respect to the preservation of the invariants of the models under consideration and compare with results from the literature and those of a pseudo-spectral code.	abstract	Reduced magnetohydrodynamics is a simplified set of magnetohydrodynamics equations with applications to both fusion and astrophysical plasmas, possessing a noncanonical Hamiltonian structure and consequently a number of conserved functionals. We propose a new discretisation strategy for these equations based on a discrete variational principle applied to a formal Lagrangian. The resulting integrator preserves important quantities like the total energy, magnetic helicity and cross helicity exactly (up to machine precision). As the integrator is free of numerical resistivity, spurious reconnection along current sheets is absent in the ideal case. If effects of electron inertia are added, reconnection of magnetic field lines is allowed, although the resulting model still possesses a noncanonical Hamiltonian structure. After reviewing the conservation laws of the model equations, the adopted variational principle with the related conservation laws are described both at the continuous and discrete level. We verify the favourable properties of the variational integrator in particular with respect to the preservation of the invariants of the models under consideration and compare with results from the literature and those of a pseudo-spectral code.		
	versions		versions			