

cases	doc_1		doc_2		decision	id
					NOT DUPLICATES	1905
	authors	<ul style="list-style-type: none">Lingling HouPengcheng Niu	authors	<ul style="list-style-type: none">Lingling HouPengcheng Niu		
	title	A Nash Type result for Divergence Parabolic Equation related to Hormander's vector fields	title	A Nash Type result for Divergence Parabolic Equation related to Hormander's vector fields		
	publication_date	2017-05-17 00:00:00	publication_date	2017-05-17 01:46:40+00:00		
	source	SupportedSources.OPENALEX	source	SupportedSources.ARXIV		
	journal	arXiv (Cornell University)	journal	None		
	volume		volume			
	doi	10.48550/arxiv.1705.06278	doi			
	urls	<ul style="list-style-type: none">https://openalex.org/W2615044592https://doi.org/10.48550/arxiv.1705.06278http://arxiv.org/pdf/1705.06278	urls	<ul style="list-style-type: none">http://arxiv.org/pdf/1705.06278v1http://arxiv.org/abs/1705.06278v1http://arxiv.org/pdf/1705.06278v1		
	id	id-5866969602453843286	id	id7047419319414157508		
	abstract		abstract	In this paper we consider the divergence parabolic equation with bounded and measurable coefficients related to Hormander's vector fields and establish a Nash type result, i.e., the local Holder regularity for weak solutions. After deriving the parabolic Sobolev inequality, (1,1) type Poincar'e inequality of Hormander's vector fields and a De Giorgi type Lemma, the Holder regularity of weak solutions to the equation is proved based on the estimates of oscillations of solutions and the isomorphism between parabolic Campanato space and parabolic Holder space. As a consequence, we give the Harnack inequality of weak solutions by showing an extension property of positivity for functions in the De Giorgi class.		
	versions		versions			