

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none"><li>Dongho Chae</li></ul>	authors	<ul style="list-style-type: none"><li>Dongho Chae</li></ul>	DUPLICATES	21
	title	Nonexistence of Self-similar Singularities in the Ideal Magnetohydrodynamics	title	Nonexistence of self-similar singularities in the ideal magnetohydrodynamics		
	publication_date	2009-12-01 00:00:00	publication_date	2007-04-09 00:00:00		
	source	SupportedSources.OPENALEX	source	SupportedSources.INTERNET_ARCHIVE		
	journal	Archive for Rational Mechanics and Analysis	journal			
	volume	194	volume			
	doi	10.1007/s00205-008-0182-9	doi			
	urls	<ul style="list-style-type: none"><li>https://openalex.org/W1976442759</li><li>https://doi.org/10.1007/s00205-008-0182-9</li><li>http://arxiv.org/pdf/math/0703317</li></ul>	urls	<ul style="list-style-type: none"><li>https://archive.org/download/arxiv-math0703317/math0703317.pdf</li></ul>		
	id	id-2560408112486590670	id	id-5898474138420081907		
	abstract		abstract	In this paper we exclude the scenario of apparition of finite time singularity in the form of self-similar singularities in the ideal magnetohydrodynamic equations, assuming suitable integrability conditions on the vorticity and the magnetic field. We also consider more sophisticated possibility of asymptotically self-similar singularities, which means that the local classical solution converges to the self-similar profile as we approaches to the possible time of singularity. The scenario of asymptotically self-similar singularity is also excluded under suitable conditions on the profile. In the 2D magnetohydrodynamics the magnetic field evolution equations reduce to a divergence free transport equation for a scalar stream function. This helps us to improve the above nonexistence theorems on the self-similar singularities, in the sense that we only need weaker integrability conditions on the profile to prove the results.		
	versions		versions			