cases	doc_1		doc_2		decision	id
	authors	• Laurent Manivel	**authors**	• Laurent Manivel	DUPLICATES	232
	title	Configurations of lines and models of Lie algebras	**title**	Configurations of lines and models of Lie algebras		
	publication_date	2005-07-06 12:24:41+00:00	**publication_date**	2005-07-06 00:00:00		
	source	SupportedSources.ARXIV	**source**	SupportedSources.INTERNET_ARCHIVE		
	journal	Journal of Algebra 304, 1 (2006) 457-486	**journal**			
	volume		**volume**			
	doi		**doi**			
	urls	• http://arxiv.org/pdf/math/0507118v1 • http://arxiv.org/abs/math/0507118v1 • http://arxiv.org/pdf/math/0507118v1	**urls**	• https://archive.org/download/arxiv-math0507118/math0507118.pdf		
	id	id-5693478141935801408	**id**	id1796320959106989510		
	abstract	The automorphism groups of the 27 lines on the smooth cubic surface or the 28 bitangents to the general quartic plane curve are well-known to be closely related to the Weyl groups of E_6 and E_7. We show how classical subconfigurations of lines, such as double-sixes, triple systems or Steiner sets, are easily constructed from certain models of the exceptional Lie algebras. For ${\mathfrak e}\_7$ and ${\mathfrak e}\_8$ we are lead to beautiful models graded over the octonions, which display these algebras as plane projective geometries of subalgebras. We also interpret the group of the bitangents as a group of transformations of the triangles in the Fano plane, and show how this allows to realize the isomorphism $PSL(3,F\_2)\simeq PSL(2,F\_7)$ in terms of harmonic cubes.	**abstract**	The automorphism groups of the 27 lines on the smooth cubic surface or the 28 bitangents to the general quartic plane curve are well-known to be closely related to the Weyl groups of E_6 and E_7. We show how classical subconfigurations of lines, such as double-sixes, triple systems or Steiner sets, are easily constructed from certain models of the exceptional Lie algebras. For e_7 and e_8 we are lead to beautiful models graded over the octonions, which display these algebras as plane projective geometries of subalgebras. We also interpret the group of the bitangents as a group of transformations of the triangles in the Fano plane, and show how this allows to realize the isomorphism PSL(3,F_2)â‰*f* PSL(2,F_7) in terms of harmonic cubes.		
	versions		**versions**			