		doc_1		doc_2	decision	id
cases	authors	Fernando Haas Kellen Alves Pascoal José Tito Mendonça				
	title	Neutrino magnetohydrodynamics				
	publication_date   2017-12-15 12:34:31+00:00					
	source	SupportedSources.ARXIV				
	journal	None	authors			
	volume		title	Neutrino magnetohydrodynamics		
	doi		publication_date	2016-01-01 00:00:00		
	urls	• http://arxiv.org/pdf/1712.05640v1	source	SupportedSources.INTERNET_ARCHIVE		
		• http://arxiv.org/abs/1712.05640v1	journal	AIP Publishing		
		• http://arxiv.org/pdf/1712.05640v1	volume			
	id	id-2214159739917135704	doi	10.1063/pt.5.7232	4	
		A new neutrino magnetohydrodynamics (NMHD) model is formulated, where the effects of the charged weak current on the electron-ion magnetohydrodynamic fluid are taken into account. The model incorporates in a systematic way the role of the Fermi neutrino weak force in magnetized plasmas. A fast neutrino-driven short wavelengths instability associated with the magnetosonic wave is derived. Such an instability should play a central role in strongly magnetized plasma as occurs in supernovae, where dense neutrino beams also exist. In addition, in the case of nonlinear or high frequency waves, the neutrino coupling is shown to be responsible for breaking the frozen-in magnetic field lines condition even in infinite conductivity plasmas. Simplified and ideal NMHD assumptions were adopted and analyzed in detail.	urls	• https://web.archive.org/web/20200310161641/https://lume.ufrgs.br/bitstream/handle/10183/142518/000983192.pdf;jsessionid=1BB2086FAC1E750E978F4611DC594335? sequence=1	DUPLICATES 1	1043
			id	id1775480276882979042		
			abstract	Sheared-flow-driven ion-acoustic drift-wave instability and the formation of quadrupolar vortices in a nonuniform electron-positron-ion magnetoplasma Phys. Plasmas 11, 4341 (2004); 10.1063/1.1774164 Effect of nonlinear coupling of right and left circularly polarized Hall magnetohydrodynamic modes on filamentation processes in the solar wind A new neutrino magnetohydrodynamics (NMHD) model is formulated, where the effects of the charged weak current on the electron-ion magnetohydrodynamic fluid are taken into account. The model incorporates in a systematic way the role of the Fermi neutrino weak force in magnetized plasmas. A fast neutrino-driven short wavelengths instability associated with the magnetosonic wave is derived. Such an instability should play a central role in strongly magnetized plasma as occurs in supernovae, where dense neutrino beams also exist. In addition, in the case of nonlinear or high frequency waves, the neutrino coupling is shown to be responsible for breaking the frozen-in magnetic field lines condition even in infinite conductivity plasmas. Simplified and ideal NMHD assumptions were adopted and analyzed in detail. V C 2016 AIP Publishing LLC.		
			versions			
	versions					$\parallel \parallel$
						<u> </u>