

cases	doc_1		doc_2		decision	id
	authors	<ul style="list-style-type: none">Olindo ZanottiMichael DumbserArturo HidalgoDinshaw S. Balsara	authors	<ul style="list-style-type: none">O. ZanottiM. DumbserA. HidalgoD. Balsara	DUPLICATES	1072
	title	An ADER-WENO Finite Volume AMR code for Astrophysics	title	An ADER-WENO Finite Volume AMR code for Astrophysics		
	publication_date	2014-01-24 00:00:00	publication_date	2014-01-24 00:00:00		
	source	SupportedSources.OPENALEX	source	SupportedSources.SEMANTIC_SCHOLAR		
	journal	8th International Conference of Numerical Modeling of Space Plasma Flows (ASTRONUM 2013)	journal	arXiv: Instrumentation and Methods for Astrophysics		
	volume	488	volume			
	doi	None	doi			
	urls	<ul style="list-style-type: none">https://openalex.org/W2963139272	urls	<ul style="list-style-type: none">https://www.semanticscholar.org/paper/1a82e8a146daf1859243697d2bd454b7f800dde6		
	id	id-8587170392853109231	id	id-976187453671677469		
	abstract		abstract	A high order one-step ADER-WENO finite volume scheme with Adaptive Mesh Refinement (AMR) in multiple space dimensions is presented. A high order one-step time discretization is achieved using a local space-time discontinuous Galerkin predictor method, while a high order spatial accuracy is obtained through a WENO reconstruction. Thanks to the one-step nature of the underlying scheme, the resulting algorithm can be efficiently imported within an AMR framework on space-time adaptive meshes. We provide convincing evidence that the presented high order AMR scheme behaves better than traditional second order AMR methods. Tests are shown of the new scheme for nonlinear systems of hyperbolic conservation laws, including the classical Euler equations and the equations of ideal magnetohydrodynamics. The proposed scheme is likely to become a useful tool in several astrophysical scenarios.		
	versions		versions			