

# Enhanced Patient Satisfaction With Cellular-Enabled Glucose Meters in Pregnancy

Journal of Diabetes Science and Technology 2018, Vol. 12(5) 1086–1087 © 2018 Diabetes Technology Society Article reuse guidelines: sagepub.com/journals-permissions DOI: 10.1177/1932296818769101 journals.sagepub.com/home/dst

(\$)SAGE

Alyson R. Pierick, BS<sup>1</sup>, Sarah A. Wernimont, MD, PhD<sup>2</sup>, Emily A. Boevers, MA, BA<sup>1</sup>, Avery M. Whitis, BS<sup>1</sup>, Benjamin E. Deonovic, MS, PhD<sup>3</sup>, Diedre K. Fleener, RN, BSN<sup>2</sup>, Craig H. Syrop, MD, MHCDS<sup>2</sup>, and Janet I. Andrews, MD<sup>2</sup>

#### **Keywords**

pregnancy, blood glucose meter, patient satisfaction, telemedicine

Optimal diabetes management during pregnancy requires intensive glucose monitoring, medication adjustment and frequent communication with providers. Telemedicine strategies have been increasingly used to manage diabetes in pregnancy.<sup>2,3</sup> Unfortunately, many of these methods depend on reliable patient access to cellular services and wireless internet. Cellular-enabled blood glucose meters (BGM) provide real time glucose monitoring data and obviate the need for personal cell phones, data plans and internet access. These meters have been adopted outside of pregnancy to reduce patient and provider burdens of care. In addition, glucose values are uploaded in real time to a secure website for immediate evaluation by the medical care team. Patient satisfaction with cellular-enabled BGM compared to standard BGM has not been assessed in pregnancy. Our aim was to determine patient satisfaction with using a cellular-enabled BGM compared to standard BGM during pregnancy.

Women with diabetes (gestational, type 1 and type 2) were surveyed following delivery but prior to hospital discharge (n = 100). This project was reviewed by the University of Iowa Institutional Review Board and deemed exempt (IRB No 201509749). We used a modified version of the validated Oxford Maternity Diabetes Treatment Satisfaction Questionnaire. Survey response rate was 94%. This validated survey addresses patients' satisfaction with diabetes control, use of glucose monitoring equipment, life style fit, overall understanding of diabetes and communication and relationship with the care team. Survey responses on a 5-part Likert-type scale were analyzed using a Mann-Whitney two sample test.

Women who utilized a cellular-enabled BGM reported significantly more satisfaction with lifestyle fit (P = .001) and equipment (P = .022) compared to those using standard

BGM (Figure 1). Trends of improved perception of diabetes control, relationship and communication with the healthcare team, and overall experience with the cellular-enabled BGM were noted, though these differences did not reach statistical significance. Of note, the same multidisciplinary team consisting of physicians, advanced practitioners, and nurses made weekly contact with both groups of women for blood glucose review and possible medication adjustments.

Women describe significant burdens associated with the testing and reporting of blood glucose values in pregnancy. Efforts geared towards easing patient and provider burdens of care have increasingly employed telemedicine strategies to provide health services at a distance. Whether cellularenabled BGM can significantly improve maternal glucose control in pregnancy and perinatal outcomes remains to be determined. However, when up to 37% of blood glucose values are falsified in pregnancy, cellular-enabled BGM could play a critical role in providing more accurate and timely glucose values to care teams and provide a basis for better decision making and clinical management.

This work shows that cellular-enabled BGM increased patient satisfaction with several aspects of glucose management in pregnancy. Further studies are needed to assess

### **Corresponding Author:**

Janet I. Andrews, MD, Department of Obstetrics and Gynecology, University of Iowa Hospitals and Clinics, 31315 PFP, 200 Hawkins Dr, Iowa City, IA 52242, USA.
Email: janet-andrews@uiowa.edu

<sup>&</sup>lt;sup>1</sup>Carver College of Medicine, University of Iowa, Iowa City, IA, USA <sup>2</sup>Department of Obstetrics and Gynecology, University of Iowa Hospitals

Department of Obstetrics and Gynecology, University of Iowa Hospitals and Clinics, Iowa City, IA, USA

<sup>&</sup>lt;sup>3</sup>Department of Biostatistics, University of Iowa College of Public Health, Iowa City, IA, USA

Pierick et al 1087




Figure 1. Patient satisfaction survey results of 7 questions related to diabetes management during pregnancy. The P values are displayed for each question. Women who utilized a cellular-enabled blood glucose meter reported significantly higher satisfaction with lifestyle fit (P = .001) and equipment (P = .002) compared to those using standard blood glucose meters.

whether programs employing this technology have improved perinatal outcomes, efficiency, and provider satisfaction.

## **Abbreviation**

BGM, blood glucose meter.

# **Declaration of Conflicting Interests**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

# **Funding**

The author(s) received no financial support for the research, authorship, and/or publication of this article.

#### **ORCID iD**

Janet I Andrews https://orcid.org/0000-0003-0868-928X

## References

- 1. American Diabetes Association. Management of diabetes in pregnancy. *Diabetes Care*. 2017;40:S114-S119.
- Lau Y, Htun TP, Wong SN, Tam WS, Klainin-Yobas P. Efficacy of internet-based self-monitoring interventions on maternal and neonatal outcomes in perinatal diabetic women: a systematic review and meta-analysis. *J Med Internet Res*. 2016;18:e220.
- Ming WK, Mackillop LH, Farmer AJ, et al. Telemedicine technologies for diabetes in pregnancy: a systematic review and meta-analysis. *J Med Internet Res.* 2016;18:e290.
- 4. Hirst JE, Mackillop L, Loerup L, et al. Acceptability and user satisfaction of a smartphone-based, interactive blood glucose management system in women with gestational diabetes mellitus. *J Diabetes Sci Technol*. 2015;9:111-115.
- Kendrick JM, Wilson C, Elder RF, Smith CS. Reliability of reporting of self-monitoring of blood glucose in pregnant women. J Obstet Gynecol Neonatal Nurs. 2005;34:329-334.