




Ordinary XY calibration: $d = 3$, $\varepsilon/\pi = 1$, $N \geq 6$
Leading correction
REJECTED: $\chi^2/\mathrm{dof} = 330.0/8$
Leading + subleading correction
REJECTED: $\chi^2/\mathrm{dof} = 320.4/7$
N=6
N=8
N=12
N=16
$U = 1 - \langle m^4 \rangle / (2\langle m^2 \rangle^2)$
$(\mathrm{data} - \mathrm{fit})/\sigma$
β
Dots: paired-start measurements; vertical bars: 1 bootstrap SD. Dashed curves: saved joint fits.
Black rings: sampling-diagnostic flags. Residual shading: ±2 SD. These fits give no accepted critical coupling.


Ordinary XY calibration: $d = 3$, $\varepsilon/\pi = 1$, $N \geq 8$
Leading correction
REJECTED: $\chi^2/\mathrm{dof} = 290.0/5$
Leading + subleading correction
Unidentifiable: no unique fit to overlay
N=8
N=12
N=16
$U = 1 - \langle m^4 \rangle / (2\langle m^2 \rangle^2)$
Rank 4 for 5 parameters
$(\mathrm{data} - \mathrm{fit})/\sigma$
Residuals undefined
β
Dots: paired-start measurements; vertical bars: 1 bootstrap SD. Dashed curves: saved joint fits.
Black rings: sampling-diagnostic flags. Residual shading: ±2 SD. These fits give no accepted critical coupling.



Dots: paired-start measurements; vertical bars: 1 bootstrap SD. Dashed curves: saved joint fits.
Black rings: sampling-diagnostic flags. Residual shading: ±2 SD. These fits give no accepted critical coupling.


Ordinary XY calibration: $d = 4$, $\varepsilon/\pi = 1$, $N \geq 8$
Leading correction
REJECTED: $\chi^2/\mathrm{dof} = 1159.0/8$
Leading + subleading correction
REJECTED: $\chi^2/\mathrm{dof} = 1122.4/7$
N=8
N=10
N=12
N=16
$Q = \langle m^2 \rangle^2 / \langle m^4 \rangle$
(data − fit)/σ
β
Dots: paired-start measurements; vertical bars: 1 bootstrap SD. Dashed curves: saved joint fits.
Black rings: sampling-diagnostic flags. Residual shading: ±2 SD. These fits give no accepted critical coupling.