

# On Generalizations of the Nonwindowed Scattering Transform

Albert Chua<sup>a,\*</sup>, Matthew Hirn<sup>a,b,c</sup>, Anna Little<sup>d</sup>

<sup>a</sup>Department of Mathematics, Michigan State University, East Lansing, MI, 48824 USA

<sup>b</sup>Department of Computational Mathematics, Science & Engineering, Michigan State University, East Lansing, MI, 48824 USA

<sup>c</sup>Center for Quantum Computing, Science & Engineering Michigan State University, East Lansing, MI, 48824 USA

<sup>d</sup>Department of Mathematics and the Utah Center For Data Science, University of Utah, Salt Lake City, UT, 84112 USA

---

## Abstract

In this paper, we generalize finite depth wavelet scattering transforms, which we formulate as  $\mathbf{L}^q(\mathbb{R}^n)$  norms of a cascade of continuous wavelet transforms (or dyadic wavelet transforms) and contractive nonlinearities. We then provide norms for these operators, prove that these operators are well-defined, and are Lipschitz continuous to the action of  $C^2$  diffeomorphisms in specific cases. Lastly, we extend our results to formulate an operator invariant to the action of rotations  $R \in \mathrm{SO}(n)$  and an operator that is equivariant to the action of rotations of  $R \in \mathrm{SO}(n)$ .

*Keywords:* Wavelets, Wavelet Scattering Transform, Deformation Stability

---

## 1. Introduction

In recent years, convolutional neural networks have shown strong performance on various vision tasks like image classification [1, 2, 3, 4]. The main reason for this is that they are able to capture information at multiple scales through the use of convolutions and pooling. However, the exact method in which these networks use this information is not understood very well.

In [5], the author proposed a formulation for a simpler model for a convolutional neural network through the use of handcrafted filters, wavelets, and a series of cascading wavelet transforms. This model, called the scattering transform, and its extensions have shown success in vision tasks, quantum chemistry, manifold learning, and graph-related tasks [6, 7, 8, 9, 10].

We first provide a review of scattering transforms to motivate this paper. Let  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$  be a low pass filter ( $\hat{\phi}(0) \neq 0$ ),  $\psi : \mathbb{R}^n \rightarrow \mathbb{C}$  a suitable mother wavelet ( $\hat{\psi}(0) = 0$ ), and  $G^+$  be a set of “positive” rotations with determinant 1. Define a set of rotations and dilations by

$$\Lambda_J := \{\lambda = 2^j r : r \in G^+, j > -J\} \text{ if } J \neq \infty \quad (1)$$

and

$$\Lambda_\infty := \{2^j r : r \in G^+, j \in \mathbb{Z}\}. \quad (2)$$

Let  $\lambda = 2^j r \in \Lambda_J$ . Consider the operator

$$U[\lambda]f = \left| \int_{\mathbb{R}^n} f(u) 2^{-nj} \psi(2^{-j} r^{-1}(x - u)) du \right| \quad (3)$$

---

<sup>\*</sup>This work was partially supported by the National Science Foundation [A.C. and M.H. by grant DMS-1845856; A.L. and M.H. by grant DMS-1912906], the National Institutes of Health [M.H. by grant NIGMS-R01GM135929], and the Department of Energy [M.H. by grant DE-SC0021152].

<sup>\*</sup>Corresponding author

Email addresses: chuaalbe@msu.edu (Albert Chua), mhirn@msu.edu (Matthew Hirn), little@math.utah.edu (Anna Little)

For a tuple of rotations and dilations in  $\Lambda_J$ , define a path of length  $m$  as the tuple  $p := (\lambda_1, \dots, \lambda_m)$  and let  $\mathcal{P}_J$  be the set of all finite paths. The scattering propagator for  $f \in \mathbf{L}^2(\mathbb{R}^n)$  and  $p \in \mathcal{P}_J$  is

$$U[p]f := U[\lambda_m] \cdots U[\lambda_1]f, \quad (4)$$

which gathers high frequency information via a cascade of wavelet transforms and nonlinearities. The scattering operator is

$$\bar{S}f(p) = \frac{1}{\mu_p} \int_{\mathbb{R}^n} U[p]f(x) dx \quad (5)$$

with  $\mu_p := \int_{\mathbb{R}^n} U[p]\delta(x) dx$ . Additionally, to aggregate features similar to pooling, the author of [5] define the scattering operator for  $f \in \mathbf{L}^2(\mathbb{R}^n)$  and  $p \in \mathcal{P}_J$  as

$$S_J[p]f(x) = \int_{\mathbb{R}^n} U[p]f(u) 2^{-nJ} \phi(2^{-J}(x - u)) du. \quad (6)$$

Additionally, the windowed scattering transform is the set of functions

$$S_J[\mathcal{P}_J]f = \{S_J[p]f\}_{p \in \mathcal{P}_J}. \quad (7)$$

This operator is similar to a convolution neural network because along each path (analogous to each layer of a convolutional neural network) a convolution, a nonlinearity is applied, and feature aggregation occurs via the low pass filter. The scattering norm for any set of paths  $\Omega$  is

$$\|S_J[\Omega]f\|^2 = \sum_{p \in \Omega} \|S_J[p]f\|_2^2. \quad (8)$$

Under very stringent conditions on the mother wavelet, the author of [5] was able to prove an isometry property for the windowed scattering transform. However, the problem with the admissibility condition in [5] is that there are very few classes of wavelets that are admissible. The author of [5] mentions an analytic cubic spline Battle-Lemarié wavelet is admissible in one dimension, but provides no other examples. On a related note, [11] has shown that scattering coefficients have exponential decay for  $n = 1$  under relatively mild assumptions, but her proof only applies for  $n = 1$ , which makes the admissibility condition still necessary for  $n \geq 2$ . Additionally, to our knowledge, there are no examples in the literature of wavelets that satisfy the admissibility condition when  $n > 1$ .

The windowed scattering transform has three important properties that are helpful for certain machine learning tasks. The first two are the following:

1. The windowed scattering transform is a well-defined mapping on  $\mathbf{L}^2(\mathbb{R}^n)$  and nonexpansive. In particular, for all  $f, h \in \mathbf{L}^2(\mathbb{R}^n)$ ,

$$\|S_J[\mathcal{P}_J]f - S_J[\mathcal{P}_J]h\| \leq \|f - h\|_2. \quad (9)$$

2. Let the translation of a function be denoted as  $L_c f(u) = f(u - c)$ . For certain classes of wavelets, we have

$$\lim_{J \rightarrow \infty} \|S_J[\mathcal{P}_J]f - S_J[\mathcal{P}_J]L_c f\| = 0 \quad (10)$$

for all  $c \in \mathbb{R}^n$  and for all  $f \in \mathbf{L}^2(\mathbb{R}^n)$ . One can think of this as local translation invariance.

Finally, for the last property, the following definition was used in [5] for Lipschitz continuity to the action of  $C^2$  diffeomorphisms. Let  $\mathcal{H}$  be a Hilbert space,  $\tau \in C^2$ , and define the operator  $L_\tau f(x) = f(x - \tau(x))$ . A translation invariant operator  $\Phi$  is said to be Lipschitz continuous to the action of  $C^2$  diffeomorphisms if for any compact  $\Omega \subset \mathbb{R}^n$ , there exists  $C_\Omega$  such that for all  $f \in \mathbf{L}^2(\mathbb{R}^n)$  supported in  $\Omega$  and all  $\tau \in C^2(\mathbb{R}^n)$ , we have

$$\|\Phi(f) - \Phi(L_\tau f)\|_{\mathcal{H}} \leq C_\Omega \left( \|D\tau\|_\infty + \|D^2\tau\|_\infty \right) \|f\|_2. \quad (11)$$

The idea is that the difference in norm is proportional to the size of  $\|D\tau\|_\infty + \|D^2\tau\|_\infty$ , which indicates how much  $L_\tau$  deforms  $f$ . In particular, the author of [5] show that (11) holds for the windowed scattering transform.

The concept of stability to diffeomorphisms has become a major point of study after the publication of [5]. Based on the definition above, there has been a lot of interest in exploring the stability of various operators related to machine learning and data science. For example, [7, 12] extend the scattering transform and stability of the scattering transform to graphs and compact Riemannian manifolds, respectively; the authors in [13] loosen the restriction on the regularity of  $\tau$ . Other papers explore stability for different operators with desirable properties for machine learning [14, 15, 16, 17].

Although much work has appeared in recent years about operators similar to the scattering transform and about generalizations of the scattering transform, there are still some loose ends left in [5] that have not been explored yet. First, while the author of [5] does explore creating a norm for the nonwindowed scattering transform, he does not actually prove the norm is stable to diffeomorphisms. We consider a less stringent definition for stability to diffeomorphisms in the same spirit as the definition in [5] for this paper. Let  $V_1$  and  $V_2$  be normed vector spaces. Then we say that a translation invariant operator  $\Phi : V_1 \rightarrow V_2$  is said to be Lipschitz continuous to the action of  $C^2$  diffeomorphisms if for any compact  $\Omega \subset \mathbb{R}^n$ , there exists  $C_{\Omega,\tau}$  such that for all  $f \in V_1$  supported in  $\Omega$  and all  $\tau \in C^2(\mathbb{R}^n)$ , we have

$$\|\Phi(f) - \Phi(L_\tau f)\|_{V_2} \leq C_{\Omega,\tau} \|f\|_{V_1}, \quad (12)$$

where  $C_{\Omega,\tau} \rightarrow 0$  as  $\|D\tau\|_\infty + \|D^2\tau\|_\infty \rightarrow 0$ . Like with equation (11),  $\|\Phi(f) - \Phi(L_\tau f)\|_{V_2}$  depends on  $\|D\tau\|_\infty + \|D^2\tau\|_\infty$ .

Using this definition, we consider a slightly different problem than the author of [5] did for the nonwindowed scattering transform. The scattering transform introduced in [5] was a collection of  $L^1(\mathbb{R}^n)$  norms of various cascades of dyadic wavelet convolutions and modulus nonlinearities applied to a signal. Here, we extend the definition of the scattering transform to the continuous wavelet transform and for  $L^q(\mathbb{R}^n)$  norms with  $q \in [1, 2]$ . For a continuous dilation parameter  $\lambda \in \mathbb{R}_+$  we define the dilations of  $\psi$  as:

$$\forall \lambda \in \mathbb{R}_+, \quad \psi_\lambda(x) := \lambda^{-n/2} \psi(\lambda^{-1}x),$$

which preserves the  $L^2(\mathbb{R}^n)$  norm of  $\psi$ :

$$\|\psi_\lambda\|_2 = \|\psi\|_2, \quad \forall \lambda \in \mathbb{R}_+.$$

For the continuous wavelet transform, the one layer wavelet scattering transform with  $L^q(\mathbb{R}^n)$  norm is the function  $S_{\text{cont},q} : \mathbb{R}_+ \rightarrow \mathbb{R}$  defined as:

$$\forall \lambda \in \mathbb{R}_+, \quad S_{\text{cont},q}f(\lambda) := \|f * \psi_\lambda\|_q. \quad (13)$$

For a dyadic dilation parameter  $j \in \mathbb{Z}$  we define dilations of  $\psi$  as:

$$\forall j \in \mathbb{Z}, \quad \psi_j(x) = 2^{-nj} \psi(2^{-j}x),$$

which preserves the  $L^1(\mathbb{R}^n)$  norm of  $\psi$ :

$$\|\psi_j\|_1 = \|\psi\|_1, \quad \forall j \in \mathbb{Z}.$$

The one layer wavelet scattering transform for the dyadic wavelet transform is the function  $S_{\text{dyad},q}f : \mathbb{Z} \rightarrow \mathbb{R}$  defined as:

$$\forall j \in \mathbb{Z}, \quad S_{\text{dyad},q}f(j) := \|f * \psi_j\|_q. \quad (14)$$

More generally, the  $m$ -layer wavelet scattering transforms  $S_{\text{cont},q}^m f : \mathbb{R}_+^m \rightarrow \mathbb{R}$  and  $S_{\text{dyad},q}^m f : \mathbb{Z}^m \rightarrow \mathbb{R}$  are defined as

$$S_{\text{cont},q}^m f(\lambda_1, \dots, \lambda_m) := |||f * \psi_{\lambda_1} * \psi_{\lambda_2} * \dots * \psi_{\lambda_m}|||_q, \quad (15)$$

$$S_{\text{dyad},q}^m f(j_1, \dots, j_m) := |||f * \psi_{j_1} * \psi_{j_2} * \dots * \psi_{j_m}|||_q. \quad (16)$$

This is similar to working with a windowed scattering transform with a finite number of layers. However, our operator is different from the operator  $S_J$  in [5] because it does not contain the filter  $A_J$  to aggregate low frequency information, so the scale parameter in our formulation is not bounded above or below. Additionally, because the averaging filter is replaced  $L^q(\mathbb{R}^n)$  norms, our representation is fully translation invariant rather than translation invariant as  $J \rightarrow \infty$ .

As for the significance of using  $L^q(\mathbb{R}^n)$  norms to replace the averaging filter, there is one area with direct application: quantum energy regression tasks [8], where a representation that is similar to the rotation invariant representation in Section 6.2 has already been used for quantum energy regression.

Given a configuration of atoms, we would like to estimate the ground state energy of the configuration. Suppose we have a molecule with  $K$  atoms with nuclear charges  $z_k$  and nuclear positions  $p_k$  with  $k = 1, \dots, K$ . The state  $x$  of a molecule is given by

$$x = \{(p_k, z_k) \in \mathbb{R}^3 \times \mathbb{R} : k = 1 \dots, K\}, \quad (17)$$

Due to how we have defined our state, we would like our representation to have the following properties:

- **Permutation Invariance:** the energy should not depend on the index of the molecules.
- **Deformation Stability:** small deformations of the molecule should only lead to small changes in energy of the system.
- **Isometry Invariance:** the energy should be invariant to group actions such as translations, rotations, and other general isometries.
- **Multiscale Interactions:** molecules have many interactions terms, and these interaction terms depend on the pairwise distance between atoms (i.e. short range covalent bonds and longer range Van Der Waals interactions).

The rotation invariant version of our scattering transform in section 6 satisfies permutation invariance, deformation stability, and has multiscale interactions based on the proofs we’ve provided. We do not prove isometry invariance, but the operator is rotation and translation invariant.

Motivated by DFT theory, the paper [8] uses a dictionary of one and two layer scattering norms with  $q = 1$  and  $q = 2$  to get (at the time) state-of-the-art results for energy regression tasks for planar molecules. In particular, scattering operators with  $q = 1$  scaled with the number of atoms in the system and  $q = 2$  encoded pairwise interactions. The motivation for using  $1 < q < 2$  comes from [9, 10], which based on the Thomas–Fermi–Dirac–von Weizsäcker model [18], also use scattering norms with  $q = 4/3, 5/3$ . Later papers, like [9, 10], use a similar representation, involving spherical harmonics, for 3D quantum energy regression.

Generalizing to stochastic processes, one can also consider scattering moments [5, 19], which have similar desirable properties as the nonwindowed scattering transform. Applications include, but are not limited to, audio texture synthesis [19] and cosmology [20]. The main idea in all these applications is that the nonwindowed scattering transform has desirable mathematical properties and provides a small number of relevant descriptors for high dimensional, complicated data.

**Remark 1.** We can replace all the modulus operators with any contraction mapping (or use different contraction mappings in each layer) in the definition above, and all the proofs in the rest of this paper will still work. In particular, the modulus can be replaced with a complex version of the rectified linear unit (ReLU) nonlinearity,  $\max(0, \operatorname{Re}(a_i))_{i=1, \dots, n}$  for  $a \in \mathbb{C}^n$ , which is a popular choice for complex neural networks. Nonetheless, we will use the modulus operator throughout this paper without any loss of generality.

We provide a general roadmap for this paper. Section 2 will cover notation, basic properties about wavelets and the wavelet scattering operator, and harmonic analysis that will be necessary for the paper. In Section 3, we provide norms for an  $m$ -layer wavelet scattering transforms and prove that the operators are well defined mappings into specific spaces when  $1 \leq q \leq 2$ . For Section 4, we explore conditions

under which the  $m$ -layer scattering transform is stable to dilations, and we generalize our results to diffeomorphisms in Section 5. Lastly, in Section 6, we formulate two new translation invariant operators that are stable to diffeomorphisms. The first is rotation equivariant, and the second is rotation invariant. Our contributions include, but are not limited to, the following:

- We formulate an extension of the dyadic wavelet scattering operator for a finite, arbitrary number of layers with parameter  $q \in [1, 2]$  by applying  $\mathbf{L}^q(\mathbb{R}^n)$  norms instead of  $\mathbf{L}^1(\mathbb{R}^n)$  norms. Additionally, we formulate a wavelet scattering operator with  $q \in [1, 2]$  that uses a continuous scale parameter, like the continuous wavelet transform.
- We create a new finite depth scattering norm using dyadic and continuous scales in the case when  $q \in [1, 2]$ , and prove that the mappings are well defined and provide theoretical justification for a broader class of wavelets that make the scattering transform Lipschitz continuous to the action of  $C^2$  diffeomorphisms. However, the trade-off is that our stability bound depends on the number of layers.
- We provide a condition for norm equivalence in the case of  $q = 2$  that is less stringent.
- In the case of  $q \in (1, 2]$ , we prove that our norm is stable to diffeomorphisms  $\tau \in C^2(\mathbb{R}^n)$  provided that  $\|D\tau\|_\infty < \frac{1}{2n}$  and the wavelet and its first and second partial derivatives have sufficient decay. In the case of  $q = 1$ , we show stability to dilations; for  $m \geq 2$  this is done in a damped scale measure, which is necessary because the undamped  $q = 1$  scattering norm is infinite for essentially every  $f$ .
- We extend our formulation to include invariance or equivariance to the action of rotations  $R \in \text{SO}(n)$ .

## 2. Notation and Basic Properties

We start by providing basic notation that we will use in this paper and proceed to give basic definitions and properties that will be necessary for our results.

### 2.1. Function Spaces

Set  $\mathbb{R}_+$  to be the positive real numbers, i.e.  $\mathbb{R}_+ := (0, \infty)$ . The gradient of a function  $f : \mathbb{R}^n \rightarrow \mathbb{C}$  is given by  $\nabla f$ , the Jacobian of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is given by  $Df$ , and the Hessian is given by  $D^2f$ . For  $1 \leq q < \infty$ , the  $\mathbf{L}^q(\mathbb{R}^n)$  norm of a function  $f : \mathbb{R}^n \rightarrow \mathbb{C}$  is  $\|f\|_q := [\int_{\mathbb{R}^n} |f(x)|^q dx]^{1/q}$ . When  $q = \infty$ ,  $\|f\|_\infty := \text{ess sup } |f|$ . We will also use the notation,  $\|\Delta f\|_\infty = \sup_{x, y \in \mathbb{R}^n} |f(x) - f(y)|$ , which should not be mistaken for applying a Laplacian operator. Greek letters with a vector symbol, such as  $\vec{\alpha} = (\alpha_1, \dots, \alpha_n)$ , will be a multi-index of nonnegative integers; additionally, we write  $|\vec{\alpha}| = \alpha_1 + \dots + \alpha_n$ , and the usage will be clear from context. The operator  $D^{\vec{\alpha}}$  is a multi-index of derivatives:  $D^{\vec{\alpha}}f = \frac{\partial^{|\vec{\alpha}|}}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}} f$ . For integer  $s \geq 0$ , we define the function space  $\mathbf{H}^s(\mathbb{R}^n) = \{f \in \mathbf{L}^2(\mathbb{R}^n) : D^{\vec{\alpha}}f \in \mathbf{L}^2(\mathbb{R}^n) \text{ for } |\vec{\alpha}| \leq s\}$ .

The Fourier transform of a function  $f \in \mathbf{L}^1(\mathbb{R}^n)$  is the function  $\hat{f} \in \mathbf{L}^\infty(\mathbb{R}^n)$  defined as:

$$\forall \omega \in \mathbb{R}^n, \quad \hat{f}(\omega) := \int_{\mathbb{R}^n} f(x) e^{-ix \cdot \omega} dx.$$

The Hilbert transform of a function  $f \in \mathbf{L}^1(\mathbb{R})$  is denoted by  $Hf$  and is defined as:

$$Hf(x) := \lim_{\epsilon \rightarrow 0} \int_{|x-y|>\epsilon} \frac{f(y)}{x-y} dy.$$

The map  $H$  is a convolution operator in which  $f$  is convolved against the function  $1/x$ . We note that

$$H : \mathbf{L}^q(\mathbb{R}) \rightarrow \mathbf{L}^q(\mathbb{R}), \quad \forall 1 < q < \infty,$$

however the result is not true for  $q = 1$ , i.e., if  $f \in \mathbf{L}^1(\mathbb{R})$  it is not necessarily true that  $Hf \in \mathbf{L}^1(\mathbb{R})$ . We thus introduce the Hardy space. We denote the Hardy space as  $\mathbf{H}^1(\mathbb{R})$  and it consists of those functions  $f \in \mathbf{L}^1(\mathbb{R})$  such that  $Hf \in \mathbf{L}^1(\mathbb{R})$  as well. For  $f \in \mathbf{H}^1(\mathbb{R})$  the Hardy space norm is  $\|f\|_{\mathbf{H}^1(\mathbb{R})}$ , which we define as (see Corollary 2.4.7 of [21])

$$\|f\|_{\mathbf{H}^1(\mathbb{R})} := \|f\|_1 + \|Hf\|_1. \quad (18)$$

One can show that if  $f \in \mathbf{H}^1(\mathbb{R})$ , then  $f$  must necessarily have zero average. An important property of the Hilbert transform and convolution is the following:

$$H(f * g) = Hf * g = f * Hg, \quad f \in \mathbf{L}^p(\mathbb{R}), \quad g \in \mathbf{L}^q(\mathbb{R}), \quad 1 < \frac{1}{p} + \frac{1}{q}.$$

We have a similar definition for Hardy spaces when  $n \geq 2$ . For  $1 \leq j \leq n$ , define the  $j^{\text{th}}$  Riesz transform as

$$R_j f(x) = \lim_{\varepsilon \rightarrow 0} \int_{|x-y| > \varepsilon} \frac{x_j - y_j}{|x-y|^{n+1}} f(y) dy, \quad (19)$$

where  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$ . The Hardy space  $f \in \mathbf{H}^1(\mathbb{R}^n)$  consists of functions  $f$  such that  $f \in \mathbf{L}^1(\mathbb{R}^n)$  and  $R_j f \in \mathbf{L}^1(\mathbb{R}^n)$  for  $1 \leq j \leq n$  as well. For  $f \in \mathbf{H}^1(\mathbb{R}^n)$  the Hardy space norm is  $\|f\|_{\mathbf{H}^1(\mathbb{R}^n)}$ , which we define as (see Corollary 2.4.7 of [21])

$$\|f\|_{\mathbf{H}^1(\mathbb{R}^n)} := \|f\|_1 + \sum_{j=1}^n \|R_j f\|_1. \quad (20)$$

## 2.2. Wavelets

We let  $\psi \in \mathbf{L}^1(\mathbb{R}^n) \cap \mathbf{L}^2(\mathbb{R}^n)$  be a wavelet, which means it is a function that is localized in both space and frequency and has zero average, i.e.,

$$\int_{\mathbb{R}^n} \psi(x) dx = 0.$$

Assume  $f \in \mathbf{L}^2(\mathbb{R}^n)$ . The continuous wavelet transform  $\mathcal{W}f \in \mathbf{L}^2(\mathbb{R}^n \times \mathbb{R}_+)$  is defined as:

$$\forall (x, \lambda) \in \mathbb{R}^n \times \mathbb{R}_+, \quad \mathcal{W}f(x, \lambda) := f * \psi_\lambda(x).$$

Furthermore, if  $\psi$  satisfies the following admissibility condition

$$\int_0^\infty \frac{|\widehat{\psi}(\lambda\omega)|^2}{\lambda} d\lambda = \mathcal{C}_\psi, \quad \forall \omega \in \mathbb{R}^n \setminus \{0\}, \quad (21)$$

for some  $\mathcal{C}_\psi > 0$ , then we will say that  $\psi$  is a Littlewood-Paley wavelet for the continuous wavelet transform. If  $\psi$  satisfies (21), one can show that the norm  $\mathcal{W}f$  computed with a weighted measure  $(dx, d\lambda/\lambda^{n+1})$  on  $\mathbb{R}^n \times \mathbb{R}_+$  is well defined:

$$\|\mathcal{W}f\|_{\mathbf{L}^2(\mathbb{R}^n \times \mathbb{R}_+)}^2 := \int_0^\infty \int_{\mathbb{R}^n} |\mathcal{W}f(x, \lambda)|^2 dx \frac{d\lambda}{\lambda^{n+1}} = \int_0^\infty \int_{\mathbb{R}^n} |f * \psi_\lambda(x)|^2 dx \frac{d\lambda}{\lambda^{n+1}} = \int_0^\infty \|f * \psi_\lambda\|_2^2 \frac{d\lambda}{\lambda^{n+1}}.$$

We note, in fact, that one can show:

$$\|\mathcal{W}f\|_{\mathbf{L}^2(\mathbb{R}^n \times \mathbb{R}_+)}^2 = \beta \cdot \mathcal{C}_\psi \|f\|_2^2.$$

where

$$\beta = \begin{cases} 1/2 & \text{if } \psi \text{ is real valued} \\ 1 & \text{if } \psi \text{ is complex valued} \end{cases}. \quad (22)$$

For a function  $f \in \mathbf{L}^2(\mathbb{R}^n)$  we define the dyadic wavelet transform  $Wf \in \ell^2(\mathbf{L}^2(\mathbb{R}^n))$  as

$$Wf = (f * \psi_j)_{j \in \mathbb{Z}}.$$

If  $\psi$  satisfies

$$\sum_{j \in \mathbb{Z}} |\widehat{\psi}(2^j \omega)|^2 = \hat{C}_\psi, \quad \forall \omega \in \mathbb{R}^n \setminus \{0\}, \quad (23)$$

for some  $\hat{C}_\psi > 0$ , then we will say that  $\psi$  is a Littlewood-Paley wavelet for the dyadic wavelet transform. If  $\psi$  satisfies (23), one can show that the norm  $Wf$  given below is well defined:

$$\|Wf\|_{\ell^2(\mathbf{L}^2(\mathbb{R}^n))}^2 := \sum_{j \in \mathbb{Z}} \|f * \psi_j\|_2^2.$$

In fact, we have the following norm equivalence:

$$\|Wf\|_{\ell^2(\mathbf{L}^2(\mathbb{R}^n))}^2 = \beta \cdot \hat{C}_\psi \|f\|_2^2,$$

where  $\beta$  is defined in (22).

### 2.3. Operator Valued Spaces

Consider a Banach space  $\mathcal{B}$ . Suppose  $f : \mathbb{R}^n \rightarrow \mathcal{B}$  and  $x \rightarrow \|f(x)\|_{\mathcal{B}}$  is measurable in the Lebesgue sense. Define  $\mathbf{L}_{\mathcal{B}}^p(\mathbb{R}^n)$  for  $1 \leq p < \infty$  to be

$$\|f\|_{\mathbf{L}_{\mathcal{B}}^p(\mathbb{R}^n)}^p = \int_{\mathbb{R}^n} \|f(x)\|_{\mathcal{B}}^p dx.$$

Also, for  $1 \leq p < \infty$ , define

$$\|f\|_{\mathbf{L}_{\mathcal{B}}^{p,\infty}(\mathbb{R}^n)} = \sup_{\delta > 0} \delta \cdot m(\{x \in \mathbb{R}^n : \|f(x)\|_{\mathcal{B}} > \delta\})^{1/p}.$$

We also have the following relation:

$$\|f\|_{\mathbf{L}_{\mathcal{B}}^{p,\infty}(\mathbb{R}^n)} \leq \|f\|_{\mathbf{L}_{\mathcal{B}}^p(\mathbb{R}^n)}.$$

Note that for  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ,

$$\|f\|_{\mathbf{L}_{\mathbb{R}^n}^p(\mathbb{R}^n)}^p = \int_{\mathbb{R}^n} \|f(x)\|_{\mathbb{R}^n}^p dx = \int_{\mathbb{R}^n} |f(x)|^p dx = \|f\|_p^p.$$

## 3. Wavelet Scattering is a Bounded Operator

In this section we explore for which  $q > 0$  and  $m \geq 1$  the wavelet scattering transforms  $S_{\text{cont},q}^m f$  and  $S_{\text{dyad},q}^m f$  are well-defined as functions in some Banach space (i.e., have finite norm), and under what circumstances.

Let  $\psi$  be a wavelet. We assume that  $\psi$  has the following properties:

$$|\psi(x)| \leq A(1 + |x|)^{-n-\varepsilon} \quad (24)$$

$$\int_{\mathbb{R}^n} |\psi(x-y) - \psi(x)| dx \leq A|y|^{\varepsilon'}, \quad (25)$$

for some constants  $A, \varepsilon', \varepsilon > 0$  and for all  $y \neq 0$ .

Consider the Littlewood-Paley  $G$ -function

$$G_\psi(f)(x) = \left( \int_{(0,\infty)} |f * t^{-n} \psi(x/t)|^2 \frac{dt}{t} \right)^{1/2}. \quad (26)$$

Let  $\mathcal{B} = \mathbf{L}^2\left((0, \infty), \frac{dt}{t}\right)$ . We can rewrite this as a Bochner integral by considering the function  $K(x) = (t^{-n/2}\psi_t(x))_{t>0}$ . This is a mapping  $K : \mathbb{R}^n \rightarrow \mathcal{B}$  and the function  $x \rightarrow \|K(x)\|_{\mathcal{B}}$  is measurable. Also, if we let

$$\mathcal{T}(f)(x) = \left( \int_{\mathbb{R}^n} t^{-n/2} \psi_t(x-y) f(y) dy \right)_{t>0} = \left( (t^{-n/2} \psi_t * f)(x) \right)_{t>0},$$

we observe that

$$G_\psi(f)(x) = \|\mathcal{T}(f)(x)\|_{\mathcal{B}}$$

and

$$\|G_\psi(f)\|_p^p = \|\mathcal{T}(f)\|_{L_B^p(\mathbb{R}^n)}^p.$$

From Problem 6.1.4 of [22], the two properties above for the wavelet  $\psi$  imply that

$$\|K(x)\|_{\mathcal{B}} \leq \frac{c_n A}{|x|^n}, \quad (27)$$

and

$$\sup_{y \in \mathbb{R}^n \setminus \{0\}} \int_{|x| \geq 2|y|} \|K(x-y) - K(x)\|_{\mathcal{B}} dx \leq c'_n A, \quad (28)$$

where  $c_n$  and  $c'_n$  depend only on  $n$ ,  $\varepsilon$ , and  $\varepsilon'$ . We will omit the dependence on  $\varepsilon$  and  $\varepsilon'$  throughout the rest of this paper, and this will have no effect on any of our proofs.

**Remark 2.** For the rest of this paper, we will write  $G$  in place of  $G_\psi$  when referring to the  $G$ -function because the dependence on the mother wavelet is clear.

**Remark 3.** Note that (25) holds under the alternative condition

$$|\nabla \psi(x)| \leq A(1 + |x|)^{-n-1-\varepsilon'}. \quad (29)$$

This is a consequence of Mean Value Theorem.

We have the following result taken from Problem 6.1.4 of [22] and from Chapter V of [23].

**Lemma 1** ([22, 23]). *Assume that  $\psi$  is defined as above and satisfies (27) and (28). Then the operator  $G$  is bounded from  $\mathbf{L}^2(\mathbb{R}^n)$  to  $\mathbf{L}^2(\mathbb{R}^n)$ . Also, for  $p \in (1, \infty)$  and  $\mathcal{B} = \mathbf{L}^2(\mathbb{R}_+, dt/t)$ , we have*

$$\|\mathcal{T}f\|_{L_B^p(\mathbb{R}^n)} \leq C_n A \max(p, (p-1)^{-1}) \|f\|_{L^p(\mathbb{R}^n)}, \quad (30)$$

for some  $C_n$ . For all  $f \in \mathbf{L}^1(\mathbb{R}^n)$ , we also have

$$\|\mathcal{T}f\|_{L_B^{1,\infty}(\mathbb{R}^n)} \leq C'_n A \|f\|_{L^1(\mathbb{R}^n)} \quad (31)$$

and

$$\|\mathcal{T}f\|_{L_B^1(\mathbb{R}^n)} \leq C'_n A \|f\|_{H^1(\mathbb{R}^n)}, \quad (32)$$

for some  $C'_n$ .

**Remark 4.** We can also formulate similar bounds for the Littlewood-Paley  $g$  operator

$$g(f)(x) := \left[ \sum_{j \in \mathbb{Z}} |\psi_j * f(x)|^2 \right]^{1/2} \quad (33)$$

using similar arguments.

**Remark 5.** Let  $\psi$  be a wavelet that has properties (24) and (25). Then with the  $\mathbf{L}^2$  normalized dilations, the Littlewood-Paley  $G$ -function can be written as:

$$G(f)(x) = \left[ \int_0^\infty |f * \psi_\lambda(x)|^2 \frac{d\lambda}{\lambda^{n+1}} \right]^{1/2}. \quad (34)$$

Note that the  $\lambda$  measure for  $G(f)$  matches the measure in defining the norm of  $\mathcal{W}f$ .

### 3.1. The $\mathbf{L}^2(\mathbb{R}^n)$ Wavelet Scattering Transform

In this subsection we prove the  $\mathbf{L}^2(\mathbb{R}^n)$  scattering transforms are bounded operators. More specifically, we prove that  $S_{\text{cont},2}^m : \mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+^m)$ , where  $\mathbf{L}^2(\mathbb{R}_+^m)$  has the weighted measure defined by

$$\|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 := \int_0^\infty \cdots \int_0^\infty |S_{\text{cont},2}^m f(\lambda_1, \dots, \lambda_m)|^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}}$$

and we show that  $\|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \leq C \|f\|_{\mathbf{L}^2(\mathbb{R}^n)}$ . We also show that  $S_{\text{dyad},2}^m : \mathbf{L}^2(\mathbb{R}^n) \rightarrow \ell^2(\mathbb{Z}^m)$ , where

$$\|S_{\text{dyad},2}^m f\|_{\ell^2(\mathbb{Z}^m)}^2 := \sum_{j_m \in \mathbb{Z}} \cdots \sum_{j_1 \in \mathbb{Z}} |S_{\text{dyad},2}^m f(j_1, \dots, j_m)|^2.$$

**Proposition 2.** *For any wavelet satisfying (24) and (25), we have  $S_{\text{cont},2}^m : \mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+^m)$  and  $S_{\text{dyad},2}^m : \mathbf{L}^2(\mathbb{R}^n) \rightarrow \ell^2(\mathbb{Z}^m)$ .*

*Proof.* The proof of the dyadic case is essentially identical to the proof given below and is thus omitted. The case of  $m = 1$  follows by an application of Fubini's Theorem:

$$\begin{aligned} \|S_{\text{cont},2} f\|_{\mathbf{L}^2(\mathbb{R}_+)}^2 &= \int_0^\infty \|f * \psi_\lambda\|_2^2 \frac{d\lambda}{\lambda^{n+1}} \\ &= \int_0^\infty \int_{\mathbb{R}^n} |(f * \psi_\lambda)(x)|^2 dx \frac{d\lambda}{\lambda^{n+1}} \\ &= \int_{\mathbb{R}^n} |G(f)(x)|^2 dx \\ &\leq C \|f\|_2^2 \end{aligned}$$

by boundedness of the G-function. Now we proceed by using induction. Assume that we have  $\|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 \leq C_m \|f\|_2^2$ . Let  $\mathcal{W}_t f = f * \psi_t$ , define  $Mf = |f|$ , and  $U_\lambda = M\mathcal{W}_\lambda$  for notational brevity. Then notice that

$$\| |f * \psi_{\lambda_1}| * \psi_{\lambda_2} | * \cdots * \psi_{\lambda_m} | * \psi_{\lambda_{m+1}} \|_2^2 = \|\mathcal{W}_{\lambda_{m+1}} U_{\lambda_m} \cdots U_{\lambda_1} f\|_2^2.$$

Substituting yields

$$\begin{aligned} \|S_{\text{cont},2}^{m+1} f\|_{\mathbf{L}^2(\mathbb{R}_+^{m+1})}^2 &= \int_0^\infty \cdots \int_0^\infty \|\mathcal{W}_{\lambda_{m+1}} U_{\lambda_m} \cdots U_{\lambda_1} f\|_2^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_{m+1}}{\lambda_{m+1}^{n+1}} \\ &= \int_0^\infty \cdots \int_0^\infty \int_0^\infty \|(U_{\lambda_m} \cdots U_{\lambda_1} f) * \psi_{\lambda_{m+1}}\|_2^2 \frac{d\lambda_{m+1}}{\lambda_{m+1}^{n+1}} \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \\ &= \int_0^\infty \cdots \int_0^\infty \|U_{\lambda_m} \cdots U_{\lambda_1} f\|_{\mathbf{L}^2(\mathbb{R}_+)}^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \\ &\leq C \int_0^\infty \cdots \int_0^\infty \|U_{\lambda_m} \cdots U_{\lambda_1} f\|_2^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \\ &= C \int_0^\infty \cdots \int_0^\infty |S_{\text{cont},2}^m(\lambda_1, \dots, \lambda_m)|^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \\ &\leq C^{m+1} \|f\|_2^2, \end{aligned}$$

where we used the induction hypothesis in the last line. This completes the proof.  $\square$

**Proposition 3.** *Suppose  $\psi$  is a Littlewood-Paley wavelet satisfying (24) and (25). Then  $S_{\text{cont},2}^m f : \mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+^m)$  and specifically  $\|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 = C_\psi^m \|f\|_2^2$ . Also,  $S_{\text{dyad},2}^m : \mathbf{L}^2(\mathbb{R}^n) \rightarrow \ell^2(\mathbb{Z}^m)$  and  $\|S_{\text{dyad},2}^m f\|_{\ell^2(\mathbb{Z}^m)}^2 = \hat{C}_\psi^m \|f\|_2^2$ .*

*Proof.* We only provide the proof of the continuous case again. First consider the case  $m = 1$ . We have:

$$\begin{aligned}
\|S_{\text{cont},2}f\|_{\mathbf{L}^2(\mathbb{R}_+)}^2 &= \int_0^\infty \|f * \psi_\lambda\|_2^2 \frac{d\lambda}{\lambda^{n+1}} \\
&= \frac{1}{(2\pi)^n} \int_0^\infty \|\hat{f} \cdot \hat{\psi}_\lambda\|_2^2 \frac{d\lambda}{\lambda^{n+1}} \\
&= \frac{1}{(2\pi)^n} \int_0^\infty \left( \int_{\mathbb{R}^n} |\hat{f}(\omega)|^2 |\hat{\psi}_\lambda(\omega)|^2 d\omega \right) \frac{d\lambda}{\lambda^{n+1}} \\
&= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \left( \int_0^\infty |\hat{\psi}(\lambda\omega)|^2 \frac{d\lambda}{\lambda} \right) |\hat{f}(\omega)|^2 d\omega \\
&= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} (C_\psi |\hat{f}(\omega)|^2) d\omega \\
&= \frac{1}{(2\pi)^n} C_\psi \|\hat{f}\|_2^2 \\
&= C_\psi \|f\|_2^2.
\end{aligned}$$

Thus the claim holds for  $m = 1$ . Now assume that it holds through  $m$ . Then by the inductive hypothesis,

$$\|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 = \int_0^\infty \cdots \int_0^\infty \| |f * \psi_{\lambda_1}| * \psi_{\lambda_2} | * \cdots * \psi_{\lambda_m} \|_2^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} = C_\psi^m \|f\|_2^2.$$

Now consider the case of  $m + 1$ . Similar to the previous proposition, we have

$$\begin{aligned}
\|S_{\text{cont},2}^{m+1} f\|_{\mathbf{L}^2(\mathbb{R}_+^{m+1})}^2 &= \int_0^\infty \cdots \int_0^\infty \left( \int_0^\infty \| (U_{\lambda_m} \cdots U_{\lambda_1} f) * \psi_{\lambda_{m+1}} \|_2^2 \frac{d\lambda_{m+1}}{\lambda_{m+1}^{n+1}} \right) \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \\
&= C_\psi \int_0^\infty \cdots \int_0^\infty |S_{\text{cont},2}^m f(\lambda_1, \dots, \lambda_m)|^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \\
&= C_\psi \|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 \\
&= C_\psi^{m+1} \|f\|_2^2.
\end{aligned}$$

Thus, the claim is proven by induction.  $\square$

### 3.2. The $\mathbf{L}^1(\mathbb{R}^n)$ Wavelet Scattering Transform

Define the notation  $\mathcal{W}_t f = f * \psi_t$ ,  $Mf = |f|$ , and  $U_t = M\mathcal{W}_t$ . We now study the mapping properties of  $S_{\text{cont},1}^m$  on  $\mathbf{H}^1(\mathbb{R}^n)$ . We will see that  $S_{\text{cont},1} : \mathbf{H}^1(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+)$  for one layer, that for  $m \geq 2$  the norm below is infinite for essentially every  $f$  (Proposition 7), and that a damped measure on the scale variables restores boundedness for all  $m$  (Theorem 11). The norm for  $S_{\text{cont},1}^m f$  is:

$$\begin{aligned}
\|S_{\text{cont},1}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} &:= \left( \int_0^\infty \int_0^\infty \cdots \int_0^\infty |S_{\text{cont},1}^m f(\lambda_1, \lambda_2, \dots, \lambda_m)|^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \frac{d\lambda_2}{\lambda_2^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{1/2} \\
&= \left( \int_0^\infty \int_0^\infty \cdots \int_0^\infty \| (U_{\lambda_{m-1}} \cdots U_{\lambda_1} f) * \psi_{\lambda_m} \|_1^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \frac{d\lambda_2}{\lambda_2^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{1/2}.
\end{aligned}$$

An analogous result will also hold for the operator  $\mathbf{H}^1(\mathbb{R}^n) \rightarrow \ell^2(\mathbb{Z}_+^m)$  with norm

$$\|S_{\text{dyad},1}^m f\|_{\ell^2(\mathbb{Z}^m)} := \left( \sum_{j_m \in \mathbb{Z}} \cdots \sum_{j_1 \in \mathbb{Z}} |S_{\text{dyad},1}^m f(j_1, \dots, j_m)|^2 \right)^{1/2}.$$

Before we begin, we assume that  $n \geq 2$ , which will be clear from context; we will need an important multiplier property of the individual Riesz Transforms:

$$\widehat{R_j f}(\omega) = -i \frac{\omega_j}{|\omega|} \hat{f}(\omega). \quad (35)$$

Let  $\vec{\alpha} = (\alpha_1, \dots, \alpha_n)$  be a multi-index with  $n$ -elements, and let  $t = (t_1, \dots, t_n) \in \mathbb{R}^n$ . We say that  $\psi$  has  $k$  vanishing moments if for all  $|\vec{\alpha}| < k$ , we have

$$\int_{\mathbb{R}^n} (\prod_{i=1}^n t_i^{\alpha_i}) \psi(t) dt = 0. \quad (36)$$

The following lemmas will be necessary.

**Lemma 4** ([24]). *Suppose that  $\psi$  has  $N$  vanishing moments, let  $M > 1$  be an integer, let  $\vec{\alpha}$  be defined as before, and let  $\vec{\beta} = (\beta_1, \dots, \beta_n)$  be a multi-index. Assume that  $\psi$  satisfies the following properties:*

- $\psi \in \mathbf{H}^s(\mathbb{R}^n) \cap C(\mathbb{R}^n)$  for some  $s > M + \frac{n}{2}$ .
- There exists  $A > 0$  and  $\epsilon \in [0, 1)$  such that  $\psi$  satisfies

$$|D^{\vec{\alpha}} \psi| \leq A(1 + |x|)^{-n-N-|\vec{\alpha}|+\epsilon} \text{ for } 0 \leq |\vec{\alpha}| \leq M.$$

- For  $0 \leq |\vec{\alpha}| \leq M - 1$  and  $|\vec{\beta}| < N + |\vec{\alpha}|$ ,

$$\int_{\mathbb{R}^n} \prod_{i=1}^n t_i^{\beta_i} D^{\vec{\alpha}} \psi(t) dt = 0.$$

Then

$$|D^{\vec{\alpha}} R_i \psi(x)| = |R_i D^{\vec{\alpha}} \psi(x)| \leq A(1 + |x|)^{-n-N-|\vec{\alpha}|+\epsilon+\delta}$$

for some  $0 < \delta < 1 - \epsilon$  and  $D^{\vec{\alpha}} R_i \psi$  has vanishing moments up to degree  $N - 1 + |\vec{\alpha}|$ .

An immediate consequence is the following Lemma, which we will provide without proof.

**Lemma 5.** *Suppose that  $\psi$  satisfies the following conditions:*

- $\psi \in \mathbf{H}^s(\mathbb{R}^n) \cap C(\mathbb{R}^n)$  for some  $s > 2 + \frac{n}{2}$ .
- There exists  $A > 0$  and  $\epsilon \in [0, 1)$  such that  $\psi$  satisfies

$$|D^{\vec{\alpha}} \psi| \leq A(1 + |x|)^{-n-2-|\vec{\alpha}|+\epsilon} \text{ for } 0 \leq |\vec{\alpha}| \leq 3.$$

- For  $0 \leq |\vec{\alpha}| \leq 2$  and  $|\vec{\beta}| < 2 + |\vec{\alpha}|$ ,

$$\int_{\mathbb{R}^n} \prod_{i=1}^n t_i^{\beta_i} D^{\vec{\alpha}} \psi(t) dt = 0.$$

Then  $R_j \psi$  and all of its first and second partial derivatives have  $O((1 + |x|)^{-n-1+\eta})$  decay for some  $\eta \in (0, 1)$ .

The first implication to take note of is that  $R_j \psi$  is a wavelet with "good" decay of itself and all its first and second partial derivatives. Note that the strict decay on the partial derivatives is necessary for technical reasons in later proofs, but decay on all second partial derivatives can be relaxed for the following theorem.

**Theorem 6.** *Let  $\psi$  be a wavelet satisfying (24) and (25), and let  $S_{\text{cont},1}$  and  $S_{\text{dyad},1}$  be defined as above. Then for  $f \in \mathbf{H}^1(\mathbb{R}^n)$ , there exists a constant  $C_1$  such that*

$$\|S_{\text{cont},1} f\|_{\mathbf{L}^2(\mathbb{R}_+)} \leq C_1 \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}.$$

Additionally,

$$\|S_{\text{dyad},1} f\|_{\ell^2(\mathbb{Z})} \leq C_1 \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}.$$

*Proof.* We only provide a proof for the continuous case because the dyadic case follows by almost identical reasoning. Let  $f \in \mathbf{H}^1(\mathbb{R}^n)$ . By Minkowski's integral inequality ([25], Theorem 202), we have

$$\begin{aligned}
\|S_{\text{cont},1}f\|_{\mathbf{L}^2(\mathbb{R}_+)} &= \left( \int_0^\infty \|f * \psi_\lambda\|_1^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{1/2} \\
&= \left( \int_0^\infty \left( \int_{\mathbb{R}^n} |f * \psi_\lambda(x)| dx \right)^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{1/2} \\
&\leq \left( \int_{\mathbb{R}^n} \left( \int_0^\infty |f * \psi_\lambda(x)|^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{1/2} dx \right) \\
&= \int_{\mathbb{R}^n} G(f)(x) dx \\
&= \|G(f)\|_1 \\
&\leq C_1 \|f\|_{\mathbf{H}^1(\mathbb{R}^n)},
\end{aligned}$$

where in the last inequality we used Lemma 1.  $\square$

**Remark 6.** Notice that the proof above only uses (24) and (25). In particular, we do not need Lemmas 4 and 5 here; they will only be used in Corollary 15 and Corollary 18.

Unfortunately, the argument above does not extend to  $m \geq 2$ . Let  $V_0$  be the identity operator and let  $V_k = U_{\lambda_k} \cdots U_{\lambda_1}$  for  $k \geq 1$ , so that

$$S_{\text{cont},1}^m f(\lambda_1, \dots, \lambda_m) = \|\mathcal{W}_{\lambda_m} V_{m-1} f\|_1.$$

The input to the  $m$ -th layer is  $V_{m-1}f$ , which is nonnegative. Thus, if  $V_{m-1}f$  does not vanish identically, then

$$\int_{\mathbb{R}^n} V_{m-1}f(x) dx > 0,$$

and  $V_{m-1}f \notin \mathbf{H}^1(\mathbb{R}^n)$  because every function in  $\mathbf{H}^1(\mathbb{R}^n)$  has zero average. In other words, the wavelet only provides cancellation in the first layer, which is exactly what the step above needs. In fact, the following proposition shows that the norm is infinite when  $m \geq 2$ .

**Proposition 7.** Let  $\psi \not\equiv 0$ , let  $m \geq 2$ , and let  $f \in \mathbf{L}^1(\mathbb{R}^n)$ . Assume that  $V_{m-1}f \neq 0$  for all  $(\lambda_1, \dots, \lambda_{m-1})$  in a set of positive Lebesgue measure. Then

$$\|S_{\text{cont},1}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} = \infty.$$

Similarly, if  $V_{m-1}f \neq 0$  for some  $(j_1, \dots, j_{m-1}) \in \mathbb{Z}^{m-1}$ , then

$$\|S_{\text{dyad},1}^m f\|_{\ell^2(\mathbb{Z}^m)} = \infty.$$

*Proof.* We only provide a proof for the continuous case first, and then indicate the changes for the dyadic case.

Fix  $\lambda' = (\lambda_1, \dots, \lambda_{m-1})$  such that  $g := V_{m-1}f \neq 0$ . Since  $g \geq 0$  and  $g \in \mathbf{L}^1(\mathbb{R}^n)$ , the function  $\widehat{g}$  is continuous and

$$\widehat{g}(0) = \int_{\mathbb{R}^n} g(x) dx = \|g\|_1 > 0.$$

Choose  $\omega_0 \in \mathbb{R}^n$  such that  $\widehat{\psi}(\omega_0) \neq 0$ , and use continuity of  $\widehat{g}$  at the origin to choose  $\Lambda = \Lambda(\lambda') > 0$  such that

$$\left| \widehat{g}\left(\frac{\omega_0}{\lambda}\right) \right| \geq \frac{\|g\|_1}{2}, \quad \lambda \geq \Lambda.$$

Now notice that  $\widehat{\psi}_\lambda(\omega) = \lambda^{n/2} \widehat{\psi}(\lambda\omega)$ . Evaluating at  $\omega = \omega_0/\lambda$  and using  $\|h\|_1 \geq \|\widehat{h}\|_\infty$ , we get

$$\begin{aligned} S_{\text{cont},1}^m f(\lambda', \lambda) &= \|g * \psi_\lambda\|_1 \\ &\geq \left| \widehat{g * \psi_\lambda} \left( \frac{\omega_0}{\lambda} \right) \right| \\ &= \lambda^{n/2} \left| \widehat{g} \left( \frac{\omega_0}{\lambda} \right) \right| |\widehat{\psi}(\omega_0)| \\ &\geq \frac{\lambda^{n/2}}{2} \|g\|_1 |\widehat{\psi}(\omega_0)| \end{aligned}$$

for all  $\lambda \geq \Lambda$ . It follows that

$$\begin{aligned} \int_0^\infty |S_{\text{cont},1}^m f(\lambda', \lambda)|^2 \frac{d\lambda}{\lambda^{n+1}} &\geq \frac{\|g\|_1^2 |\widehat{\psi}(\omega_0)|^2}{4} \int_\Lambda^\infty \lambda^n \frac{d\lambda}{\lambda^{n+1}} \\ &= \frac{\|g\|_1^2 |\widehat{\psi}(\omega_0)|^2}{4} \int_\Lambda^\infty \frac{d\lambda}{\lambda} \\ &= \infty. \end{aligned}$$

Since this holds for all  $\lambda'$  in a set of positive measure, Tonelli's Theorem gives the first claim.

For the dyadic case, recall that  $\|\psi_j\|_1 = \|\psi\|_1$  and  $\widehat{\psi}_j(\omega) = \widehat{\psi}(2^j\omega)$ . A similar argument gives

$$\|g * \psi_j\|_1 \geq \left| \widehat{g} \left( 2^{-j}\omega_0 \right) \right| |\widehat{\psi}(\omega_0)| \geq \frac{\|g\|_1 |\widehat{\psi}(\omega_0)|}{2}$$

for all sufficiently large  $j$ , so the sum over  $j \in \mathbb{Z}$  diverges. This proves the desired claim.  $\square$

**Remark 7.** The hypothesis of Proposition 7 is mild. Suppose that  $f * \psi_{\lambda_1} \neq 0$  for  $\lambda_1$  in a set of positive measure. If  $V_{k-1}f \neq 0$  for some  $k \geq 2$ , then  $V_{k-1}f$  is nonnegative with positive integral, so the argument above shows that  $(V_{k-1}f) * \psi_{\lambda_k} \neq 0$  for all sufficiently large  $\lambda_k$ . Thus,  $V_k f \neq 0$  on a set of positive measure, and induction on  $k$  gives the hypothesis. In particular, the hypothesis holds for every  $f \neq 0$  when  $\widehat{\psi}$  does not vanish identically on any ray  $\{\lambda\omega : \lambda > 0\}$ .

The case of  $n = 1$  is a little trickier. We have the following multiplier property for the Hilbert Transform:

$$\widehat{H}f(\omega) = \begin{cases} +i\widehat{f}(\omega) & \omega < 0 \\ -i\widehat{f}(\omega) & \omega > 0 \end{cases} \quad (37)$$

Unfortunately, this yields less regularity for  $\widehat{H}f$  at the origin without additional assumptions. However, notice that the Hilbert transform commutes with dilations, so in particular:

$$H(\psi_\lambda) = H(\psi)_\lambda \quad \text{and} \quad H(\psi_j) = H(\psi)_j.$$

Using the calculation of  $\widehat{H}f$  in (37) we see that

$$H\psi = -i\psi, \quad \text{if } \psi \text{ is complex analytic.}$$

Thus, we have the following corollary.

**Corollary 8.** Let  $\psi$  be a complex analytic wavelet such that (24) and (25) hold. Then for every  $h \in \mathbf{L}^1(\mathbb{R})$  and every  $\lambda > 0$ , we have  $\mathcal{W}_\lambda h \in \mathbf{H}^1(\mathbb{R})$  with

$$\|\mathcal{W}_\lambda h\|_{\mathbf{H}^1(\mathbb{R})} = 2 \|\mathcal{W}_\lambda h\|_1.$$

Additionally, for  $f \in \mathbf{H}^1(\mathbb{R})$  there exists a constant  $C_1$  such that

$$\begin{aligned} \|S_{\text{cont},1}f\|_{\mathbf{L}^2(\mathbb{R}_+)} &\leq C_1 \|f\|_{\mathbf{H}^1(\mathbb{R})}, \\ \|S_{\text{dyad},1}f\|_{\ell^2(\mathbb{Z})} &\leq C_1 \|f\|_{\mathbf{H}^1(\mathbb{R})}. \end{aligned}$$

Proposition 7 tells us that we need a different measure on the scale variables when  $m \geq 2$ . To see which scales cause the problem, let  $g = V_{k-1}f$  and suppose that  $\lambda_k$  is much larger than  $\lambda_{k-1}$ . Then  $\psi_{\lambda_k}$  is nearly constant on the region where  $g$  is concentrated, so that

$$\begin{aligned}\|g * \psi_{\lambda_k}\|_1 &\approx \lambda_k^{n/2} \|\psi\|_1 \|g\|_1 \\ &= \lambda_k^{n/2} \|\psi\|_1 S_{\text{cont},1}^{k-1} f(\lambda_1, \dots, \lambda_{k-1}).\end{aligned}$$

In other words, these coarse scales only reproduce the coefficient from the previous layer and provide no new information about  $f$ . We therefore damp scales that are far apart.

**Definition 9.** For  $\varepsilon > 0$ , let  $\theta_\varepsilon := \sqrt{2/\varepsilon}$  and define

$$\begin{aligned}\omega_\varepsilon(s, t) &:= \left( \frac{\min(s, t)}{\max(s, t)} \right)^\varepsilon, \\ \Omega_\varepsilon^m(\lambda_1, \dots, \lambda_m) &:= \prod_{k=2}^m \omega_\varepsilon(\lambda_{k-1}, \lambda_k),\end{aligned}$$

with  $\Omega_\varepsilon^1 := 1$ . For a measurable function  $F$  on  $\mathbb{R}_+^m$ , the damped norm is

$$\|F\|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^m)}^2 := \int_0^\infty \cdots \int_0^\infty |F(\lambda_1, \dots, \lambda_m)|^2 \Omega_\varepsilon^m(\lambda_1, \dots, \lambda_m) \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}},$$

and for a function  $F$  on  $\mathbb{Z}^m$  we define

$$\|F\|_{\ell_\varepsilon^2(\mathbb{Z}^m)}^2 := \sum_{j_m \in \mathbb{Z}} \cdots \sum_{j_1 \in \mathbb{Z}} |F(j_1, \dots, j_m)|^2 \prod_{k=2}^m 2^{-\varepsilon|j_k - j_{k-1}|}.$$

When  $m = 1$ , these are the norms used in Theorem 6.

The weight only depends on ratios of consecutive scales, so it does not interfere with the dilation arguments in Section 4. The following lemma collects what we need about it.

**Lemma 10.** Let  $\varepsilon > 0$ . Then for every  $s > 0$  and every  $i \in \mathbb{Z}$ , we have

$$\begin{aligned}\int_0^\infty \omega_\varepsilon(s, t) \frac{dt}{t} &= \frac{2}{\varepsilon}, \\ \sum_{j \in \mathbb{Z}} 2^{-\varepsilon|j-i|} &= \frac{1 + 2^{-\varepsilon}}{1 - 2^{-\varepsilon}} =: \hat{\theta}_\varepsilon^2.\end{aligned}$$

Moreover, if  $1 \leq k \leq m$  and  $H \geq 0$  depends only on  $(\lambda_1, \dots, \lambda_k)$ , then

$$\left\| \left( \prod_{j=k+1}^m \lambda_j^{n/2} \right) H \right\|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^m)} = \theta_\varepsilon^{m-k} \|H\|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^k)},$$

and similarly  $\|H\|_{\ell_\varepsilon^2(\mathbb{Z}^m)} = \hat{\theta}_\varepsilon^{m-k} \|H\|_{\ell_\varepsilon^2(\mathbb{Z}^k)}$  for  $H \geq 0$  depending only on  $(j_1, \dots, j_k)$ .

*Proof.* For the first identity, we split the integral at  $t = s$  and compute

$$\begin{aligned}\int_0^\infty \omega_\varepsilon(s, t) \frac{dt}{t} &= \int_0^s \left( \frac{t}{s} \right)^\varepsilon \frac{dt}{t} + \int_s^\infty \left( \frac{s}{t} \right)^\varepsilon \frac{dt}{t} \\ &= \frac{1}{\varepsilon} + \frac{1}{\varepsilon}.\end{aligned}$$

The second identity is the sum of two geometric series. For the third, notice that

$$\lambda_j^n \frac{d\lambda_j}{\lambda_j^{n+1}} = \frac{d\lambda_j}{\lambda_j},$$

so integrating out  $\lambda_m, \lambda_{m-1}, \dots, \lambda_{k+1}$  in this order and using the first identity at each step produces a factor of  $2/\varepsilon$  per variable. Tonelli's Theorem justifies the interchange since the integrand is nonnegative. The dyadic case is identical, using  $\|\psi_j\|_1 = \|\psi\|_1$  in place of the factors  $\lambda_j^{n/2}$ .  $\square$

We can now state the boundedness result for all  $m$ .

**Theorem 11.** *Let  $\psi$  be a wavelet satisfying (24) and (25), let  $\varphi \in \mathbf{L}^1(\mathbb{R}^n)$ , let  $\varepsilon > 0$ , and let  $m \geq 2$ . Then for every  $f \in \mathbf{L}^1(\mathbb{R}^n)$ ,*

$$\| (V_{m-1}f) * \varphi_{\lambda_m} \|_1 \| \varphi \|_1 \theta_\varepsilon \| S_{\text{cont},1}^{m-1} f \|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^{m-1})}.$$

In particular, for every  $m \geq 1$  and every  $f \in \mathbf{H}^1(\mathbb{R}^n)$ ,

$$\| S_{\text{cont},1}^m f \|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^m)} \leq (\|\psi\|_1 \theta_\varepsilon)^{m-1} C_1 \|f\|_{\mathbf{H}^1(\mathbb{R}^n)},$$

where  $C_1$  is the constant from Theorem 6. The same statements hold for the dyadic transform, with  $\theta_\varepsilon$  replaced by  $\hat{\theta}_\varepsilon$ .

*Proof.* We only provide a proof for the continuous case because the dyadic case follows by almost identical reasoning. Since  $V_{m-1}f = |\mathcal{W}_{\lambda_{m-1}} V_{m-2}f|$ , Young's inequality and  $\|\varphi_{\lambda_m}\|_1 = \lambda_m^{n/2} \|\varphi\|_1$  give

$$\begin{aligned} \| (V_{m-1}f) * \varphi_{\lambda_m} \|_1 &\leq \lambda_m^{n/2} \|\varphi\|_1 \|V_{m-1}f\|_1 \\ &= \lambda_m^{n/2} \|\varphi\|_1 S_{\text{cont},1}^{m-1} f(\lambda_1, \dots, \lambda_{m-1}). \end{aligned}$$

The right hand side does not depend on  $\lambda_m$ , so the first claim follows from Lemma 10 with  $k = m - 1$ . Taking  $\varphi = \psi$  and iterating, we get

$$\begin{aligned} \| S_{\text{cont},1}^m f \|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^m)} &\leq \|\psi\|_1 \theta_\varepsilon \| S_{\text{cont},1}^{m-1} f \|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^{m-1})} \\ &\leq (\|\psi\|_1 \theta_\varepsilon)^{m-1} \| S_{\text{cont},1} f \|_{\mathbf{L}^2(\mathbb{R}_+)} , \end{aligned}$$

and Theorem 6 bounds the last quantity. This proves the desired claim.  $\square$

**Remark 8.** The case  $\varepsilon = 0$  gives an infinite left hand side when  $m \geq 2$  by Proposition 7, and the constant  $\theta_\varepsilon$  blows up as  $\varepsilon \rightarrow 0$ . Alternatively, one can restrict the scales by

$$R^{-1} \leq \frac{\lambda_k}{\lambda_{k-1}} \leq R, \quad k = 2, \dots, m,$$

instead of damping them, which gives the same bounds with  $\theta_\varepsilon^2$  replaced by  $2 \log R$ . This is closer to what is done in practice, where a bounded range of scales is used.

### 3.3. $\mathbf{L}^q(\mathbb{R}^n)$ Wavelet Scattering Transform

In this subsection, assume  $1 < q < 2$ . We prove that for  $m \in \mathbb{N}$ ,  $S_{\text{cont},q}^m : \mathbf{L}^q(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+^m)$ . The norm for  $S_{\text{cont},q}^m f$  is:

$$\begin{aligned} \| S_{\text{cont},q}^m f \|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q &:= \left( \int_0^\infty \int_0^\infty \cdots \int_0^\infty |S_{\text{cont},q}^m f(\lambda_1, \lambda_2, \dots, \lambda_m)|^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \frac{d\lambda_2}{\lambda_2^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{q/2} \\ &= \left( \int_0^\infty \int_0^\infty \cdots \int_0^\infty \left( \| (U_{\lambda_{m-1}} \cdots U_{\lambda_1} f) * \psi_{\lambda_m} \|_q \right)^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \frac{d\lambda_2}{\lambda_2^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{q/2}. \end{aligned}$$

There is also an analagous result for

$$\|S_{\text{dyad},q}^m f\|_{\ell^2(\mathbb{Z}^m)}^q := \left( \sum_{j_m \in \mathbb{Z}} \cdots \sum_{j_1 \in \mathbb{Z}} |S_{\text{dyad},q}^m f(j_1, j_2, \dots, j_m)|^2 \right)^{q/2}.$$

**Theorem 12.** *Let  $1 < q < 2$ . Also, let  $\psi$  be a wavelet that satisfies properties (24) and (25) and let  $S_{\text{cont},q}^m$  and  $S_{\text{dyad},q}^m$  be defined as above. Then there exists a universal constant  $C_m > 0$  such that  $\|S_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \leq C_m \|f\|_q^q$  for all  $f \in \mathbf{L}^q(\mathbb{R}^n)$ , and furthermore  $\|S_{\text{dyad},q}^m f\|_{\ell^2(\mathbb{Z}^m)}^q \leq C_m \|f\|_q^q$ .*

*Proof.* We proceed by induction and consider the case of  $m = 1$  first. Let  $f \in \mathbf{L}^q(\mathbb{R}^n)$ . For the continuous wavelet transform, we apply Minkowski's integral inequality:

$$\begin{aligned} \|S_{\text{cont},q} f\|_{\mathbf{L}^2(\mathbb{R}_+)}^q &= \left[ \int_0^\infty (\|f * \psi_\lambda\|_q)^2 \frac{d\lambda}{\lambda^{n+1}} \right]^{q/2} \\ &= \left[ \int_0^\infty \left( \int_{\mathbb{R}^n} |f * \psi_\lambda(x)|^q dx \right)^{2/q} \frac{d\lambda}{\lambda^{n+1}} \right]^{q/2} \\ &\leq \int_{\mathbb{R}^n} \left( \int_0^\infty |f * \psi_\lambda(x)|^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{q/2} dx \\ &= \|G(f)\|_q^q \\ &\leq C \|f\|_q^q. \end{aligned}$$

where in the last inequality we used Lemma 1.

Now, let us assume that

$$\|S_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \leq C^{m \cdot q} \|f\|_{\mathbf{L}^q(\mathbb{R}^n)}^q.$$

We apply Minkowski's Integral inequality [25] to swap and then bound:

$$\begin{aligned} \|S_{\text{cont},q}^{m+1} f\|_{\mathbf{L}^2(\mathbb{R}_+^{m+1})}^q &= \left[ \int_0^\infty \cdots \int_0^\infty \left( \|(U_{\lambda_m} \cdots U_{\lambda_1} f) * \psi_{\lambda_{m+1}}\|_q \right)^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_{m+1}}{\lambda_{m+1}^{n+1}} \right]^{q/2} \\ &= \left[ \int_0^\infty \cdots \int_0^\infty \left( \int_{\mathbb{R}^n} |(U_{\lambda_m} \cdots U_{\lambda_1} f) * \psi_{\lambda_{m+1}}(x)|^q dx \right)^{2/q} \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_{m+1}}{\lambda_{m+1}^{n+1}} \right]^{q/2} \\ &= \left[ \int_0^\infty \cdots \int_0^\infty \left[ \int_0^\infty \left( \int_{\mathbb{R}^n} |(U_{\lambda_m} \cdots U_{\lambda_1} f) * \psi_{\lambda_{m+1}}(x)|^q dx \right)^{2/q} \frac{d\lambda_{m+1}}{\lambda_{m+1}^{n+1}} \right]^{\frac{q}{2} \cdot \frac{2}{q}} \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right]^{q/2} \\ &\leq \left[ \int_0^\infty \cdots \int_0^\infty \left[ \int_{\mathbb{R}^n} \left( \int_0^\infty |(U_{\lambda_m} \cdots U_{\lambda_1} f) * \psi_{\lambda_{m+1}}(x)|^2 \frac{d\lambda_{m+1}}{\lambda_{m+1}^{n+1}} \right)^{q/2} dx \right]^{\frac{2}{q}} \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right]^{q/2} \\ &= \left[ \int_0^\infty \cdots \int_0^\infty \|G(U_{\lambda_m} \cdots U_{\lambda_1} f)\|_q^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right]^{q/2} \\ &\leq C^q \left[ \int_0^\infty \cdots \int_0^\infty \|(U_{\lambda_m} \cdots U_{\lambda_1} f)\|_q^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right]^{q/2} \\ &= C^q \|S_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \\ &\leq C^{(m+1)q} \|f\|_q^q. \end{aligned}$$

This proves the desired claim.  $\square$

#### 4. Stability to Dilations

We now consider dilations defined by  $\tau(x) = cx$  for some constant  $c$ , so that  $L_\tau f(x) = f((1-c)x)$ . We will start by proving a lemma that will be useful for our work.

**Lemma 13.** *Assume  $L_\tau$  is defined as above. Then*

$$L_\tau f * \psi_\lambda(x) = (1-c)^{-n/2} \left( f * \psi_{(1-c)\lambda} \right) ((1-c)x).$$

*Proof.* Notice that

$$L_\tau f * \psi_\lambda(x) = \int_{\mathbb{R}^n} f((1-c)y) \psi_\lambda(x-y) dy.$$

We make the substitution  $z = (1-c)y$ . Then it follows that

$$\begin{aligned} L_\tau f * \psi_\lambda(x) &= (1-c)^{-n} \int_{\mathbb{R}^n} f(z) \psi_\lambda(x - (1-c)^{-1}z) dz \\ &= (1-c)^{-n} \int_{\mathbb{R}^n} f(z) \lambda^{-n/2} \psi \left( \lambda^{-1}(x - (1-c)^{-1}z) \right) dz \\ &= (1-c)^{-n/2} \int_{\mathbb{R}^n} f(z) [(1-c)\lambda]^{-n/2} \psi \left( [(1-c)\lambda]^{-1} ((1-c)x - z) \right) dz \\ &= (1-c)^{-n/2} \int_{\mathbb{R}^n} f(z) \psi_{(1-c)\lambda} ((1-c)x - z) dz \\ &= (1-c)^{-n/2} f * \psi_{(1-c)\lambda} ((1-c)x) \\ &= (1-c)^{-n/2} L_\tau \left( f * \psi_{(1-c)\lambda} \right) (x). \end{aligned}$$

□

**Remark 9.** We also have

$$L_\tau \mathcal{W}_\lambda f(x) = (f * \psi_\lambda)(x(1-c)).$$

Before we begin the next Lemma, we explain the general idea behind our approach to explain the necessity of Lemma 14. Define

$$\Psi(x) = (1-c)^{-n/2} \psi_{(1-c)}(x) - \psi(x). \quad (38)$$

We want to prove that  $\Psi$  satisfies (24) and (25) with a linear dependence on  $c$  for future stability lemmas.

**Lemma 14.** *Suppose that  $\psi$  is a wavelet that satisfies the following three conditions:*

$$|\psi(x)| \leq \frac{A}{(1+|x|)^{n+1+\alpha}} \quad x \in \mathbb{R}^n, \quad (39)$$

$$|\nabla \psi(x)| \leq \frac{A}{(1+|x|)^{n+1+\beta}} \quad x \in \mathbb{R}^n, \quad (40)$$

$$\|D^2 \psi(x)\|_\infty \leq \frac{A}{(1+|x|)^{n+1+\kappa}} \quad x \in \mathbb{R}^n, \quad (41)$$

for  $\alpha, \beta, \kappa > 0$ . Consider

$$\Psi(x) = (1-c)^{-n/2} \psi_{(1-c)}(x) - \psi(x).$$

for  $|c| < \frac{1}{2^n}$ . Then  $\Psi$  is a wavelet satisfying (24) and (25).

*Proof.* Without loss of generality, assume  $\alpha < \beta < \kappa < 1$ . First, it's clear that  $\int_{\mathbb{R}^n} \Psi = 0$ . We now just need to verify properties (24) and (25). Assume  $c > 0$ . We can modify the proof accordingly if  $c < 0$ . Then

$$\begin{aligned} |\Psi(x)| &= \left| (1-c)^{-n/2} \psi_{(1-c)}(x) - \psi(x) \right| \\ &= (1-c)^{-n} \left| \psi\left(\frac{x}{(1-c)}\right) - (1-c)^n \psi(x) \right| \\ &\leq (1-c)^{-n} \left| \psi\left(\frac{x}{1-c}\right) - \psi\left(\frac{1-c}{1-c}x\right) \right| + (1-c)^{-n} \sum_{j=1}^n \binom{n}{j} c^j |\psi(x)|. \end{aligned}$$

Now use mean value theorem on the first term to choose a point  $z$  on the segment connecting  $\frac{x}{1-c}$  and  $x$  such that

$$\frac{c}{1-c} |[\nabla \psi(z)]^T x| = \left| \psi\left(\frac{x}{1-c}\right) - \psi\left(\frac{1-c}{1-c}x\right) \right|.$$

We now use Cauchy-Schwarz to bound the left side:

$$\frac{c}{1-c} |[\nabla \psi(z)]^T x| \leq \frac{c}{1-c} \frac{A|x|}{(1+|z|)^{n+1+\beta}}.$$

Since  $z$  lies on the segment connecting  $\frac{x}{1-c}$  and  $x$ , we see that for some  $t \in [0, 1]$ , we have

$$\begin{aligned} z &= (1-t)\frac{x}{1-c} + tx \\ &= \frac{1-t}{1-c}x + \frac{t-tc}{1-c}x \\ &= \frac{1-t+t-tc}{1-c}x \\ &= \frac{1-tc}{1-c}x. \end{aligned}$$

Thus,  $|z| \geq |x|$ . It now follows that

$$\frac{c}{1-c} \frac{A|x|}{(1+|z|)^{n+1+\beta}} \leq \frac{c}{1-c} \frac{A}{(1+|x|)^{n+\beta}}.$$

Finally, we get

$$\begin{aligned} |\Psi(x)| &\leq \frac{c}{(1-c)^{n+1}} \frac{A}{(1+|x|)^{n+\beta}} + \frac{\sum_{j=1}^n \binom{n}{j} c^j}{(1-c)^{n+1}} \frac{A}{(1+|x|)^{n+\alpha}} \\ &\leq 2A \left( \frac{2n}{2n-1} \right)^{-n-1} \frac{\sum_{j=1}^n \binom{n}{j} c^j}{(1+|x|)^{n+\alpha}} \\ &\leq \frac{A_n c}{(1+|x|)^{n+\alpha}} \end{aligned}$$

for some constant  $A_n$  since we assume  $\alpha < \beta$  and  $c < \frac{1}{2n}$ . Thus, (24) is satisfied.

We use a similar idea for proving (25) holds. Assume  $c > 0$  without loss of generality and further assume that  $|x| \geq 2|y|$ . By Mean Value Theorem, there exists  $z$  on the line segment connecting  $x$  and  $x-y$  such that

$$|\Psi(x-y) - \Psi(x)| = |\nabla \Psi(z)| |y|.$$

Like before, we notice that

$$\begin{aligned}
|\nabla \Psi(z)| &= \left| (1-c)^{-n/2} \nabla \psi_{(1-c)}(z) - \nabla \psi(z) \right| \\
&= \left| (1-c)^{-n-1} \nabla \psi\left(\frac{z}{1-c}\right) - \nabla \psi(z) \right| \\
&= (1-c)^{-n-1} \left| \nabla \psi\left(\frac{z}{1-c}\right) - (1-c)^{n+1} \nabla \psi(z) \right| \\
&\leq (1-c)^{-n-1} \left| \nabla \psi\left(\frac{z}{1-c}\right) - \nabla \psi\left(\frac{1-c}{1-c} z\right) \right| + (1-c)^{-n-1} \sum_{j=1}^{n+1} \binom{n+1}{j} c^j |\nabla \psi(z)|.
\end{aligned}$$

Let  $S$  be the set of points on the segment connecting  $\frac{z}{1-c}$  and  $z$ . By Mean Value Inequality, since  $S$  is closed and bounded, we have

$$\left| \nabla \psi\left(\frac{z}{1-c}\right) - \nabla \psi\left(\frac{1-c}{1-c} z\right) \right| \leq \frac{c}{1-c} \max_{w \in S} \|D^2 \psi(w)\|_{\infty} |z|.$$

The maximum for the quantity above is attained in  $S$ , so let us say the maximizer is  $w_1 = (1-t)\frac{z}{1-c} + tz$  for some  $t \in [0, 1]$ . Now use decay of the Hessian to bound the right side:

$$\frac{c}{1-c} \max_{w \in S} \|D^2 \psi(w)\|_{\infty} |z| \leq \frac{c}{1-c} \frac{A|z|}{(1+|w_1|)^{n+1+\kappa}}.$$

It follows that

$$\begin{aligned}
w_1 &= (1-t)\frac{z}{1-c} + tz \\
&= \frac{1-t}{1-c} z + \frac{t-tc}{1-c} z \\
&= \frac{1-t+t-tc}{1-c} z \\
&= \frac{1-tc}{1-c} z.
\end{aligned}$$

Thus,  $|w_1| \geq |z|$ . We conclude

$$\frac{c}{1-c} \frac{A|z|}{(1+|w_1|)^{n+1+\kappa}} \leq \frac{c}{1-c} \frac{A}{(1+|z|)^{n+\kappa}}.$$

For bounding  $|\nabla \Psi(z)|$ , we see

$$\begin{aligned}
|\nabla \Psi(z)| &\leq \frac{c}{(1-c)^{n+2}} \frac{A}{(1+|z|)^{n+\kappa}} + \frac{\sum_{j=1}^{n+1} \binom{n+1}{j} c^j}{(1-c)^{n+1}} \frac{A}{(1+|z|)^{n+1+\beta}} \\
&\leq A(1-c)^{-n-2} \frac{2 \sum_{j=1}^{n+1} \binom{n+1}{j} c^j}{(1+|z|)^{n+\kappa}} \\
&\leq \left( \frac{2n}{2n-1} \right)^{n+2} \frac{2A \sum_{j=1}^{n+1} \binom{n+1}{j} c^j}{(1+|z|)^{n+\kappa}}.
\end{aligned}$$

Going back to proving (25) holds for  $\Psi$ ,

$$|\Psi(x-y) - \Psi(x)| = |\nabla \Psi(z)| |y| \leq \left( \frac{2n}{2n-1} \right)^{n+2} \frac{2A \sum_{j=1}^{n+1} \binom{n+1}{j} c^j |y|}{(1+|z|)^{n+\kappa}}.$$

since the point  $z$  lies on the lines on a line segment connecting  $x - y$  and  $x$  with  $|x| \geq 2|y|$ , we can use an argument similar to above to conclude

$$|\Psi(x - y) - \Psi(x)| \leq 2^{n+1+\kappa} \left( \frac{2n}{2n-1} \right)^{n+2} \frac{A \sum_{j=1}^{n+1} \binom{n+1}{j} c^j}{(1 + |x|)^{n+\kappa}} |y|.$$

Now integrate to get

$$\begin{aligned} \int_{|x| \geq 2|y|} |\Psi(x - y) - \Psi(x)| dx &\leq 2^{n+1+\kappa} \left( \frac{2n}{2n-1} \right)^{n+2} A \sum_{j=1}^{n+1} \binom{n+1}{j} c^j |y| \int_{|x| \geq 2|y|} \frac{dx}{|x|^{n+\kappa}} \\ &= 2^{n+1+\kappa} \left( \frac{2n}{2n-1} \right)^{n+2} A I_n \sum_{j=1}^{n+1} \binom{n+1}{j} c^j |y|^{1-\kappa}, \end{aligned}$$

where  $I_n$  is some constant associated with the integration. Finally, we have a bound of

$$\int_{|x| \geq 2|y|} |\Psi(x - y) - \Psi(x)| dx \leq \tilde{A}_n c |y|^{1-\kappa}.$$

for some constant  $\tilde{A}_n$  only dependent on the dimension  $n$ . Thus, (25) holds with exponent  $1 - \kappa \in (0, 1)$ . Let  $\hat{A}_n = \max\{A_n, \tilde{A}_n\}$ . It follows that

$$\begin{aligned} |\Psi(x)| &\leq \frac{\hat{A}_n |c|}{(1 + |x|)^{n+\alpha}} \\ \int_{|x| \geq 2|y|} |\Psi(x - y) - \Psi(x)| dx &\leq \hat{A}_n |c| |y|^{1-\kappa}. \end{aligned}$$

□

It follows from Problem 6.1.2 in [22] that the bound in the G-function depends linearly on the constant  $A$  from Lemma 1 when proving  $\mathbf{L}^2(\mathbb{R}^n)$  boundedness. Thus, the following corollaries hold.

**Corollary 15.** Assume  $|c| < \frac{1}{2n}$ . For  $\psi$  satisfying the conditions of Lemma 14, when  $1 < p < \infty$ , there exist constants  $C_{n,p}$  and  $\hat{C}_{n,p}$  such that

$$\left\| \left( \int_0^\infty |f * \Psi_\lambda(x)|^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{1/2} \right\|_{\mathbf{L}^p(\mathbb{R}^n)} \leq |c| \cdot C_{n,p} \max\{p, (p-1)^{-1}\} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}$$

and

$$\left\| \left( \sum_{j \in \mathbb{Z}} |f * \Psi_j(x)|^2 \right)^{1/2} \right\|_{\mathbf{L}^p(\mathbb{R}^n)} \leq |c| \cdot \hat{C}_n \max\{p, (p-1)^{-1}\} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}.$$

Alternatively, if one of the following holds:

- $n = 1$ ,  $\psi$  is complex analytic and satisfies the conditions of Lemma 14,
- $n \geq 2$  and  $\psi$  satisfies the conditions of Lemma 5,

there exist constants  $H_n$  and  $\hat{H}_n$  such that

$$\left\| \left( \int_0^\infty |f * \Psi_\lambda(x)|^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{1/2} \right\|_{\mathbf{L}^1(\mathbb{R}^n)} \leq |c| \cdot H_n \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}$$

and

$$\left\| \left( \sum_{j \in \mathbb{Z}} |f * \Psi_j(x)|^2 \right)^{1/2} \right\|_{\mathbf{L}^1(\mathbb{R}^n)} \leq |c| \cdot \hat{H}_n \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}.$$

Now we can use the results above for our main dilation stability results.

**Theorem 16.** Suppose that  $\psi$  is a wavelet that satisfies the conditions of Lemma 14. Then there exists a constants  $K_{n,m}$  and  $\hat{K}_{n,m}$  only dependent on  $n$  and  $m$  such that

$$\|S_{\text{cont},2}^m f - S_{\text{cont},2}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \leq |c| \cdot K_{n,m} \|f\|_2$$

and

$$\|S_{\text{dyad},2}^m f - S_{\text{dyad},2}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m)} \leq |c| \cdot \hat{K}_{n,m} \|f\|_2$$

for any  $|c| < \frac{1}{2^n}$ .

*Proof.* Without loss of generality, assume  $c > 0$ . Let

$$\begin{aligned} \mathcal{W}_t f &= f * \psi_t \\ Mf &= |f| \\ U_t &= M\mathcal{W}_t \\ A_q f &= \left( \int_{\mathbb{R}^n} f^q(x) dx \right)^{1/q}. \end{aligned}$$

It follows that  $S_{\text{cont},2}^m = A_2 M \mathcal{W}_{\lambda_m} U_{\lambda_{m-1}} \cdots U_{\lambda_1}$ . We will also let  $V_{m-1} = U_{\lambda_{m-1}} \cdots U_{\lambda_1}$ , with  $V_0$  being the identity operator, and make a slight abuse of notation by denoting  $\mathcal{W}_{\lambda_m}$  as  $\mathcal{W}$ . First, we will add and subtract  $A_2 M L_\tau \mathcal{W} V_{m-1} f$  and apply triangle inequality:

$$\begin{aligned} \|S_{\text{cont},2}^m f - S_{\text{cont},2}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} &= \|A_2 M \mathcal{W} V_{m-1} f - A_2 M \mathcal{W} V_{m-1} L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \\ &\leq \|A_2 M \mathcal{W} V_{m-1} f - A_2 M L_\tau \mathcal{W} V_{m-1} f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \\ &\quad + \|A_2 M L_\tau \mathcal{W} V_{m-1} f - A_2 M \mathcal{W} V_{m-1} L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}. \end{aligned}$$

We'll start by bounding the first term. We see that  $g = \mathcal{W} V_{m-1} f \in \mathbf{L}^2(\mathbb{R}^n)$ . Thus

$$|A_2 M \mathcal{W} V_{m-1} f - A_2 M L_\tau \mathcal{W} V_{m-1} f| = \|\|g\|_2 - \|L_\tau g\|_2\|.$$

Now use a change of variables:

$$\|L_\tau g\|_2^2 = \int_{\mathbb{R}^n} |g((1-c)x)|^2 dx = (1-c)^{-n} \|g\|_2^2.$$

It then follows that

$$\|\|L_\tau g\|_2 - \|g\|_2\| = \|g\|_2 \left( \frac{1}{(1-c)^{n/2}} - 1 \right) \leq \|g\|_2 \left( \frac{1}{(1-c)^n} - 1 \right).$$

Taking the scattering norm yields

$$\begin{aligned} \|A_2 M \mathcal{W} V_{m-1} f - A_2 M L_\tau \mathcal{W} V_{m-1} f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 &\leq \left( \frac{1}{(1-c)^n} - 1 \right)^2 \|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 \\ &= \left( \frac{1 - (1-c)^n}{(1-c)^n} \right)^2 \|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 \\ &= \left( \frac{1}{(1-c)^n} \sum_{j=1}^n \binom{n}{j} c^j \right)^2 \|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 \\ &\leq \left[ \left( \frac{2n}{2n-1} \right)^n \sum_{j=1}^n \binom{n}{j} c^j \right]^2 \|S_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 \\ &\leq c^2 \cdot C_{m,n} \|f\|_2^2. \end{aligned}$$

For the second term, apply Minkowski's inequality for 2 norms:

$$\begin{aligned}
& \|A_2 M L_\tau \mathcal{W} V_{m-1} f - A_2 M \mathcal{W} V_{m-1} L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \\
&= \left( \int_0^\infty \cdots \int_0^\infty \|L_\tau \mathcal{W} V_{m-1} f\|_2 - \|\mathcal{W} L_\tau V_{m-1} f\|_2 \|^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{1/2} \\
&\leq \left( \int_0^\infty \cdots \int_0^\infty \|L_\tau \mathcal{W} V_{m-1} f - \mathcal{W} L_\tau V_{m-1} f\|_2^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{1/2} \\
&= \|A_2 M[\mathcal{W} V_{m-1}, L_\tau] f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}.
\end{aligned}$$

Now this is a commutator term, and we can now bound:

$$\begin{aligned}
\|A_2 M[\mathcal{W} V_{m-1}, L_\tau] f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 &= \int_0^\infty \cdots \int_0^\infty \|[\mathcal{W} V_{m-1}, L_\tau] f\|_2^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \\
&= \|[\mathcal{W} V_{m-1}, L_\tau] f\|_{\mathbf{L}^2(\mathbb{R}_+^m \times \mathbb{R}^n)}^2 \\
&\leq \|[\mathcal{W} V_{m-1}, L_\tau]\|_{\mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+^m \times \mathbb{R}^n)}^2 \|f\|_2^2.
\end{aligned}$$

We examine the commutator term more closely. Without a loss of generality, assume  $m \geq 2$ . By expanding it, we see that each term contains  $[\mathcal{W}, L_\tau]$ . It follows that

$$\begin{aligned}
\|[\mathcal{W} V_{m-1}, L_\tau]\|_{\mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+^m \times \mathbb{R}^n)} &\leq m \|\mathcal{W}\|_{\mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)}^{m-1} \|M\|_{\mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}^n)}^{m-1} \|[\mathcal{W}, L_\tau]\|_{\mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)} \\
&\leq C_m \|[\mathcal{W}, L_\tau]\|_{\mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)}.
\end{aligned}$$

Thus, once we bound this quantity appropriately, we will finish the proof. We start by writing

$$\|[\mathcal{W}, L_\tau] f\|_{\mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)}^2 = \int_0^\infty \|(L_\tau f) * \psi_\lambda - L_\tau(f * \psi_\lambda)\|_2^2 \frac{d\lambda}{\lambda^{n+1}}.$$

By substitution with  $z = (1 - c)x$  and Lemma 13,

$$\begin{aligned}
\|(L_\tau f) * \psi_\lambda - L_\tau(f * \psi_\lambda)\|_2^2 &= \int_{\mathbb{R}^n} |(L_\tau f * \psi_\lambda)(x) - L_\tau(f * \psi_\lambda)(x)|^2 dx \\
&= \int_{\mathbb{R}^n} \left| (1 - c)^{-n/2} (f * \psi_{(1-c)\lambda})((1 - c)x) - (f * \psi_\lambda)((1 - c)x) \right|^2 dx \\
&= (1 - c)^{-n} \int_{\mathbb{R}^n} \left| (1 - c)^{-n/2} (f * \psi_{(1-c)\lambda})(z) - (f * \psi_\lambda)(z) \right|^2 dz \\
&= (1 - c)^{-n} \int_{\mathbb{R}^n} \left| f * \left( (1 - c)^{-n/2} \psi_{(1-c)\lambda} - \psi_\lambda \right) \right|^2 dz \\
&= (1 - c)^{-n} \int_{\mathbb{R}^n} |(f * \Psi_\lambda)(z)|^2 dz, \\
&= (1 - c)^{-n} \|f * \Psi_\lambda\|_2^2.
\end{aligned}$$

Thus, we obtain

$$\begin{aligned}
\int_0^\infty \|(L_\tau f) * \psi_\lambda - L_\tau(f * \psi_\lambda)\|_2^2 \frac{d\lambda}{\lambda^{n+1}} &= (1 - c)^{-n} \int_0^\infty \|f * \Psi_\lambda\|_2^2 \frac{d\lambda}{\lambda^{n+1}} \\
&= (1 - c)^{-n} \int_{\mathbb{R}^n} \int_0^\infty |f * \Psi_\lambda(x)|^2 \frac{d\lambda}{\lambda^{n+1}} dx \\
&= (1 - c)^{-n} \left\| \left( \int_0^\infty |f * \Psi_\lambda(x)|^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{1/2} \right\|_2^2 \\
&\leq c^2 \cdot \left( \frac{2n}{2n - 1} \right)^n C_{n,p} \|f\|_2^2.
\end{aligned}$$

It follows that

$$\|S_{\text{cont},2}^m f - S_{\text{cont},2}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \leq |c| \cdot K_{n,m} \|f\|_2$$

for any  $|c| < \frac{1}{2n}$ . □

As is customary at this point, we have the following corollaries. We start with the case where  $1 < q < 2$ .

**Corollary 17.** Assume  $|c| < \frac{1}{2n}$ . For  $q \in (1, 2)$ , there exists constants  $K_{n,m,q}$  and  $\hat{K}_{n,m,q}$  such that

$$\|S_{\text{cont},q}^m f - S_{\text{cont},q}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \leq |c|^q \cdot K_{n,m,q} \|f\|_q^q$$

and

$$\|S_{\text{dyad},q}^m f - S_{\text{dyad},q}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m)}^q \leq |c|^q \cdot \hat{K}_{n,m,q} \|f\|_q^q.$$

*Proof.* Without loss of generality again, assume  $c > 0$ . First, we will add and subtract  $A_q M L_\tau \mathcal{W} V_{m-1} f$  and apply triangle inequality:

$$\begin{aligned} \|S_{\text{cont},q}^m f - S_{\text{cont},q}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} &= \|A_q M \mathcal{W} V_{m-1} f - A_q M \mathcal{W} V_{m-1} L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \\ &\leq \|A_q M \mathcal{W} V_{m-1} f - A_q M L_\tau \mathcal{W} V_{m-1} f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \\ &\quad + \|A_q M L_\tau \mathcal{W} V_{m-1} f - A_q M \mathcal{W} V_{m-1} L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}. \end{aligned}$$

We'll start by bounding the first term again. Define  $g = \mathcal{W} V_{m-1} f \in \mathbf{L}^q(\mathbb{R}^n)$ . and we have

$$|A_q M \mathcal{W} V_{m-1} f - A_q M L_\tau \mathcal{W} V_{m-1} f| = \|\|g\|_q - \|L_\tau g\|_q\|.$$

By change of variables,

$$\|\|g\|_q - \|L_\tau g\|_q\| = \|g\|_q \left( \frac{1}{(1-c)^{n/q}} - 1 \right) \leq \|g\|_q \left( \frac{1}{(1-c)^n} - 1 \right).$$

Again, we have

$$\begin{aligned} \|A_q M \mathcal{W} V_{m-1} f - A_q M L_\tau \mathcal{W} V_{m-1} f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q &\leq \left( \frac{1}{(1-c)^{n/q}} - 1 \right)^q \|S_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \\ &\leq \left( \frac{1}{(1-c)^n} - 1 \right)^q \|S_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \\ &= \left[ \frac{1 - (1-c)^n}{(1-c)^n} \right]^q \|S_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \\ &= \left[ \frac{1}{(1-c)^n} \sum_{j=1}^n \binom{n}{j} c^j \right]^q \|S_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \\ &\leq \left[ \left( \frac{2n}{2n-1} \right)^n \sum_{j=1}^n \binom{n}{j} c^j \right]^q \|S_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \\ &\leq |c|^q \cdot C_{m,n} \|f\|_q^q. \end{aligned}$$

For the second term, apply Minkoski's inequality for  $q$  norms:

$$\begin{aligned} &\|A_q M L_\tau \mathcal{W} V_{m-1} f - A_q M \mathcal{W} V_{m-1} L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \\ &= \left( \int_0^\infty \cdots \int_0^\infty \left| \|L_\tau \mathcal{W} V_{m-1} f\|_q - \|\mathcal{W} L_\tau V_{m-1} f\|_q \right|^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{1/2} \\ &\leq \left( \int_0^\infty \cdots \int_0^\infty \|L_\tau \mathcal{W} V_{m-1} f - \mathcal{W} L_\tau V_{m-1} f\|_q^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{1/2} \\ &= \|A_q M[\mathcal{W} V_{m-1}, L_\tau] f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}. \end{aligned}$$

Via a similar reduction technique for Theorem 16, we can reduce to a commutator bound  $\|A_q M[\mathcal{W}, L_\tau]f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}$ . Additionally, we have

$$\|(L_\tau f) * \psi_\lambda - L_\tau(f * \psi_\lambda)\|_q^q = (1-c)^{-n} \|f * \Psi_\lambda\|_q^q.$$

Thus,

$$\begin{aligned} \|A_q M[\mathcal{W}, L_\tau]f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q &= \left( \int_0^\infty \|(L_\tau f) * \psi_\lambda - L_\tau(f * \psi_\lambda)\|_q^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{q/2} \\ &= (1-c)^{-n} \left( \int_0^\infty \|f * \Psi_\lambda\|_q^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{q/2} \\ &\leq (1-c)^{-n} \left\| \left( \int_0^\infty |f * \Psi_\lambda(x)|^2 \frac{d\lambda}{\lambda^{n+1}} \right)^{1/2} \right\|_q^q \\ &\leq |c|^q \cdot \tilde{C}_n \|f\|_q^q. \end{aligned}$$

It follows that

$$\|S_{\text{cont},q}^m f - S_{\text{cont},q}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \leq |c|^q \cdot K_{n,m} \|f\|_q^q$$

for any  $|c| < \frac{1}{2n}$ . □

Additionally, we have the following result for the case  $q = 1$ . By Proposition 7, for  $m \geq 2$  the statement must be formulated in the damped norm of Definition 9; for  $m = 1$  that norm is the norm used in the previous corollaries.

**Corollary 18.** Assume  $|c| < \frac{1}{2n}$  and let  $\varepsilon > 0$ . Suppose one of the following holds:

- $n = 1$ ,  $\psi$  is complex analytic and satisfies the conditions of Lemma 14,
- $n \geq 2$  and  $\psi$  satisfies the conditions of Lemma 5,

then there exist constants  $K_{H,m}$  and  $\hat{K}_{H,m}$ , depending only on  $n, m, \varepsilon$  and  $\psi$ , such that

$$\|S_{\text{cont},1}^m f - S_{\text{cont},1}^m L_\tau f\|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^n)} \leq |c| \cdot K_{H,m} \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}$$

and

$$\|S_{\text{dyad},1}^m f - S_{\text{dyad},1}^m L_\tau f\|_{\ell_\varepsilon^2(\mathbb{Z}^m)} \leq |c| \cdot \hat{K}_{H,m} \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}.$$

*Proof.* We only provide a proof for the continuous case because the dyadic case follows by almost identical reasoning, with  $\theta_\varepsilon$  replaced by  $\hat{\theta}_\varepsilon$ . Throughout,  $C$  denotes a constant depending only on  $n, m, \varepsilon$ , and  $\psi$ , which may change from line to line.

Define

$$\begin{aligned} \psi^c &:= (1-c)^{-n/2} \psi_{(1-c)}, \\ \mathcal{W}_\lambda^c h &:= h * \psi_\lambda^c, \\ U_\lambda^c &:= M \mathcal{W}_\lambda^c, \\ V_k^c &:= U_{\lambda_k}^c \cdots U_{\lambda_1}^c, \end{aligned}$$

with  $V_0^c$  the identity operator, and let  $S_{\text{cont},1}^{m,c} f(\lambda) := \left\| \mathcal{W}_{\lambda_m}^c V_{m-1}^c f \right\|_1$  denote the scattering transform built from  $\psi^c$ . By the definition of  $\Psi$  in Lemma 14, we have

$$\psi^c = \psi + \Psi, \quad \psi_\lambda^c = \psi_\lambda + \Psi_\lambda, \quad \|\psi^c\|_1 = \|\psi\|_1.$$

We start with the scale covariance. By Lemma 13,

$$\begin{aligned}\mathcal{W}_\lambda L_\tau h &= (1-c)^{-n/2} L_\tau \left( h * \psi_{(1-c)\lambda} \right) \\ &= L_\tau \mathcal{W}_\lambda^c h,\end{aligned}$$

and since  $|L_\tau h| = L_\tau |h|$ , we also get  $U_\lambda L_\tau = L_\tau U_\lambda^c$ . Induction on  $k$  now gives  $V_k L_\tau = L_\tau V_k^c$ , and using  $\|L_\tau F\|_1 = (1-c)^{-n} \|F\|_1$ , it follows that

$$\begin{aligned}S_{\text{cont},1}^m L_\tau f &= \|\mathcal{W}_{\lambda_m} V_{m-1} L_\tau f\|_1 \\ &= \|L_\tau (\mathcal{W}_{\lambda_m}^c V_{m-1}^c f)\|_1 \\ &= (1-c)^{-n} S_{\text{cont},1}^{m,c} f.\end{aligned}$$

Thus, we can write

$$S_{\text{cont},1}^m f - S_{\text{cont},1}^m L_\tau f = \left( S_{\text{cont},1}^m f - S_{\text{cont},1}^{m,c} f \right) + (1 - (1-c)^{-n}) S_{\text{cont},1}^{m,c} f,$$

and we bound each term separately.

We'll start by bounding the second term. By the Mean Value Theorem and  $|c| < \frac{1}{2n}$ , we have

$$|1 - (1-c)^{-n}| \leq n \left( 1 - \frac{1}{2n} \right)^{-n-1} |c| \leq 4n|c|.$$

Notice that  $\psi^c = \psi + \Psi$  is a wavelet satisfying (24) and (25), and that  $G_{\psi^c}(f) \leq G(f) + G_\Psi(f)$  pointwise. Since the bound in Lemma 1 depends linearly on the constant  $A$ , Theorem 6 and Corollary 15 give

$$\|G_{\psi^c}(f)\|_1 \leq (C_1 + |c|H_n) \|f\|_{\mathbf{H}^1(\mathbb{R}^n)},$$

so Theorem 11, applied with  $\psi$  replaced by  $\psi^c$ , yields

$$\begin{aligned}\|S_{\text{cont},1}^{m,c} f\|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^m)} &\leq (\|\psi\|_1 \theta_\varepsilon)^{m-1} (C_1 + |c|H_n) \|f\|_{\mathbf{H}^1(\mathbb{R}^n)} \\ &\leq C \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}.\end{aligned}$$

For the first term, we telescope over the layers. Let

$$E_k(\lambda_1, \dots, \lambda_k) := \left\| \mathcal{W}_{\lambda_k} V_{k-1} f - \mathcal{W}_{\lambda_k}^c V_{k-1}^c f \right\|_1,$$

and notice that the reverse triangle inequality gives  $|S_{\text{cont},1}^m f - S_{\text{cont},1}^{m,c} f| \leq E_m$ . Since  $\psi_{\lambda_k}^c = \psi_{\lambda_k} + \Psi_{\lambda_k}$ , we can write

$$\mathcal{W}_{\lambda_k} V_{k-1} f - \mathcal{W}_{\lambda_k}^c V_{k-1}^c f = \mathcal{W}_{\lambda_k} (V_{k-1} f - V_{k-1}^c f) - (V_{k-1}^c f) * \Psi_{\lambda_k}.$$

When  $k = 1$ , the first term vanishes because  $V_0 = V_0^c$  is the identity operator. When  $k \geq 2$ , the modulus is nonexpansive on  $\mathbf{L}^1(\mathbb{R}^n)$ , so that

$$\begin{aligned}\|V_{k-1} f - V_{k-1}^c f\|_1 &= \left\| \left| \mathcal{W}_{\lambda_{k-1}} V_{k-2} f \right| - \left| \mathcal{W}_{\lambda_{k-1}}^c V_{k-2}^c f \right| \right\|_1 \\ &\leq E_{k-1}.\end{aligned}$$

Now use Young's inequality and  $\|\psi_{\lambda_k}\|_1 = \lambda_k^{n/2} \|\psi\|_1$  to get

$$E_k \leq \lambda_k^{n/2} \|\psi\|_1 E_{k-1} + \|(V_{k-1}^c f) * \Psi_{\lambda_k}\|_1,$$

and iterating this bound yields

$$\left| S_{\text{cont},1}^m f - S_{\text{cont},1}^{m,c} f \right| \leq E_m \leq \sum_{k=1}^m \|\psi\|_1^{m-k} \left( \prod_{j=k+1}^m \lambda_j^{n/2} \right) \|(V_{k-1}^c f) * \Psi_{\lambda_k}\|_1.$$

Finally, we take norms. Minkowski's inequality and Lemma 10 give

$$\left\| S_{\text{cont},1}^m f - S_{\text{cont},1}^{m,c} f \right\|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^m)} \leq \sum_{k=1}^m \|\psi\|_1^{m-k} \theta_\varepsilon^{m-k} \left\| \|(V_{k-1}^c f) * \Psi_{\lambda_k}\|_1 \right\|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^k)}.$$

When  $k = 1$ , the inner quantity is  $\|f * \Psi_{\lambda_1}\|_1$ , so Minkowski's integral inequality and Corollary 15 give

$$\begin{aligned} \left\| \|f * \Psi_{\lambda_1}\|_1 \right\|_{\mathbf{L}^2(\mathbb{R}_+)} &\leq \|G_\Psi(f)\|_1 \\ &\leq |c| H_n \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}. \end{aligned}$$

When  $k \geq 2$ , Theorem 11 applied with  $\psi$  replaced by  $\psi^c$  and with  $\varphi = \Psi$ , together with the bound  $\|\Psi\|_1 \leq |c| \cdot C$  obtained by integrating (24) for  $\Psi$ , gives

$$\begin{aligned} \left\| \|(V_{k-1}^c f) * \Psi_{\lambda_k}\|_1 \right\|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^k)} &\leq \|\Psi\|_1 \theta_\varepsilon (\|\psi\|_1 \theta_\varepsilon)^{k-2} (C_1 + |c| H_n) \|f\|_{\mathbf{H}^1(\mathbb{R}^n)} \\ &\leq |c| \cdot C \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}. \end{aligned}$$

Summing over  $k$  and combining with the bound for the second term proves the desired claim.  $\square$

## 5. Stability to Diffeomorphisms

We now focus on the stability of  $S_{\text{cont},q}^m f$  for general diffeomorphisms with  $\|D\tau\|_\infty < \frac{1}{2n}$ . The corresponding operator for diffeomorphisms is defined as  $L_\tau f(x) = f(x - \tau(x))$ .

### 5.1. Stability to Diffeomorphisms When $q = 2$

**Proposition 19.** Assume  $\psi$  and its first and second order derivatives have decay<sup>1</sup> in  $O((1 + |x|)^{-n-3})$ , and  $\int_{\mathbb{R}^n} \psi(x) dx = 0$ . Then for every  $\tau \in C^2(\mathbb{R}^n)$  with  $\|D\tau\|_\infty \leq \frac{1}{2n}$ , there exists  $\tilde{C}_n > 0$  such that:

$$\|[\mathcal{W}, L_\tau]\|_{\mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)} \leq \tilde{C}_n \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right).$$

*Proof.* The argument is a continuous version of Lemma 2.14 in [5]. We will first show how to transform our commutator term into an analogous commutator term from [5]. To shorten notation, we will denote  $\|[\mathcal{W}, L_\tau]\|_{\mathbf{L}^2(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)}$  as  $\|[\mathcal{W}, L_\tau]\|$ . We have

$$\begin{aligned} \|[\mathcal{W}, L_\tau]f\|_{\mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)}^2 &= \int_0^\infty \|[\mathcal{W}_t, L_\tau]f\|_2^2 \frac{dt}{t^{n+1}} \\ &= \int_0^\infty \|\psi_t * (L_\tau f) - L_\tau(\psi_t * f)\|_2^2 \frac{dt}{t^{n+1}} \\ &= \int_0^\infty \int_{\mathbb{R}^n} |\psi_t * (L_\tau f) - L_\tau(\psi_t * f)|^2 dx \frac{dt}{t^{n+1}}. \end{aligned}$$

<sup>1</sup>Similar to [13], we have found that there needs to be  $O((1 + |x|)^{-n-2-\alpha})$  decay for some  $\alpha > 0$  to bound (E.26) in [5].

Notice that  $\psi_{\frac{1}{t}}(x) = t^{n/2}\psi(tx)$ . Use the change of variables  $t = \frac{1}{\lambda}$  to get

$$\begin{aligned}\|[\mathcal{W}, L_\tau]f\|_{\mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)}^2 &= \int_0^\infty \left\| \psi_{\frac{1}{\lambda}} * (L_\tau f) - L_\tau(\psi_{\frac{1}{\lambda}} * f) \right\|_2^2 \lambda^{n-1} d\lambda \\ &= \int_0^\infty \left\| \lambda^{n/2} \psi_{\frac{1}{\lambda}} * (L_\tau f) - L_\tau(\lambda^{n/2} \psi_{\frac{1}{\lambda}} * f) \right\|_2^2 \frac{d\lambda}{\lambda}.\end{aligned}$$

Define  $\mathscr{W}_\lambda f = f * \lambda^{n/2} \psi_{\frac{1}{\lambda}}$  with  $\lambda^{n/2} \psi_{\frac{1}{\lambda}}(x) = \lambda^n \psi(\lambda x)$ . In other words,  $\mathscr{W}_t$  is a convolution with an  $\mathbf{L}^1$  normalized wavelet, which matches with the normalization in [5]. Now we have

$$\|[\mathcal{W}, L_\tau]f\|_{\mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)}^2 = \int_0^\infty \|[\mathscr{W}_\lambda, L_\tau]f\|_2^2 \frac{d\lambda}{\lambda}.$$

This implies

$$[\mathcal{W}, L_\tau]^* [\mathcal{W}, L_\tau] = \int_0^\infty [\mathscr{W}_\lambda, L_\tau]^* [\mathscr{W}_\lambda, L_\tau] \frac{d\lambda}{\lambda}$$

Defining  $K_\lambda = \mathscr{W}_\lambda - L_\tau \mathscr{W}_\lambda L_\tau^{-1}$  so that  $[\mathscr{W}_\lambda, L_\tau] = K_\lambda L_\tau$ , we have:

$$\begin{aligned}\|[\mathcal{W}, L_\tau]\| &= \|[\mathcal{W}, L_\tau]^* [\mathcal{W}, L_\tau]\|^{1/2} \\ &= \left\| \int_0^\infty [\mathscr{W}_\lambda, L_\tau]^* [\mathscr{W}_\lambda, L_\tau] \frac{d\lambda}{\lambda} \right\|^{1/2} \\ &= \left\| \int_0^\infty L_\tau^* K_\lambda^* K_\lambda L_\tau \frac{d\lambda}{\lambda} \right\|^{1/2} \\ &\leq \|L_\tau\| \cdot \left\| \int_0^\infty K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2},\end{aligned}$$

Since  $\|L_\tau f\|_2^2 \leq \left( \frac{1}{1-n\|D\tau\|_\infty} \right) \|f\|_2^2$ ,

$$\|L_\tau\| \leq \frac{1}{1-n\|D\tau\|_\infty} \leq 2$$

and the problem is reduced to bounding  $\left\| \int_0^\infty K_\lambda^* K_\lambda \lambda^{-1} d\lambda \right\|^{1/2}$ . Let  $\gamma \geq 1$ . The integral is divided into three pieces:

$$\begin{aligned}\left\| \int_0^\infty K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2} &\leq \left( \left\| \int_0^{2^{-\gamma}} K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\| + \left\| \int_{2^{-\gamma}}^1 K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\| + \left\| \int_1^\infty K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\| \right)^{1/2} \\ &\leq \left\| \int_0^{2^{-\gamma}} K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2} + \left\| \int_{2^{-\gamma}}^1 K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2} + \left\| \int_1^\infty K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2} \\ &= P_1 + P_2 + P_3.\end{aligned}$$

To bound  $P_1$ , we decompose  $K_\lambda = \tilde{K}_{\lambda,1} + \tilde{K}_{\lambda,2}$ , where the kernels defining  $\tilde{K}_{\lambda,1}, \tilde{K}_{\lambda,2}$  are

$$\begin{aligned}\tilde{k}_{\lambda,1}(x, u) &:= (1 - \det(I - D\tau(u))) \lambda^n \psi(\lambda(x - u)) \\ &:= a(u) \lambda^n \psi(\lambda(x - u)), \\ \tilde{k}_{\lambda,2}(x, u) &:= \det(I - D\tau(u)) (\lambda^n \psi(\lambda(x - u)) - \lambda^n \psi(\lambda(x - \tau(x) - u + \tau(u))),\end{aligned}$$

respectively. Since our normalization matches with [5], E.13 implies that there exists a constant  $C_n$  such that

$$\|\tilde{K}_{\lambda,2}\| \leq C_n \lambda \|\Delta\tau\|_\infty.$$

We want to prove that

$$\left\| \int_0^1 \tilde{K}_{\lambda,1}^* \tilde{K}_{\lambda,1} \frac{d\lambda}{\lambda} \right\|^{1/2} \leq C_n \|D\tau\|_\infty.$$

Notice that the operator

$$T := \int_0^1 \tilde{K}_{\lambda,1}^* \tilde{K}_{\lambda,1} \frac{d\lambda}{\lambda}$$

is positive, so Tonelli's Theorem gives

$$\begin{aligned} \|T\| &= \sup_{\|f\|_2=1} \langle Tf, f \rangle \\ &= \sup_{\|f\|_2=1} \int_0^1 \|\tilde{K}_{\lambda,1} f\|_2^2 \frac{d\lambda}{\lambda}. \end{aligned}$$

Thus, it is sufficient to bound  $\int_0^1 \|\tilde{K}_{\lambda,1} f\|_2^2 \frac{d\lambda}{\lambda}$  for  $f \in \mathbf{L}^2(\mathbb{R}^n)$  with  $\|f\|_2 = 1$ . Since

$$\tilde{K}_{\lambda,1}(x, u) = a(u) \lambda^n \psi(\lambda(x - u)), \quad a(u) = 1 - \det(I - D\tau(u)),$$

the operator  $\tilde{K}_{\lambda,1}$  multiplies by  $a$  and then convolves with the  $\mathbf{L}^1$  normalized wavelet  $\lambda^{n/2} \psi_{\frac{1}{\lambda}}$ , that is,

$$\tilde{K}_{\lambda,1} f = \left( \lambda^{n/2} \psi_{\frac{1}{\lambda}} \right) * (af).$$

Let  $F := af \in \mathbf{L}^2(\mathbb{R}^n)$  and let  $\mathcal{F}$  denote the Fourier Transform, so that  $\mathcal{F} \left( \lambda^{n/2} \psi_{\frac{1}{\lambda}} \right) (\omega) = \hat{\psi} \left( \frac{\omega}{\lambda} \right)$ . Now use Plancherel's Theorem and Tonelli's Theorem to get

$$\begin{aligned} \int_0^1 \|\tilde{K}_{\lambda,1} f\|_2^2 \frac{d\lambda}{\lambda} &= \frac{1}{(2\pi)^n} \int_0^1 \int_{\mathbb{R}^n} |\hat{\psi} \left( \frac{\omega}{\lambda} \right)|^2 |\hat{F}(\omega)|^2 d\omega \frac{d\lambda}{\lambda} \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} |\hat{F}(\omega)|^2 \left( \int_0^1 |\hat{\psi} \left( \frac{\omega}{\lambda} \right)|^2 \frac{d\lambda}{\lambda} \right) d\omega. \end{aligned}$$

To finish up the argument, we make a substitution to rewrite

$$\int_0^1 |\hat{\psi} \left( \frac{\omega}{\lambda} \right)|^2 \frac{d\lambda}{\lambda} = \int_1^\infty |\hat{\psi}(\lambda\omega)|^2 \frac{d\lambda}{\lambda}.$$

Using our decay assumptions on  $\psi$  and its partial derivatives, from Problem 6.1.3 in [22], we know that

$$|\hat{\psi}(\omega)| \leq M_\psi \min\{|\omega|, |\omega|^{-2}\}$$

for some constant  $M_\psi$ . Now, consider the quantity  $\int_0^\infty |\hat{\psi}(\lambda\omega)|^2 \frac{d\lambda}{\lambda}$ . Without loss of generality, assume that  $|\omega| = 1$  since dilations of  $\omega$  do not change the integral. It follows that

$$\int_0^\infty |\hat{\psi}(\lambda\omega)|^2 \frac{d\lambda}{\lambda} \leq M_\psi^2 \int_0^1 \lambda d\lambda + M_\psi^2 \int_1^\infty \lambda^{-5} d\lambda < \infty,$$

and we conclude that

$$\int_1^\infty |\hat{\psi}(\lambda\omega)|^2 \frac{d\lambda}{\lambda} \leq A_\psi$$

for some constant  $A_\psi$  independent of  $\omega$ . Finally, we see that  $\|D\tau\|_\infty \leq \frac{1}{2^n}$  implies

$$\begin{aligned}\|a\|_\infty &\leq (1 + \|D\tau\|_\infty)^n - 1 \\ &\leq n\|D\tau\|_\infty(1 + \|D\tau\|_\infty)^{n-1} \\ &\leq 2n\|D\tau\|_\infty,\end{aligned}$$

so Plancherel's Theorem gives

$$\begin{aligned}\frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} |\hat{F}(\omega)|^2 \left( \int_0^1 |\hat{\psi}(\frac{\omega}{\lambda})|^2 \frac{d\lambda}{\lambda} \right) d\omega &\leq \frac{A_\psi}{(2\pi)^n} \int_{\mathbb{R}^n} |\hat{F}(\omega)|^2 d\omega \\ &= A_\psi \int_{\mathbb{R}^n} |a(z)f(z)|^2 dz \\ &\leq 4n^2 \|D\tau\|_\infty^2 A_\psi \|f\|_2^2.\end{aligned}$$

Thus, we have the desired bound on  $\left\| \int_0^1 \tilde{K}_{\lambda,1}^* \tilde{K}_{\lambda,1} \frac{d\lambda}{\lambda} \right\|^{1/2}$ , with  $C_n = 2n \sqrt{A_\psi}$ .

Substituting everything in yields

$$\begin{aligned}\left\| \int_0^{2^{-\gamma}} K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2} &= \left\| \int_0^{2^{-\gamma}} (\tilde{K}_{\lambda,1} + \tilde{K}_{\lambda,2})^* (\tilde{K}_{\lambda,1} + \tilde{K}_{\lambda,2}) \frac{d\lambda}{\lambda} \right\|^{1/2} \\ &= \left\| \int_0^{2^{-\gamma}} (\tilde{K}_{\lambda,1}^* \tilde{K}_{\lambda,1} + \tilde{K}_{\lambda,1}^* \tilde{K}_{\lambda,2} + \tilde{K}_{\lambda,2}^* \tilde{K}_{\lambda,1} + \tilde{K}_{\lambda,2}^* \tilde{K}_{\lambda,2}) \frac{d\lambda}{\lambda} \right\|^{1/2} \\ &\leq \left( \left\| \int_0^{2^{-\gamma}} \tilde{K}_{\lambda,1}^* \tilde{K}_{\lambda,1} \frac{d\lambda}{\lambda} \right\| + \left\| \int_0^{2^{-\gamma}} \tilde{K}_{\lambda,1}^* \tilde{K}_{\lambda,2} + \tilde{K}_{\lambda,2}^* \tilde{K}_{\lambda,1} + \tilde{K}_{\lambda,2}^* \tilde{K}_{\lambda,2} \frac{d\lambda}{\lambda} \right\| \right)^{1/2} \\ &\leq \left( \left\| \int_0^{2^{-\gamma}} \tilde{K}_{\lambda,1}^* \tilde{K}_{\lambda,1} \frac{d\lambda}{\lambda} \right\| + \int_0^{2^{-\gamma}} \|\tilde{K}_{\lambda,2}\|^2 \frac{d\lambda}{\lambda} + \int_0^{2^{-\gamma}} 2\|\tilde{K}_{\lambda,1}\| \|\tilde{K}_{\lambda,2}\| \frac{d\lambda}{\lambda} \right)^{1/2} \\ &\leq \left\| \int_0^{2^{-\gamma}} \tilde{K}_{\lambda,1}^* \tilde{K}_{\lambda,1} \frac{d\lambda}{\lambda} \right\|^{1/2} + \left( \int_0^{2^{-\gamma}} \|\tilde{K}_{\lambda,2}\|^2 \frac{d\lambda}{\lambda} \right)^{1/2} + \left( \int_0^{2^{-\gamma}} 2\|\tilde{K}_{\lambda,1}\| \|\tilde{K}_{\lambda,2}\| \frac{d\lambda}{\lambda} \right)^{1/2} \\ &\leq 2C_n \left( \|D\tau\|_\infty + \|\Delta\tau\|_\infty \left( \int_0^{2^{-\gamma}} \lambda^2 \frac{d\lambda}{\lambda} \right)^{1/2} + \|D\tau\|_\infty^{1/2} \|\Delta\tau\|_\infty^{1/2} \left( \int_0^{2^{-\gamma}} 2\lambda \frac{d\lambda}{\lambda} \right)^{1/2} \right) \\ &\leq 2C_n \left( \|D\tau\|_\infty + 2^{-\gamma} \|\Delta\tau\|_\infty + 2^{-\gamma/2} \|D\tau\|_\infty^{1/2} \|\Delta\tau\|_\infty^{1/2} \right) \\ &\leq 4C_n (\|D\tau\|_\infty + 2^{-\gamma} \|\Delta\tau\|_\infty).\end{aligned}$$

To bound  $P_3$ , we decompose  $K_\lambda = K_{\lambda,1} + K_{\lambda,2}$ , where the kernels defining  $K_{\lambda,1}, K_{\lambda,2}$  are

$$\begin{aligned}k_{\lambda,1}(x, u) &= \lambda^n \psi(\lambda(x - u)) - \lambda^n \psi(\lambda(I - D\tau(u))(x - u)) \det(I - D\tau(u)) \\ k_{\lambda,2}(x, u) &= \det(I - D\tau(u)) \lambda^n \psi(\lambda(I - D\tau(u))(x - u)) - \lambda^n \psi(\lambda(x - \tau(x) - u + \tau(u))).\end{aligned}$$

A similar computation to the one for  $P_1$  shows that:

$$\left\| \int_1^\infty K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2} \leq \left\| \int_1^\infty K_{\lambda,1}^* K_{\lambda,1} \frac{d\lambda}{\lambda} \right\|^{1/2} + \left( \int_1^\infty \|K_{\lambda,2}\|^2 \frac{d\lambda}{\lambda} \right)^{1/2} + \left( \int_1^\infty 2\|K_{\lambda,1}\| \|K_{\lambda,2}\| \frac{d\lambda}{\lambda} \right)^{1/2}$$

Letting  $Q_j = K_{2^j,1}^* K_{2^j,1}$ , it is shown in [5] that:

$$\begin{aligned}\|K_{\lambda,1}\| &\leq C_n \|D\tau\|_\infty \\ \|K_{\lambda,2}\| &\leq \min\{\lambda^{-1} \|D^2\tau\|_\infty, \|D\tau\|_\infty\} \\ \|Q_j Q_\ell\| &\leq C_n^2 2^{-|j-\ell|} (\|D\tau\|_\infty + \|D^2\tau\|_\infty)^4\end{aligned}$$

so that

$$\begin{aligned} \left\| \int_1^\infty K_{\lambda,1}^* K_{\lambda,1} \frac{d\lambda}{\lambda} \right\|^{1/2} &= \left\| \int_0^\infty K_{2^j,1}^* K_{2^j,1} \log(2) dj \right\|^{1/2} \\ &= \sqrt{\log(2)} \left\| \int_0^\infty Q_j dj \right\|^{1/2}. \end{aligned}$$

We now apply a continuous version of Cotlar's Lemma (see Ch. 7 of [26], Sec. 5.5 for the continuous extension). We define:

$$\beta(j, \ell) = \begin{cases} C_n 2^{-|j-\ell|/2} (\|D\tau\|_\infty + \|D^2\tau\|_\infty)^2 & j \geq 0 \text{ and } \ell \geq 0 \\ 0 & \text{otherwise} \end{cases}.$$

Defining  $Q_j = 0$  for  $j < 0$ , we have  $\|Q_j^* Q_\ell\| \leq \beta(j, \ell)^2$  and  $\|Q_j Q_\ell^*\| \leq \beta(j, \ell)^2$  for all  $j, \ell$ . Thus by Cotlar's Lemma:

$$\begin{aligned} \left\| \int_{\mathbb{R}} Q_j dj \right\| &\leq \sup_{j \in \mathbb{R}} \int_{\mathbb{R}} \beta(j, \ell) d\ell, \\ \left\| \int_0^\infty Q_j dj \right\| &\leq \sup_{j \geq 0} \int_0^\infty \beta(j, \ell) d\ell \\ &\leq C_n (\|D\tau\|_\infty + \|D^2\tau\|_\infty)^2 \left( \sup_{j \geq 0} \int_0^\infty 2^{-|j-\ell|/2} d\ell \right). \end{aligned}$$

Now observing that with the change of variable  $\lambda_1 = 2^j, \lambda_2 = 2^\ell$ , we have  $2^{-|j-\ell|/2} = \frac{\lambda_1}{\lambda_2} \wedge \frac{\lambda_2}{\lambda_1}$ , we obtain:

$$\begin{aligned} \sup_{j \geq 0} \int_0^\infty 2^{-|j-\ell|/2} d\ell &= \sup_{\lambda_1 \geq 1} \int_1^\infty \frac{(\lambda_1 \wedge \lambda_2)}{\sqrt{\lambda_1 \lambda_2}} \frac{d\lambda_2}{\ln(2) \lambda_2} \\ &= \frac{1}{\ln(2)} \sup_{\lambda_1 \geq 1} \left( \int_1^{\lambda_1} \frac{1}{\sqrt{\lambda_1 \lambda_2}} d\lambda_2 + \int_{\lambda_1}^\infty \frac{\sqrt{\lambda_1}}{\lambda_2^{3/2}} d\lambda_2 \right) \\ &= \frac{1}{\ln(2)} \sup_{\lambda_1 \geq 1} \left( \frac{1}{\sqrt{\lambda_1}} (2\sqrt{\lambda_1} - 2) + \sqrt{\lambda_1} \left( \frac{2}{\sqrt{\lambda_1}} \right) \right) \\ &= \frac{1}{\ln(2)} \sup_{\lambda_1 \geq 1} \left( 4 - \frac{2}{\sqrt{\lambda_1}} \right) \\ &= \frac{4}{\ln(2)} \end{aligned}$$

and conclude that

$$\left\| \int_1^\infty K_{\lambda,1}^* K_{\lambda,1} \frac{d\lambda}{\lambda} \right\|^{1/2} \leq 3C_n (\|D\tau\|_\infty + \|D^2\tau\|_\infty).$$

Thus we have:

$$\left\| \int_1^\infty K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2} \leq \left\| \int_1^\infty K_{\lambda,1}^* K_{\lambda,1} \frac{d\lambda}{\lambda} \right\|^{1/2} + \left( \int_1^\infty \|K_{\lambda,2}\|^2 \frac{d\lambda}{\lambda} \right)^{1/2} + \left( \int_1^\infty 2\|K_{\lambda,1}\| \|K_{\lambda,2}\| \frac{d\lambda}{\lambda} \right)^{1/2}.$$

Now we see that there exists a constant  $C_n$  such that

$$\begin{aligned} \left\| \int_1^\infty K_{\lambda,1}^* K_{\lambda,1} \frac{d\lambda}{\lambda} \right\|^{1/2} &\leq C_n (\|D\tau\|_\infty + \|D^2\tau\|_\infty) \\ \left( \int_1^\infty \|K_{\lambda,2}\|^2 \frac{d\lambda}{\lambda} \right)^{1/2} &\leq C_n \|D^2\tau\|_\infty \left( \int_1^\infty \lambda^{-2} \frac{d\lambda}{\lambda} \right)^{1/2} \\ \left( \int_1^\infty 2\|K_{\lambda,1}\| \|K_{\lambda,2}\| \frac{d\lambda}{\lambda} \right)^{1/2} &\leq C_n \|D\tau\|_\infty^{1/2} \|D^2\tau\|_\infty^{1/2} \left( \int_1^\infty 2\lambda^{-1} \frac{d\lambda}{\lambda} \right)^{1/2}. \end{aligned}$$

and

$$\begin{aligned} \left\| \int_1^\infty K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2} &\leq C_n \left( \|D\tau\|_\infty + \frac{1}{\sqrt{2}} \|D^2\tau\|_\infty + \sqrt{2} \|D\tau\|_\infty^{1/2} \|D^2\tau\|_\infty^{1/2} \right) \\ &\leq C_n \left( \|D\tau\|_\infty + \frac{1}{\sqrt{2}} \|D^2\tau\|_\infty + \frac{1}{\sqrt{2}} \|D\tau\|_\infty + \frac{1}{\sqrt{2}} \|D^2\tau\|_\infty \right) \\ &\leq 2C_n (\|D\tau\|_\infty + \|D^2\tau\|_\infty). \end{aligned}$$

Finally, we bound  $P_2$ . Note that in the previous section it was observed (shown in [5]) that

$$\begin{aligned} \|K_{\lambda,1}\| &\leq C_n \|D\tau\|_\infty \\ \|K_{\lambda,2}\| &\leq \min\{\lambda^{-1} \|D^2\tau\|_\infty, \|D\tau\|_\infty\}. \end{aligned}$$

The above two inequalities imply

$$\|K_\lambda\| = \|K_{\lambda,1} + K_{\lambda,2}\| \leq \|K_{\lambda,1}\| + \|K_{\lambda,2}\| \leq 2C_n \|D\tau\|_\infty$$

so that

$$\begin{aligned} \left\| \int_{2^{-\gamma}}^1 K_\lambda^* K_\lambda \frac{d\lambda}{\lambda} \right\|^{1/2} &\leq \left( \int_{2^{-\gamma}}^1 \|K_\lambda\|^2 \frac{d\lambda}{\lambda} \right)^{1/2} \\ &\leq 2C_n \|D\tau\|_\infty \left( \int_{2^{-\gamma}}^1 \frac{d\lambda}{\lambda} \right)^{1/2} \\ &\leq 2C_n \|D\tau\|_\infty (-\ln(2^{-\gamma}))^{1/2} \\ &\leq 2C_n \gamma^{1/2} \|D\tau\|_\infty. \end{aligned}$$

Putting everything together and since  $\gamma \geq 1$ , we obtain:

$$\begin{aligned} \|[\mathcal{W}, L_\tau]\| &\leq 2(P_1 + P_2 + P_3) \\ &\leq 4C_n (\|D\tau\|_\infty + 2^{-\gamma} \|\Delta\tau\|_\infty) + 2C_n \gamma^{1/2} \|D\tau\|_\infty + 3C_n (\|D\tau\|_\infty + \|D^2\tau\|_\infty) \\ &\leq \tilde{C}_n \left( \gamma \|D\tau\|_\infty + 2^{-\gamma} \|\Delta\tau\|_\infty + \|D^2\tau\|_\infty \right). \end{aligned}$$

Choosing  $\gamma = \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \right) \vee 1$  gives

$$\|[\mathcal{W}, L_\tau]\| \leq \tilde{C}_n \left( \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) \|D\tau\|_\infty + \|D^2\tau\|_\infty \right),$$

and the lemma is proved.  $\square$

**Theorem 20.** Assume  $\psi$  and its first and second order derivatives have decay in  $O((1+|x|)^{-n-3})$  and  $\int_{\mathbb{R}^n} \psi(x) dx = 0$ . Then for every  $\tau \in C^2(\mathbb{R}^n)$  with  $\|D\tau\|_\infty < \frac{1}{2n}$ , there exists  $C_{m,n} > 0$  and  $\hat{C}_{m,n} > 0$  such that

$$\|S_{\text{cont},2}^m f - S_{\text{cont},2}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 \leq C_{m,n} \left( \|D\tau\|_\infty^2 + \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \right) \|f\|_2^2.$$

and

$$\|S_{\text{dyad},2}^m f - S_{\text{dyad},2}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m)}^2 \leq \hat{C}_{m,n} \left( \|D\tau\|_\infty^2 + \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \right) \|f\|_2^2.$$

*Proof.* The proof is only provided for the continuous case. We have the following bound for some  $C_m$ :

$$\begin{aligned} \|S_{\text{cont},2}^m f - S_{\text{cont},2}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} &\leq \|A_2 M W V_{m-1} f - A_2 M L_\tau W V_{m-1} f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} + \|A_2 M [\mathcal{W} V_{m-1}, L_\tau] f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} \\ &\leq \|A_2 M W V_{m-1} f - A_2 M L_\tau W V_{m-1} f\|_{\mathbf{L}^2(\mathbb{R}_+^m)} + C_m \|[\mathcal{W}, L_\tau]\|_{\mathbf{L}^2(\mathbb{R}_+^m \times \mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}^n)} \|f\|_2. \end{aligned}$$

For the first term, we can mimic the dilation argument to get

$$|A_2 M W V_{m-1} f - A_2 M L_\tau W V_{m-1} f| = \|\|g\|_2 - \|L_\tau g\|_2\|.$$

The difference is the term with the diffeomorphism. Let  $y = \gamma(x) = x - \tau(x)$ . Then it follows that  $\gamma^{-1}(y) = x$  and change of variables implies that

$$\|L_\tau f\|_2^2 = \int_{\mathbb{R}^n} |f(x - \tau(x))|^2 dx = \int_{\mathbb{R}^n} |f(y)|^2 \frac{dy}{|\det(I - D\tau(\gamma^{-1}(y)))|}.$$

From [27], we have (with an obvious adjustment when  $n = 1$ )

$$1 - n\|D\tau\|_\infty \leq |\det(I - D\tau(\gamma^{-1}(y)))|.$$

Regarding an upper bound on the determinant, if  $n = 1$ , we simply see that

$$|\det(I - D\tau(\gamma^{-1}(y)))| \leq 1 + n\|D\tau\|_\infty.$$

When  $n \geq 2$ , using an upper bound in [27], we get

$$|\det(I - D\tau(\gamma^{-1}(y)))| \leq \left(1 + 2\|D\tau\|_\infty + n\|D\tau\|_\infty^2\right)^{n/2} \leq \left(1 + \frac{5}{2}\|D\tau\|_\infty\right)^{n/2}.$$

Since  $\frac{5}{2}\|D\tau\|_\infty < 1$ , set  $Q = \lceil n/2 \rceil$ , so that  $Q \leq \frac{n+1}{2}$  and, because  $1 + \frac{5}{2}\|D\tau\|_\infty > 1$ ,

$$\left(1 + \frac{5}{2}\|D\tau\|_\infty\right)^{n/2} \leq \left(1 + \frac{5}{2}\|D\tau\|_\infty\right)^Q.$$

Expanding binomially and using  $\binom{Q}{k} \leq \frac{Q^k}{k!}$  together with  $\|D\tau\|_\infty^{k-1} < (2n)^{-(k-1)}$ , we have

$$\begin{aligned} \left(1 + \frac{5}{2}\|D\tau\|_\infty\right)^Q &= 1 + \|D\tau\|_\infty \sum_{k=1}^Q \binom{Q}{k} \left(\frac{5}{2}\right)^k \|D\tau\|_\infty^{k-1} \\ &\leq 1 + \|D\tau\|_\infty \sum_{k=1}^Q \frac{Q^k}{k!} \left(\frac{5}{2}\right)^k \frac{1}{(2n)^{k-1}} \\ &= 1 + 2n\|D\tau\|_\infty \sum_{k=1}^Q \frac{1}{k!} \left(\frac{5Q}{4n}\right)^k \\ &\leq 1 + 2n\|D\tau\|_\infty \sum_{k=1}^\infty \frac{1}{k!} \left(\frac{15}{16}\right)^k \\ &= 1 + 2n \left(e^{15/16} - 1\right) \|D\tau\|_\infty, \end{aligned}$$

where we used  $\frac{5Q}{4n} \leq \frac{5(n+1)}{8n} = \frac{5}{8} \left(1 + \frac{1}{n}\right) \leq \frac{15}{16} < 1$  for  $n \geq 2$ . Writing  $C_n := 2n \left(e^{15/16} - 1\right)$ , we obtain

$$1 - n\|D\tau\|_\infty \leq |\det(I - D\tau(\gamma^{-1}(y)))| \leq 1 + C_n\|D\tau\|_\infty.$$

From the determinant bounds we have

$$\frac{1}{1 + C_n\|D\tau\|_\infty} \|f\|_2^2 \leq \|L_\tau f\|_2^2 \leq \frac{1}{1 - n\|D\tau\|_\infty} \|f\|_2^2,$$

and hence

$$(1 + C_n\|D\tau\|_\infty)^{-1/2} \|f\|_2 \leq \|L_\tau f\|_2 \leq (1 - n\|D\tau\|_\infty)^{-1/2} \|f\|_2.$$

Using the elementary inequalities  $(1+x)^{-1/2} \geq 1 - \frac{x}{2}$  for  $x \geq 0$  and  $(1-x)^{-1/2} \leq 1+x$  for  $x \in [0, \frac{1}{2}]$ , the lower bound gives

$$\|f\|_2 - \|L_\tau f\|_2 \leq \|f\|_2 \left(1 - (1 + C_n\|D\tau\|_\infty)^{-1/2}\right) \leq \frac{1}{2} C_n\|D\tau\|_\infty \|f\|_2,$$

while the upper bound gives

$$\|L_\tau f\|_2 - \|f\|_2 \leq \|f\|_2 \left((1 - n\|D\tau\|_\infty)^{-1/2} - 1\right) \leq n\|D\tau\|_\infty \|f\|_2,$$

where we used  $n\|D\tau\|_\infty \leq \frac{1}{2}$ . Combining the two, we obtain

$$|\|f\|_2 - \|L_\tau f\|_2| \leq \frac{1}{2} C_n\|D\tau\|_\infty \|f\|_2 \quad \text{for any } f \in L^2(\mathbb{R}^n),$$

since  $\frac{1}{2} C_n \geq n$ . Now we mimic the argument given for dilation stability to get

$$\|A_2 M W V_{m-1} f - A_2 M L_\tau W V_{m-1} f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^2 \leq C \|D\tau\|_\infty^2 \|f\|_2^2$$

for some constant  $C$ . For the second term, we have

$$C_m^2 \|\llbracket \mathcal{W}, L_\tau \rrbracket\|_{\mathbf{L}^2(\mathbb{R}_+^m \times \mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}^n)}^2 \|f\|_2^2 \leq C' \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \|f\|_2^2$$

for some constant  $C'$ . We now choose  $C_{n,m} = \max\{C', C\}$  to get the desired bound.  $\square$

## 5.2. Stability to Diffeomorphisms When $1 < q < 2$

**Lemma 21.** Let  $\gamma(z) = z - \tau(z)$ ,  $g(z) = f(\gamma(z))$ , and

$$K_\lambda(x, z) = \det(D\gamma(z)) \psi_\lambda(\gamma(x) - \gamma(z)) - \psi_\lambda(x - z).$$

Additionally, define

$$T_\lambda g(x) = \int_{\mathbb{R}^n} g(z) K_\lambda(x, z) dz$$

and consider  $Tg : \mathbb{R}^n \rightarrow \mathbf{L}^2(\mathbb{R}_+, \frac{d\lambda}{\lambda^{n+1}})$  defined by  $Tg(x) = (T_\lambda g(x))_{\lambda \in \mathbb{R}_+}$ . Then for the Banach space  $\mathcal{X} = \mathbf{L}^2(\mathbb{R}_+, \frac{d\lambda}{\lambda^{n+1}})$ ,

$$\|Tg\|_{\mathbf{L}_\lambda^2(\mathbb{R}^n)}^2 \leq C_{n,m} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \|f\|_2^2$$

for some constant  $C_{n,m} > 0$ .

*Proof.* Notice that

$$\begin{aligned}
\|Tg\|_{\mathbf{L}_X^2(\mathbb{R}^n)}^2 &= \int_{\mathbb{R}^n} \int_0^\infty |T_\lambda g(x)|^2 \frac{d\lambda}{\lambda^{n+1}} dx \\
&= \int_{\mathbb{R}^n} \int_0^\infty \left| \int_{\mathbb{R}^n} K_\lambda(x, z) g(z) dz \right|^2 \frac{d\lambda}{\lambda^{n+1}} dx \\
&= \int_{\mathbb{R}^n} \int_0^\infty \left| \int_{\mathbb{R}^n} f(\gamma(z)) [\det(D\gamma(z)) \psi_\lambda(\gamma(x) - \gamma(z)) - \psi_\lambda(x - z)] dz \right|^2 \frac{d\lambda}{\lambda^{n+1}} dx \\
&= \int_{\mathbb{R}^n} \int_0^\infty \left| \int_{\mathbb{R}^n} \det(D\gamma(z)) f(\gamma(z)) \psi_\lambda(\gamma(x) - \gamma(z)) dz - \int_{\mathbb{R}^n} f(\gamma(z)) \psi_\lambda(x - z) dz \right|^2 \frac{d\lambda}{\lambda^{n+1}} dx.
\end{aligned}$$

Using the change of variables  $u = \gamma(z)$ , we get

$$\begin{aligned}
\|Tg\|_{\mathbf{L}_X^2(\mathbb{R}^n)}^2 &= \int_{\mathbb{R}^n} \int_0^\infty |L_\tau(f * \psi_\lambda)(x) - (L_\tau f * \psi_\lambda)(x)|^2 \frac{d\lambda}{\lambda^{n+1}} dx \\
&= \int_{\mathbb{R}^n} \int_0^\infty |[\mathcal{W}_\lambda, L_\tau]f(x)|^2 \frac{d\lambda}{\lambda^{n+1}} dx \\
&= \int_0^\infty \int_{\mathbb{R}^n} |[\mathcal{W}_\lambda, L_\tau]f(x)|^2 dx \frac{d\lambda}{\lambda^{n+1}} \\
&= \int_0^\infty \|[\mathcal{W}_\lambda, L_\tau]f\|_2^2 \frac{d\lambda}{\lambda^{n+1}} \\
&= \|[\mathcal{W}, L_\tau]f\|_{\mathbf{L}^2(\mathbb{R}_+ \times \mathbb{R}^n)}^2 \\
&\leq C_{n,m} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \|f\|_2^2,
\end{aligned}$$

where the last inequality follows from the  $q = 2$  case.  $\square$

**Lemma 22** ([23], Marcinkiewicz Interpolation). *Let  $\mathcal{A}$  and  $\mathcal{B}$  be Banach spaces and let  $T : \mathcal{A} \rightarrow \mathcal{B}$  be a quasilinear operator defined on  $\mathbf{L}_{\mathcal{A}}^{p_0}(\mathbb{R}^n)$  and  $\mathbf{L}_{\mathcal{A}}^{p_1}(\mathbb{R}^n)$  with  $0 < p_0 < p_1$ . Furthermore, if  $T$  satisfies*

$$\|Tf\|_{\mathbf{L}_{\mathcal{B}}^{p_i \infty}(\mathbb{R}^n)} \leq M_i \|f\|_{\mathbf{L}_{\mathcal{A}}^{p_i}(\mathbb{R}^n)}$$

for  $i = 0, 1$ , then for all  $p \in (p_0, p_1)$ ,

$$\|Tf\|_{\mathbf{L}_{\mathcal{B}}^p(\mathbb{R}^n)} \leq N_p \|f\|_{\mathbf{L}_{\mathcal{A}}^p(\mathbb{R}^n)},$$

where  $N_p$  only depends on  $M_0, M_1$ , and  $p$ .

**Remark 10.** Like with the scalar valued estimate, it can be shown that  $N_p = \eta M_0^\delta M_1^{1-\delta}$ , where

$$\delta = \begin{cases} \frac{p_0(p_1 - p)}{p(p_1 - p_0)} & p_1 < \infty, \\ \frac{p_0}{p} & p_1 = \infty \end{cases}$$

and

$$\eta = \begin{cases} 2 \left( \frac{p(p_1 - p_0)}{(p - p_0)(p_1 - p)} \right)^{1/p} & p_1 < \infty, \\ 2 \left( \frac{p_0}{p - p_0} \right)^{1/p} & p_1 = \infty. \end{cases}$$

**Lemma 23.** Let  $T$  be the operator defined in Lemma 21. Let  $q \in (1, 2)$  and  $r \in (1, q)$ . Then  $T$  satisfies

$$\|Tg\|_{\mathbf{L}^{r,\infty}_{\mathcal{X}}(\mathbb{R}^n)} \leq M_r \|f\|_{\mathbf{L}^r(\mathbb{R}^n)}$$

for some constant  $M_r > 0$ , which is independent of  $\|D\tau\|_\infty$  and  $\|D^2\tau\|_\infty$ . Furthermore,  $T$  also satisfies

$$\|Tg\|_{\mathbf{L}^{2,\infty}_{\mathcal{X}}(\mathbb{R}^n)}^2 \leq \tilde{C}_n \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \|f\|_{\mathbf{L}^2(\mathbb{R}^n)}^2$$

for some constant  $\tilde{C}_n > 0$ .

*Proof.* The second inequality obviously follows from strong boundedness of the operator, so we will omit the proof. For the first inequality, the norm satisfies

$$\begin{aligned} \|Tg(x)\|_{\mathcal{X}}^2 &= \int_0^\infty \left| \int_{\mathbb{R}^n} \det(D\gamma(z)) f(\gamma(z)) \psi_\lambda(\gamma(x) - \gamma(z)) dz - \int_{\mathbb{R}^n} f(\gamma(z)) \psi_\lambda(x - z) dz \right|^2 \frac{d\lambda}{\lambda^{n+1}} \\ &= \int_0^\infty \left| \int_{\mathbb{R}^n} f(z) \psi_\lambda(\gamma(x) - z) dz - \int_{\mathbb{R}^n} f(\gamma(z)) \psi_\lambda(x - z) dz \right|^2 \frac{d\lambda}{\lambda^{n+1}} \\ &\leq 4 \int_0^\infty \left| \int_{\mathbb{R}^n} f(z) \psi_\lambda(\gamma(x) - z) dz \right|^2 \frac{d\lambda}{\lambda^{n+1}} + 4 \int_0^\infty \left| \int_{\mathbb{R}^n} f(\gamma(z)) \psi_\lambda(x - z) dz \right|^2 \frac{d\lambda}{\lambda^{n+1}} \\ &= 4|(Gf)(\gamma(x))|^2 + 4|GL_\tau f(x)|^2. \end{aligned}$$

We see

$$\|Tg(x)\|_{\mathcal{X}} \leq \sqrt{4|(Gf)(\gamma(x))|^2 + 4|GL_\tau f(x)|^2} \leq 2|(Gf)(\gamma(x))| + 2|GL_\tau f(x)|.$$

For  $\delta > 0$ , Chebyshev's inequality implies that there exists  $A_r$  such that

$$\begin{aligned} m\{\|Tg(x)\|_{\mathcal{X}} > \delta\} &\leq m\{2|(Gf)(\gamma(x))| + 2|GL_\tau f(x)| > \delta\} \\ &\leq \frac{A_r}{\delta^r} (\|(Gf)(\gamma(\cdot))\|_{\mathbf{L}^r(\mathbb{R}^n)}^r + \|GL_\tau f\|_{\mathbf{L}^r(\mathbb{R}^n)}^r). \end{aligned}$$

We want to now ensure that  $\|(Gf)(\gamma(\cdot))\|_{\mathbf{L}^r(\mathbb{R}^n)}^r$  can be bounded above by a constant multiple of  $\|Gf\|_{\mathbf{L}^r(\mathbb{R}^n)}^r$ . Since  $\gamma$  is a diffeomorphism, we can use change of variables to get

$$\begin{aligned} \|(Gf)(\gamma(\cdot))\|_{\mathbf{L}^r(\mathbb{R}^n)}^r &= \int_{\mathbb{R}^n} |Gf(\gamma(x))|^r dx \\ &= \int_{\mathbb{R}^n} |Gf(u)|^r \frac{du}{\det[(D\gamma)(\gamma^{-1}(u))]} \\ &\leq 2 \int_{\mathbb{R}^n} |Gf(x)|^r dx \\ &= 2\|Gf\|_{\mathbf{L}^r(\mathbb{R}^n)}^r. \end{aligned}$$

By Lemma 1, we get

$$\|GL_\tau f\|_{\mathbf{L}^r(\mathbb{R}^n)}^r \leq C_r \|L_\tau f\|_{\mathbf{L}^r(\mathbb{R}^n)}^r \leq 2C_r \|f\|_{\mathbf{L}^r(\mathbb{R}^n)}^r$$

for some constant  $C_r$  dependent on  $r$ . Thus, we have

$$m\{\|Tg(x)\|_{\mathcal{X}} > \delta\}^{1/r} \leq \frac{M_r}{\delta} \|f\|_{\mathbf{L}^r(\mathbb{R}^n)}$$

for some constant  $M_r > 0$ . □

**Lemma 24.** Fix  $r = \frac{1+q}{2}$  so that  $r \in (1, q)$ . For some constant  $C_{n,q} > 0$ , the operator  $T$  defined in Lemma 21 satisfies the estimate

$$\|Tg\|_{\mathbf{L}^q_{\mathcal{X}}(\mathbb{R}^n)}^q \leq C_{n,q} \eta^q M_r^{q\delta} \left( \|D\tau\|_{\infty} \left( \log \frac{\|\Delta\tau\|_{\infty}}{\|D\tau\|_{\infty}} \vee 1 \right) + \|D^2\tau\|_{\infty} \right)^{q(1-\delta)} \|f\|_q^q,$$

where  $\eta$  and  $\delta$  come from interpolation, and  $M_r$  comes from the constant for weak boundedness in Lemma 23.

*Proof.* Since  $T$  is an integral operator, it is clear that is quasilinear. Using the  $\mathbf{L}^r(\mathbb{R}^n)$  and  $\mathbf{L}^2(\mathbb{R}^n)$  estimates from the previous Lemma, we interpolate using Marcinkiewicz since  $\|g\|_r \leq 2\|f\|_r \leq 4\|g\|_r$ .  $\square$

**Theorem 25.** Let  $1 < q < 2$ . Assume  $\psi$  and its first and second order derivatives have decay in  $O((1 + |x|)^{-n-3})$ , and  $\int_{\mathbb{R}^n} \psi(x) dx = 0$ . Then for every  $\tau \in C^2(\mathbb{R}^n)$  with  $\|D\tau\|_{\infty} < \frac{1}{2n}$ , there exists  $C_{n,q} > 0$  such that

$$\|S_{\text{cont},q}f - S_{\text{cont},q}L_{\tau}f\|_{\mathbf{L}^2(\mathbb{R}_+)}^q \leq C_{n,q} \left[ \|D\tau\|_{\infty}^q + \eta^q M_r^{q\delta} \left( \|D\tau\|_{\infty} \left( \log \frac{\|\Delta\tau\|_{\infty}}{\|D\tau\|_{\infty}} \vee 1 \right) + \|D^2\tau\|_{\infty} \right)^{q(1-\delta)} \right] \|f\|_q^q.$$

*Proof.* We use the same notation as Theorem 16. Using a nearly identical argument to Corollary 17, we get

$$\begin{aligned} \|S_{\text{cont},q}f - S_{\text{cont},q}L_{\tau}f\|_{\mathbf{L}^2(\mathbb{R}_+)} &= \|A_q M \mathcal{W}f - A_q M \mathcal{W}L_{\tau}f\|_{\mathbf{L}^2(\mathbb{R}_+)} \\ &= \|A_q M \mathcal{W}f - A_q M L_{\tau} \mathcal{W}f + A_q M L_{\tau} \mathcal{W}f - A_q M \mathcal{W}L_{\tau}f\|_{\mathbf{L}^2(\mathbb{R}_+)} \\ &\leq \|A_q M \mathcal{W}f - A_q M L_{\tau} \mathcal{W}f\|_{\mathbf{L}^2(\mathbb{R}_+)} + \|A_q M L_{\tau} \mathcal{W}f - A_q M \mathcal{W}L_{\tau}f\|_{\mathbf{L}^2(\mathbb{R}_+)} \\ &\leq \|(A_q M - A_q M L_{\tau}) \mathcal{W}f\|_{\mathbf{L}^2(\mathbb{R}_+)} + \|A_q M[\mathcal{W}, L_{\tau}]f\|_{\mathbf{L}^2(\mathbb{R}_+)}. \end{aligned}$$

The first term,  $\|(A_q M - A_q M L_{\tau}) \mathcal{W}f\|_{\mathbf{L}^2(\mathbb{R}_+)}$ , can be bounded using an argument identical to the  $q = 2$  case. In particular, we can prove that

$$(1 - n\|D\tau\|_{\infty})\|f\|_q \leq (1 - n\|D\tau\|_{\infty})^{1/q}\|f\|_q \leq \|L_{\tau}f\|_q$$

and

$$\|L_{\tau}f\|_q \leq (1 + 2n\|D\tau\|_{\infty})^{1/q}\|f\|_q \leq (1 + 2n\|D\tau\|_{\infty})\|f\|_q,$$

which means

$$\|(A_q M - A_q M L_{\tau}) \mathcal{W}f\|_{\mathbf{L}^2(\mathbb{R}_+)}^q \leq C \|D\tau\|_{\infty}^q \|f\|_q^q.$$

For the other term,

$$\|A_q M[\mathcal{W}, L_{\tau}]f\|_{\mathbf{L}^2(\mathbb{R}_+)}^q = \left( \int_0^{\infty} \left[ \int_{\mathbb{R}^n} |(L_{\tau}f * \psi_{\lambda})(x) - L_{\tau}(f * \psi_{\lambda})(x)|^q dx \right]^{2/q} \frac{d\lambda}{\lambda^{n+1}} \right)^{q/2}.$$

Now, expand convolution and then use change of variables to get

$$\begin{aligned}
& \|A_q M[\mathcal{W}, L_\tau] f\|_{\mathbf{L}^2(\mathbb{R}_+)}^q \\
&= \left( \int_0^\infty \left[ \int_{\mathbb{R}^n} \left| \int_{\mathbb{R}^n} f(\gamma(z)) (\det(D\gamma(z))) \psi_\lambda(\gamma(x) - \gamma(z)) - \psi_\lambda(x - z) dz \right|^q dx \right]^{2/q} \frac{d\lambda}{\lambda^{n+1}} \right)^{q/2} \\
&= \left( \int_0^\infty \left[ \int_{\mathbb{R}^n} \left| \int_{\mathbb{R}^n} g(z) K_\lambda(x, z) dz \right|^q dx \right]^{2/q} \frac{d\lambda}{\lambda^{n+1}} \right)^{q/2} \\
&= \left( \int_0^\infty \left[ \int_{\mathbb{R}^n} |T_\lambda g(x)|^q dx \right]^{2/q} \frac{d\lambda}{\lambda^{n+1}} \right)^{q/2} \\
&\leq \int_{\mathbb{R}^n} \left[ \int_0^\infty |T_\lambda g(x)|^2 \frac{d\lambda}{\lambda^{n+1}} \right]^{q/2} dx \\
&= \int_{\mathbb{R}^n} \|Tg(x)\|_{\mathbf{L}^2(\mathbb{R}^+, \frac{d\lambda}{\lambda^{n+1}})}^q dx \\
&= \|Tg\|_{\mathbf{L}^q_{\mathcal{X}}(\mathbb{R}^n)}^q \\
&\leq C_n \eta^q M_r^{q\delta} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^{q(1-\delta)} \|f\|_q^q.
\end{aligned}$$

Thus, the proof is complete.  $\square$

**Corollary 26.** Let  $1 < q < 2$ . Assume  $\psi$  and its first and second order derivatives have decay in  $O((1 + |x|)^{-n-3})$ , and  $\int_{\mathbb{R}^n} \psi(x) dx = 0$ . Then for every  $\tau \in C^2(\mathbb{R}^n)$  with  $\|D\tau\|_\infty < \frac{1}{2n}$ , there exist constants  $C_{n,m}, \hat{C}_{n,m} > 0$  such that

$$\|S_{cont,q}^m f - S_{cont,q}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m)}^q \leq C_{n,m} \left[ \|D\tau\|_\infty^q + \eta^q M_r^{q\delta} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^{q(1-\delta)} \right] \|f\|_q^q$$

and

$$\|S_{dyad,q}^m f - S_{dyad,q}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m)}^q \leq \hat{C}_{n,m} \left[ \|D\tau\|_\infty^q + \eta^q M_r^{q\delta} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^{q(1-\delta)} \right] \|f\|_q^q.$$

**Remark 11.** This bound is not exactly the same as the definition for stability to diffeomorphisms in [5], but the idea is similar. Since  $r$  is fixed, so is  $\delta$ . It is easy to confirm that  $\delta = \frac{1}{1+q} \in \left(\frac{1}{3}, \frac{1}{2}\right)$  when using Marcinkiewicz interpolation in Lemma 24, so

$$C_{n,q} \eta^q M_r^{q\delta} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^{q(1-\delta)} \rightarrow 0$$

when  $\|D\tau\|_\infty \rightarrow 0$  and  $\|D^2\tau\|_\infty \rightarrow 0$ .

## 6. Equivariance and Invariance to Rotations

We now consider adding group actions to our scattering transform and prove invariance to rotations. Let  $\text{SO}(n)$  be the group of  $n \times n$  rotation matrices. Since  $\text{SO}(n)$  is a compact Lie group, we can define a Haar measure, say  $\mu$ , with  $\mu(\text{SO}(n)) < \infty$ . We say that  $f \in \mathbf{L}^2(\text{SO}(n))$  if and only if  $f$  is  $\mu$ -measurable and  $\int_{\text{SO}(n)} |f(r)|^2 d\mu(r) < \infty$ .

### 6.1. Rotation Equivariant Representations

Let  $\psi : \mathbb{R}^n \rightarrow \mathbb{R}$  be a wavelet. Define

$$\psi_{\lambda,R}(x) = \lambda^{-n/2} \psi(\lambda^{-1} R^{-1} x),$$

where  $R \in \text{SO}(n)$  is a  $n \times n$  rotation matrix. The continuous and dyadic wavelet transforms of  $f$  are given by

$$\begin{aligned} \mathcal{W}_{\text{Rot}} f &:= \{f * \psi_{\lambda,R}(x) : x \in \mathbb{R}^n, \lambda \in (0, \infty), R \in \text{SO}(n)\}, \\ \mathcal{W}_{\text{Rot}} f &:= \{f * \psi_{j,R}(x) : x \in \mathbb{R}^n, j \in \mathbb{Z}, R \in \text{SO}(n)\}. \end{aligned}$$

We will first consider a translation invariant and rotation equivariant formulation of continuous and dyadic one-layer scattering using

$$\begin{aligned} \mathfrak{S}_{\text{cont},q} f(\lambda, R) &:= \|f * \psi_{\lambda,R}\|_q, \\ \mathfrak{S}_{\text{dyad},q} f(j, R) &:= \|f * \psi_{j,R}\|_q. \end{aligned}$$

The translation invariance of our representation follows from translation invariance of the norm. For rotation equivariance, notice that if  $f_{\tilde{R}}(x) := f(\tilde{R}^{-1}x)$ , then we have

$$\begin{aligned} \mathfrak{S}_{\text{cont},q} f_{\tilde{R}}(\lambda, R) &= \mathfrak{S}_{\text{cont},q} f(\lambda, \tilde{R}^{-1}R), \\ \mathfrak{S}_{\text{dyad},q} f_{\tilde{R}}(j, R) &= \mathfrak{S}_{\text{dyad},q} f(j, \tilde{R}^{-1}R). \end{aligned}$$

Now suppose we have  $m$  layers again. Then we define our  $m$  layer transforms by

$$\begin{aligned} \mathfrak{S}_{\text{cont},q}^m f(\lambda_1, \dots, \lambda_m, R_1, \dots, R_m) &:= \| |f * \psi_{\lambda_1, R_1}| * \dots * \psi_{\lambda_m, R_m} \|_q, \\ \mathfrak{S}_{\text{dyad},q}^m f(j_1, \dots, j_m, R_1, \dots, R_m) &:= \| |f * \psi_{j_1, R_1}| * \dots * \psi_{j_m, R_m} \|_q. \end{aligned}$$

and rotation equivariance implies

$$\begin{aligned} \mathfrak{S}_{\text{cont},q}^m f_{\tilde{R}}(\lambda_1, \dots, \lambda_m, R_1, \dots, R_m) &= \mathfrak{S}_{\text{cont},q}^m f(\lambda_1, \dots, \lambda_m, \tilde{R}^{-1}R_1, \dots, \tilde{R}^{-1}R_m), \\ \mathfrak{S}_{\text{dyad},q}^m f_{\tilde{R}}(j_1, \dots, j_m, R_1, \dots, R_m) &= \mathfrak{S}_{\text{dyad},q}^m f(j_1, \dots, j_m, \tilde{R}^{-1}R_1, \dots, \tilde{R}^{-1}R_m). \end{aligned}$$

The norm we will use is similar to our previous formulations. Denote the scattering norm for the continuous transform as

$$\|\mathfrak{S}_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^m}^q := \left( \int_0^\infty \int_{\text{SO}(n)} \dots \int_0^\infty \int_{\text{SO}(n)} \| |f * \psi_{\lambda_1, R_1}| * \dots * \psi_{\lambda_m, R_m} \|_q^2 d\mu_1(R_1) \frac{d\lambda_1}{\lambda_1^{n+1}} \dots d\mu_m(R_m) \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{q/2}$$

For the dyadic transform, we denote the norm using

$$\|\mathfrak{S}_{\text{dyad},q}^m f\|_{\ell^2(\mathbb{Z}^m) \times \text{SO}(n)^m}^q := \left( \sum_{j_1 \in \mathbb{Z}} \int_{\text{SO}(n)} \dots \sum_{j_m \in \mathbb{Z}} \int_{\text{SO}(n)} \| |f * \psi_{j_1, R_1}| * \dots * \psi_{j_m, R_m} \|_q^2 d\mu_1(R_1) \dots d\mu_m(R_m) \right)^{q/2}.$$

We will start by proving that these formulations of the scattering transform are well defined, and prove properties about stability to diffeomorphisms like in previous sections.

**Lemma 27.** *Let  $\psi$  be a wavelet that satisfies properties (24) and (25).*

- If  $1 < q \leq 2$ , we have  $\mathfrak{S}_{\text{cont},q}^m : \mathbf{L}^q(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^m$  and  $\mathfrak{S}_{\text{dyad},q}^m : \mathbf{L}^q(\mathbb{R}^n) \rightarrow \ell^2(\mathbb{Z}^m) \times \text{SO}(n)^m$ .
- If  $q = 1$ , then for every  $\varepsilon > 0$  we have  $\mathfrak{S}_{\text{cont},1}^m : \mathbf{H}^1(\mathbb{R}^n) \rightarrow \mathbf{L}_\varepsilon^2(\mathbb{R}_+^m) \times \text{SO}(n)^m$  and  $\mathfrak{S}_{\text{dyad},1}^m : \mathbf{H}^1(\mathbb{R}^n) \rightarrow \ell_\varepsilon^2(\mathbb{Z}^m) \times \text{SO}(n)^m$ , with the damped norms of Definition 9. For  $m = 1$  the damping is unnecessary, and for  $m \geq 2$  it cannot be removed, by Proposition 7.

- If  $\psi$  is also a Littlewood-Paley wavelet, we have

$$\begin{aligned}\|\mathfrak{S}_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^m}^2 &= \mu(\text{SO}(n))^m C_\psi^m \|f\|_2^2, \\ \|\mathfrak{S}_{\text{dyad},q}^m f\|_{\ell^2(\mathbb{Z}^m) \times \text{SO}(n)^m}^2 &= \mu(\text{SO}(n))^m \hat{C}_\psi^m \|f\|_2^2.\end{aligned}$$

*Proof.* We prove the first and third claim. The second claim is almost identical to the first claim, so the proof will be omitted for brevity. Note that we will only provide arguments for the continuous scattering transform since the proofs for the dyadic transform are very similar. By Fubini Theorem and boundedness of the  $m$ -layer scattering transform, there exists a constant  $C_q > 0$ , which is dependent on  $q$ , such that

$$\begin{aligned}\|\mathfrak{S}_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^m}^q &= \left[ \int_0^\infty \int_{\text{SO}(n)} \cdots \int_0^\infty \int_{\text{SO}(n)} \|f * \psi_{\lambda_1, R_1} * \cdots * \psi_{\lambda_m, R_m}\|_q^2 d\mu(R_m) \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots d\mu(R_1) \frac{d\lambda_m}{\lambda_m^{n+1}} \right]^{q/2} \\ &= \left[ \int_{\text{SO}(n)} \cdots \int_{\text{SO}(n)} \left( \int_0^\infty \cdots \int_0^\infty \|f * \psi_{\lambda_1, R_1} * \cdots * \psi_{\lambda_m, R_m}\|_q^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{\frac{q}{2} \cdot \frac{2}{q}} d\mu(R_1) \cdots d\mu(R_m) \right]^{q/2} \\ &\leq \left[ \int_{\text{SO}(n)} \cdots \int_{\text{SO}(n)} (C_q^{mq} \|f\|_q^q)^{2/q} d\mu(R_1) \cdots d\mu(R_m) \right]^{q/2} \\ &= C_q^{mq} \mu(\text{SO}(n))^{mq/2} \|f\|_q^q\end{aligned}$$

because each  $\psi_{\lambda_i, R_i}$  is still a wavelet with sufficient decay even if the rotation is applied. For the third claim, we see that

$$\begin{aligned}\|\mathfrak{S}_{\text{cont},2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^m}^2 &= \int_{\text{SO}(n)} \cdots \int_{\text{SO}(n)} \left( \int_0^\infty \cdots \int_0^\infty \|f * \psi_{\lambda_1, R_1} * \cdots * \psi_{\lambda_m, R_m}\|_{\mathbf{L}^2(\mathbb{R}^n)}^2 \frac{d\lambda_1}{\lambda_1^{n+1}} \cdots \frac{d\lambda_m}{\lambda_m^{n+1}} \right) d\mu(R_1) \cdots d\mu(R_m) \\ &= \int_{\text{SO}(n)} \cdots \int_{\text{SO}(n)} C_\psi^m \|f\|_2^2 d\mu(R_1) \cdots d\mu(R_m) \\ &= \mu(\text{SO}(n))^m C_\psi^m \|f\|_2^2.\end{aligned}$$

□

**Theorem 28.** Assume  $|c| < \frac{1}{2n}$ . Let  $\tau(x) = cx$  and  $L_\tau f(x) = f((1-c)x)$ . Suppose that  $\psi$  is a wavelet that satisfies the conditions of Lemma 14. Then there exist constants  $\tilde{K}_{n,m,q}$  and  $\tilde{K}'_{n,m,q}$  dependent only on  $n$ ,  $m$ , and  $q$  such that

$$\|\mathfrak{S}_{\text{cont},q}^m f - \mathfrak{S}_{\text{cont},q}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^m}^q \leq |c|^q \cdot \tilde{K}_{n,m,q} \|f\|_q^q$$

and

$$\|\mathfrak{S}_{\text{dyad},q}^m f - \mathfrak{S}_{\text{dyad},q}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m) \times \text{SO}(n)^m}^q \leq |c|^q \cdot \tilde{K}'_{n,m,q} \|f\|_q^q.$$

Alternatively, if one of the following holds:

- $n = 1$ ,  $\psi$  is complex analytic and satisfies the conditions of Lemma 14,
- $n \geq 2$  and  $\psi$  satisfies the conditions of Lemma 5,

then for every  $\varepsilon > 0$  there exist  $\tilde{H}_{m,n}$  and  $\tilde{H}'_{m,n}$ , depending also on  $\varepsilon$ , such that

$$\|\mathfrak{S}_{\text{cont},1}^m f - \mathfrak{S}_{\text{cont},1}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^m} \leq |c| \cdot \tilde{H}_{m,n} \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}.$$

and

$$\|\mathfrak{S}_{\text{dyad},1}^m f - \mathfrak{S}_{\text{dyad},1}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m) \times \text{SO}(n)^m} \leq |c| \cdot \tilde{H}'_{m,n} \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}$$

**Theorem 29.** Let  $\tau \in C^2(\mathbb{R}^n)$ , and let  $L_\tau f(x) = f(x - \tau(x))$  with  $\|D\tau\|_\infty < \frac{1}{2n}$ . Suppose that  $\psi$  is a wavelet such that the wavelet and all its first and second partial derivatives have  $O((1 + |x|)^{-n-3})$  decay. When  $q \in (1, 2)$ , there exists a constant  $C_{n,m,q}$  dependent on  $\mu(\text{SO}(n))$ ,  $n$ ,  $m$ , and  $q$  such that

$$\begin{aligned} \|\mathfrak{S}_{\text{cont},q}^m f - \mathfrak{S}_{\text{cont},q}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^m}^q &\leq C_{n,m,q} \left[ \|D\tau\|_\infty^q + \eta^q M_r^{q\delta} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^{q(1-\delta)} \right] \|f\|_q^q, \\ \|\mathfrak{S}_{\text{dyad},q}^m f - \mathfrak{S}_{\text{dyad},q}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m) \times \text{SO}(n)^m}^q &\leq \tilde{C}_{n,m,q} \left[ \|D\tau\|_\infty^q + \eta^q M_r^{q\delta} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^{q(1-\delta)} \right] \|f\|_q^q, \\ \|\mathfrak{S}_{\text{cont},2}^m f - \mathfrak{S}_{\text{cont},2}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^m}^2 &\leq C_{n,m} \left[ \|D\tau\|_\infty^2 + \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \right] \|f\|_2^2, \\ \|\mathfrak{S}_{\text{dyad},2}^m f - \mathfrak{S}_{\text{dyad},2}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m) \times \text{SO}(n)^m}^2 &\leq \tilde{C}_{n,m} \left[ \|D\tau\|_\infty^2 + \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \right] \|f\|_2^2. \end{aligned}$$

## 6.2. Rotation Invariant Representations

The representation before was rotation equivariant, but in some tasks, we would rather have rotation invariance. In [5], the authors choose to integrate over each group action in a group of transformations. However, this will remove the information the relative angles between each action if we have multiple layers in our transform.

In the case of one layer, since there is only one angle, we use a similar formulation to [5] and define continuous and dyadic scattering transforms for rotation invariance as

$$\begin{aligned} \mathcal{S}_{\text{cont},q} f(\lambda) &= \int_{\text{SO}(n)} \|f * \psi_{\lambda,R}\|_{\mathbf{L}^q(\mathbb{R}^n)}^q d\mu(R), \\ \mathcal{S}_{\text{dyad},q} f(j) &= \int_{\text{SO}(n)} \|f * \psi_{j,R}\|_{\mathbf{L}^q(\mathbb{R}^n)}^q d\mu(R). \end{aligned}$$

The corresponding norms are given by

$$\begin{aligned} \|\mathcal{S}_{\text{cont},q} f\|_{\mathbf{L}^2(\mathbb{R}_+)}^q &:= \left[ \int_0^\infty \left[ \int_{\text{SO}(n)} \|f * \psi_{\lambda,R}\|_{\mathbf{L}^q(\mathbb{R}^n)}^q d\mu(R) \right]^{2/q} \frac{d\lambda}{\lambda^{n+1}} \right]^{q/2}, \\ \|\mathcal{S}_{\text{dyad},q} f\|_{\ell^2(\mathbb{Z})}^q &:= \left[ \sum_{j \in \mathbb{Z}} \left[ \int_{\text{SO}(n)} \|f * \psi_{j,R}\|_{\mathbf{L}^q(\mathbb{R}^n)}^q d\mu(R) \right]^{2/q} \right]^{q/2}. \end{aligned}$$

Now we generalize to the case where  $m \geq 2$ . Let  $R_1, \dots, R_m \in \text{SO}(n)$ . Define

$$\begin{aligned} \mathcal{S}_{\text{cont},q}^m f(\lambda_1, \dots, \lambda_m, R_2, \dots, R_m) &:= \int_{\text{SO}(n)} \|f * \psi_{\lambda_1, R_2 R_1} * \dots * \psi_{\lambda_m, R_m R_1}\|_q^2 d\mu(R_1), \\ \mathcal{S}_{\text{dyad},q}^m f(j_1, \dots, j_m, R_2, \dots, R_m) &:= \int_{\text{SO}(n)} \|f * \psi_{j_1, R_2 R_1} * \dots * \psi_{j_m, R_m R_1}\|_q^2 d\mu(R_1). \end{aligned}$$

The norm for the continuous transform, the norm  $\|\mathcal{S}_{\text{cont},q}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times \text{SO}(n)^{m-1}}^q$ , is given by

$$\left( \int_0^\infty \int_{\text{SO}(n)} \dots \int_0^\infty \int_{\text{SO}(n)} \int_0^\infty \mathcal{S}_{\text{cont},q}^m f(\lambda_1, \dots, \lambda_m, R_2, \dots, R_m) \frac{d\lambda_1}{\lambda_1^{n+1}} d\mu_2(R_2) \frac{d\lambda_2}{\lambda_2^{n+1}} \dots d\mu_m(R_m) \frac{d\lambda_m}{\lambda_m^{n+1}} \right)^{q/2}$$

For the dyadic transform, the norm  $\|\mathcal{S}_{\text{dyad},q}^m f\|_{\ell^2(\mathbb{Z}) \times \text{SO}(n)^{m-1}}^q$  is given by

$$\left( \sum_{j_m \in \mathbb{Z}} \int_{\text{SO}(n)} \dots \sum_{j_2 \in \mathbb{Z}} \int_{\text{SO}(n)} \sum_{j_1 \in \mathbb{Z}} \mathcal{S}_{\text{dyad},q}^m f(j_1, \dots, j_m, R_2, \dots, R_m) d\mu_2(R_2) \dots d\mu_m(R_m) \right)^{q/2}.$$

Like before, we will discuss the well-definedness and stability of these operators to diffeomorphisms. The proofs will be omitted since they follow directly from the previous sections with minor modifications.

**Lemma 30.** *Let  $\psi$  be a wavelet that satisfies properties (24) and (25).*

- If  $1 < q \leq 2$ , we have  $\mathcal{S}_{cont,q}^m : \mathbf{L}^q(\mathbb{R}^n) \rightarrow \mathbf{L}^2(\mathbb{R}_+^m) \times SO(n)^{m-1}$  and  $\mathcal{S}_{dyad,q}^m : \mathbf{L}^q(\mathbb{R}^n) \rightarrow \ell^2(\mathbb{Z}^m) \times SO(n)^{m-1}$ .
- If  $q = 1$ , then for every  $\varepsilon > 0$  we have  $\mathcal{S}_{cont,1}^m : \mathbf{H}^1(\mathbb{R}^n) \rightarrow \mathbf{L}_\varepsilon^2(\mathbb{R}_+^m) \times SO(n)^{m-1}$  and  $\mathcal{S}_{dyad,1}^m : \mathbf{H}^1(\mathbb{R}^n) \rightarrow \ell_\varepsilon^2(\mathbb{Z}^m) \times SO(n)^{m-1}$ , with the damped norms of Definition 9; for  $m = 1$  the damping is unnecessary.
- If  $q = 2$  and  $\psi$  is also a Littlewood-Paley wavelet, we have  $\|\mathcal{S}_{cont,2}^m f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times SO(n)^{m-1}}^2 = \mu(SO(n))^{m-1} C_\psi^m \|f\|_2^2$  and  $\|\mathcal{S}_{dyad,2}^m f\|_{\ell^2(\mathbb{Z}^m) \times SO(n)^{m-1}}^2 = \mu(SO(n))^{m-1} \hat{C}_\psi^m \|f\|_2^2$ .

**Theorem 31.** *Assume  $|c| < \frac{1}{2n}$  and  $1 < q < 2$ . Let  $\tau(x) = cx$  and let  $L_\tau f(x) = f((1-c)x)$ . Suppose that  $\psi$  is a wavelet that satisfies the conditions of Lemma 14. Then there exist constants  $\hat{K}_{n,m,q}$  and  $\hat{K}'_{n,m,q}$  dependent only on  $n$ ,  $m$ , and  $q$  such that*

$$\|\mathcal{S}_{cont,q}^m f - \mathcal{S}_{cont,q}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times SO(n)^{m-1}}^q \leq |c|^q \cdot \hat{K}_{n,m,q} \|f\|_q^q$$

and

$$\|\mathcal{S}_{dyad,q}^m f - \mathcal{S}_{dyad,q}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m) \times SO(n)^{m-1}}^q \leq |c|^q \cdot \hat{K}'_{n,m,q} \|f\|_q^q.$$

Additionally, if  $q = 1$  and one of the following holds:

- $n = 1$ ,  $\psi$  is complex analytic and satisfies the conditions of Lemma 14,
- $n \geq 2$  and  $\psi$  satisfies the conditions of Lemma 5,

then for every  $\varepsilon > 0$  there exist  $\hat{H}_{m,n}$  and  $\hat{H}'_{m,n}$ , depending also on  $\varepsilon$ , such that

$$\|\mathcal{S}_{cont,1}^m f - \mathcal{S}_{cont,1}^m L_\tau f\|_{\mathbf{L}_\varepsilon^2(\mathbb{R}_+^m) \times SO(n)^{m-1}} \leq |c| \cdot \hat{H}_{m,n} \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}$$

and

$$\|\mathcal{S}_{dyad,1}^m f - \mathcal{S}_{dyad,1}^m L_\tau f\|_{\ell_\varepsilon^2(\mathbb{Z}^m) \times SO(n)^{m-1}} \leq |c| \cdot \hat{H}'_{m,n} \|f\|_{\mathbf{H}^1(\mathbb{R}^n)}.$$

**Theorem 32.** *Let  $\tau \in C^2(\mathbb{R}^n)$  and define  $L_\tau f(x) = f(x - \tau(x))$  with  $\|D\tau\|_\infty < \frac{1}{2n}$ . Suppose that  $\psi$  is a wavelet such that the wavelet and all its first and second partial derivatives have  $O((1+|x|)^{-n-3})$  decay. For  $q \in (1, 2]$ , there exist constants  $C_{m,n}$ ,  $\hat{C}_{m,n}$ ,  $C_{m,n,q}$ , and  $\hat{C}_{m,n,q}$  such that*

$$\begin{aligned} \|\mathcal{S}_{cont,2}^m f - \mathcal{S}_{cont,2}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times SO(n)^{m-1}}^2 &\leq C_{m,n} \left( \|D\tau\|_\infty^2 + \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \right) \|f\|_2^2, \\ \|\mathcal{S}_{dyad,2}^m f - \mathcal{S}_{dyad,2}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m) \times SO(n)^{m-1}}^2 &\leq \hat{C}_{m,n} \left( \|D\tau\|_\infty^2 + \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^2 \right) \|f\|_2^2, \\ \|\mathcal{S}_{cont,q}^m f - \mathcal{S}_{cont,q}^m L_\tau f\|_{\mathbf{L}^2(\mathbb{R}_+^m) \times SO(n)^{m-1}}^q &\leq C_{m,n,q} \left[ \|D\tau\|_\infty^q + \eta^q M_r^{q\delta} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^{q(1-\delta)} \right] \|f\|_q^q, \\ \|\mathcal{S}_{dyad,q}^m f - \mathcal{S}_{dyad,q}^m L_\tau f\|_{\ell^2(\mathbb{Z}^m) \times SO(n)^{m-1}}^q &\leq \hat{C}_{m,n,q} \left[ \|D\tau\|_\infty^q + \eta^q M_r^{q\delta} \left( \|D\tau\|_\infty \left( \log \frac{\|\Delta\tau\|_\infty}{\|D\tau\|_\infty} \vee 1 \right) + \|D^2\tau\|_\infty \right)^{q(1-\delta)} \right] \|f\|_q^q. \end{aligned}$$

## 7. Conclusion

For  $q = 1$  and  $m \geq 2$ , we have seen that the scale measure must be damped: the input to each layer past the first is nonnegative with positive integral, hence carries no cancellation, and the undamped norm is infinite. We have formulated operators that are translation invariant in  $\mathbf{L}^q(\mathbb{R}^n)$ , proven these operators are Lipschitz continuous to the action of  $C^2$  diffeomorphisms when  $1 < q \leq 2$  with respect to certain norms, and used these results to formulate rotation invariant/equivariant operators on  $\mathbf{L}^q(\mathbb{R}^n)$  that are Lipschitz continuous to the action of  $C^2$  diffeomorphisms.

## 8. Acknowledgments

We would like to thank Michael Perlmutter for providing suggestions that improved the clarity of our initial draft. We would also like to thank the anonymous reviewers for their careful feedback over multiple iterations. Lastly, we would like to thank Yang Yang for checking over a revision to the proof of Proposition 15. Claude Opus 5.0 was used for a correction of the proofs for the norm when  $q = 1$  without a damping measure, which were originally "true" based on a false proof. The authors have checked the output and take responsibility for all results.

## References

- [1] A. Krizhevsky, I. Sutskever, G. E. Hinton, [Imagenet classification with deep convolutional neural networks](#), Communications of the ACM 60 (6) (2017) 84–90.
- [2] K. Simonyan, A. Zisserman, Very deep convolutional networks for large-scale image recognition, in: International Conference on Learning Representations, 2015.
- [3] C. Szegedy, W. Liu, Y. Jia, P. Sermanet, S. Reed, D. Anguelov, D. Erhan, V. Vanhoucke, A. Rabinovich, [Going deeper with convolutions](#), in: Proceedings of the IEEE conference on computer vision and pattern recognition, 2015, pp. 1–9.
- [4] K. He, X. Zhang, S. Ren, J. Sun, [Deep residual learning for image recognition](#), in: Proceedings of the IEEE conference on computer vision and pattern recognition, 2016, pp. 770–778.
- [5] S. Mallat, [Group invariant scattering](#), Communications on Pure and Applied Mathematics 65 (10) (2012) 1331–1398.
- [6] F. Gao, G. Wolf, M. Hirn, [Geometric scattering for graph data analysis](#), in: International Conference on Machine Learning, PMLR, 2019, pp. 2122–2131.
- [7] M. Perlmutter, F. Gao, G. Wolf, M. Hirn, [Geometric wavelet scattering networks on compact Riemannian manifolds](#), in: Mathematical and Scientific Machine Learning, PMLR, 2020, pp. 570–604.
- [8] M. Hirn, S. Mallat, N. Poilvert, [Wavelet scattering regression of quantum chemical energies](#), Multiscale Modeling & Simulation 15 (2) (2017) 827–863.
- [9] M. Eickenberg, G. Exarchakis, M. Hirn, S. Mallat, L. Thiry, [Solid harmonic wavelet scattering for predictions of molecule properties](#), The Journal of chemical physics 148 (24) (2018) 241732.
- [10] P. Sinz, M. W. Swift, X. Brumwell, J. Liu, K. J. Kim, Y. Qi, M. Hirn, Wavelet scattering networks for atomistic systems with extrapolation of material properties, The Journal of Chemical Physics 153 (8) (2020) 084109.
- [11] I. Waldspurger, Exponential decay of scattering coefficients, in: 2017 international conference on sampling theory and applications (SampTA), IEEE, 2017, pp. 143–146.
- [12] F. Gama, A. Ribeiro, J. Bruna, [Stability of graph scattering transforms](#), Advances in Neural Information Processing Systems 32.
- [13] F. Nicola, S. I. Trapasso, Stability of the scattering transform for deformations with minimal regularity, arXiv preprint arXiv:2205.11142.
- [14] T. Wiatowski, H. Bölcskei, [A mathematical theory of deep convolutional neural networks for feature extraction](#), IEEE Transactions on Information Theory 64 (3) (2017) 1845–1866.
- [15] M. Koller, J. Großmann, U. Monich, H. Boche, [Deformation stability of deep convolutional neural networks on Sobolev spaces](#), in: 2018 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP), IEEE, 2018, pp. 6872–6876.
- [16] W. Czaja, W. Li, [Analysis of time-frequency scattering transforms](#), Applied and Computational Harmonic Analysis 47 (1) (2019) 149–171.
- [17] A. Bietti, J. Mairal, [Group invariance, stability to deformations, and complexity of deep convolutional representations](#), The Journal of Machine Learning Research 20 (1) (2019) 876–924.
- [18] E. Cancès, M. Defranceschi, W. Kutzelnigg, C. Le Bris, Y. Maday, Computational quantum chemistry: A primer, in: Special Volume, Computational Chemistry, Vol. 10 of Handbook of Numerical Analysis, Elsevier, 2003, pp. 3–270. doi:[https://doi.org/10.1016/S1570-8659\(03\)10003-8](https://doi.org/10.1016/S1570-8659(03)10003-8).
- [19] J. Bruna, S. Mallat, Audio texture synthesis with scattering moments (2013). [arXiv:1311.0407](#).
- [20] E. Allys, F. Levrier, S. Zhang, C. Colling, B. Regalado-Saint Blancard, F. Boulanger, P. Hennebelle, S. Mallat, The rwst, a comprehensive statistical description of the non-gaussian structures in the ism, Astronomy & Astrophysics 629 (2019) A115.
- [21] L. Grafakos, [Modern fourier analysis](#), Vol. 250, Springer, 2009.
- [22] L. Grafakos, [Classical fourier analysis](#), Vol. 249, Springer, 2014.

- [23] J. García-Cuerva, J. R. De Francia, [Weighted norm inequalities and related topics](#), Elsevier, 1985.
- [24] J. P. Ward, K. N. Chaudhury, M. Unser, Decay properties of riesz transforms and steerable wavelets, *SIAM Journal on Imaging Sciences* 6 (2) (2013) 984–998. [doi:10.1137/120864143](#).
- [25] G. H. Hardy, J. E. Littlewood, G. Pólya, *Inequalities*, Cambridge University Press, Cambridge, 1988.
- [26] E. M. Stein, T. S. Murphy, [Harmonic Analysis \(PMS-43\): Real-Variable Methods, Orthogonality, and Oscillatory Integrals](#), Princeton University Press, 1993.
- [27] R. P. Brent, J. anne H. Osborn, W. D. Smith, Note on best possible bounds for determinants of matrices close to the identity matrix, *Linear Algebra and its Applications* 466 (2015) 21–26. [doi:https://doi.org/10.1016/j.laa.2014.09.041](#).