

Relation: Part 1 - Definition

Adila A. Krisnadhi

Faculty of Computer Science, Universitas Indonesia

Reference: Rosen, *Discrete Mathematics and Its Applications*, 8ed, 2019, Sec. 9.1 - 9.2

What is relation?

Intuition: relation \rightsquigarrow association between several objects.

What is relation?

Intuition: relation \rightsquigarrow association between several objects.

- `hasParent` between someone and his/her parent;
`hasParent(Ilham, Rudi)`, `hasParent(Ilham, Ainun)`.

What is relation?

Intuition: relation \rightsquigarrow association between several objects.

- **hasParent** between someone and his/her parent;
`hasParent(Ilham, Rudi)`, `hasParent(Ilham, Ainun)`.
- **divisibleBy** between an integer and an(other) integer that divides it;
`divisibleBy(12,4)`, `divisibleBy(0,7)`.

What is relation?

Intuition: relation \rightsquigarrow association between several objects.

- **hasParent** between someone and his/her parent;
`hasParent(Ilham, Rudi)`, `hasParent(Ilham, Ainun)`.
- **divisibleBy** between an integer and an(other) integer that divides it;
`divisibleBy(12,4)`, `divisibleBy(0,7)`.
- **Arrival** between a flight, airline, origin airport, landing date, landing time, and debarking terminal;
`Arrival(QZ 0691, Air Asia, SUB, 2020-03-20, 13:00, 2E)`,
`Arrival(ID 6519, Batik Air, DPS, 2020-03-20, 09:30, 2E)`.

Tuple

The pair $(12, 4)$ in `divisibleBy(12, 4)` represents the objects for which the relation holds. Such a pair is called a tuple.

Tuple

The pair $(12, 4)$ in $\text{divisibleBy}(12, 4)$ represents the objects for which the relation holds. Such a pair is called a tuple.

Definition (n -tuple)

For every $n \in \mathbb{N}$, an **n -tuple** is a sequence (or ordered list) containing n elements a_1, a_2, \dots, a_n written using the notation: (a_1, a_2, \dots, a_n)

Tuple

The pair $(12, 4)$ in $\text{divisibleBy}(12, 4)$ represents the objects for which the relation holds. Such a pair is called a tuple.

Definition (n -tuple)

For every $n \in \mathbb{N}$, an **n -tuple** is a sequence (or ordered list) containing n elements a_1, a_2, \dots, a_n written using the notation: (a_1, a_2, \dots, a_n)

- There exists exactly one 0-tuple, namely the **empty tuple** $()$.

Tuple

The pair $(12, 4)$ in $\text{divisibleBy}(12, 4)$ represents the objects for which the relation holds. Such a pair is called a tuple.

Definition (n -tuple)

For every $n \in \mathbb{N}$, an **n -tuple** is a sequence (or ordered list) containing n elements a_1, a_2, \dots, a_n written using the notation: (a_1, a_2, \dots, a_n)

- There exists exactly one 0-tuple, namely the **empty tuple** $()$.
- 1-tuples: (1) , (a) , (January) . A 1-tuple is usually written without parentheses, e.g., 1 , a , January .

Tuple

The pair $(12, 4)$ in $\text{divisibleBy}(12, 4)$ represents the objects for which the relation holds. Such a pair is called a tuple.

Definition (n -tuple)

For every $n \in \mathbb{N}$, an **n -tuple** is a sequence (or ordered list) containing n elements a_1, a_2, \dots, a_n written using the notation: (a_1, a_2, \dots, a_n)

- There exists exactly one 0-tuple, namely the **empty tuple** $()$.
- 1-tuples: (1) , (a) , (January) . A 1-tuple is usually written without parentheses, e.g., 1 , a , January .
- 2-tuples or **(ordered) pairs**, e.g., $(3, 3)$, $(1, 2)$, $(2, \text{adila})$, $(\text{Fasilkom}, \text{UI})$

Tuple

The pair $(12, 4)$ in $\text{divisibleBy}(12, 4)$ represents the objects for which the relation holds. Such a pair is called a tuple.

Definition (n -tuple)

For every $n \in \mathbb{N}$, an **n -tuple** is a sequence (or ordered list) containing n elements a_1, a_2, \dots, a_n written using the notation: (a_1, a_2, \dots, a_n)

- There exists exactly one 0-tuple, namely the **empty tuple** $()$.
- 1-tuples: (1) , (a) , (January) . A 1-tuple is usually written without parentheses, e.g., 1 , a , January .
- 2-tuples or **(ordered) pairs**, e.g., $(3, 3)$, $(1, 2)$, $(2, \text{adila})$, $(\text{Fasilkom}, \text{UI})$
- 3-tuples or **triples**, e.g., $(3, 3, a)$, $(0, 1, 2)$, $(\text{adila}, \text{UI}, \text{depok})$

Cartesian product

Definition

The **Cartesian product** of n sets A_1, A_2, \dots, A_n is the following set

$$A_1 \times A_2 \times \dots \times A_n = \{(c_1, c_2, \dots, c_n) \mid c_i \in A_i \text{ for each } i = 1, 2, \dots, n\}$$

The Cartesian product $\underbrace{A \times A \times \dots \times A}_{n \text{ times}}$ is often written A^n .

Cartesian product

Definition

The **Cartesian product** of n sets A_1, A_2, \dots, A_n is the following set

$$A_1 \times A_2 \times \dots \times A_n = \{(c_1, c_2, \dots, c_n) \mid c_i \in A_i \text{ for each } i = 1, 2, \dots, n\}$$

The Cartesian product $\underbrace{A \times A \times \dots \times A}_{n \text{ times}}$ is often written A^n .

Suppose $A = \{1, 2, 3\}$ and $B = \{a, b\}$. Then:

- $A \times B =$
- $B \times B \times A =$

Cartesian product

Definition

The **Cartesian product** of n sets A_1, A_2, \dots, A_n is the following set

$$A_1 \times A_2 \times \dots \times A_n = \{(c_1, c_2, \dots, c_n) \mid c_i \in A_i \text{ for each } i = 1, 2, \dots, n\}$$

The Cartesian product $\underbrace{A \times A \times \dots \times A}_{n \text{ times}}$ is often written A^n .

Suppose $A = \{1, 2, 3\}$ and $B = \{a, b\}$. Then:

- $A \times B = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b)\}$
- $B \times B \times A =$

Cartesian product

Definition

The **Cartesian product** of n sets A_1, A_2, \dots, A_n is the following set

$$A_1 \times A_2 \times \dots \times A_n = \{(c_1, c_2, \dots, c_n) \mid c_i \in A_i \text{ for each } i = 1, 2, \dots, n\}$$

The Cartesian product $\underbrace{A \times A \times \dots \times A}_{n \text{ times}}$ is often written A^n .

Suppose $A = \{1, 2, 3\}$ and $B = \{a, b\}$. Then:

- $A \times B = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b)\}$
- $B \times B \times A = \{(a, a, 1), (a, a, 2), (a, a, 3), (a, b, 1), (a, b, 2), (a, b, 3), (b, a, 1), (b, a, 2), (b, a, 3), (b, b, 1), (b, b, 2), (b, b, 3)\}$

Suppose $A = \{1, 2, 3\}$, $B = \{a, b\}$, $C = \{0, 1\}$, and $D = \emptyset$. Then:

- $B \times A =$

- $B \times B =$

- $B \times C \times A =$

- $A \times B \times C \times D =$

Definition

Let A_1, A_2, \dots, A_n be sets.

- An **n -ary relation** R over A_1, \dots, A_n is a subset of the Cartesian product $A_1 \times \dots \times A_n$
- We write the above n -ary relation $R \subseteq A_1 \times \dots \times A_n$.
- The sets A_1, \dots, A_n are the **domain** of R and n is the **arity** of R .
- A tuple (a_1, \dots, a_n) is an **instance** of an n -ary relation R iff $(a_1, \dots, a_n) \in R$.
 - For the above, we usually write $R(a_1, \dots, a_n)$.
 - If R is binary ($n = 2$), an instance $(a, b) \in R$ is also written $a R b$.

Relation

- Can a relation be empty?
- Can a relation be infinite?

Relation

- Can a relation be empty?
Yes:
- Can a relation be infinite?

Relation

- Can a relation be empty?

Yes: \emptyset is also a subset of the Cartesian product.

- Can a relation be infinite?

Relation

- Can a relation be empty?

Yes: \emptyset is also a subset of the Cartesian product.

- Can a relation be infinite?

Yes:

Relation

- Can a relation be empty?

Yes: \emptyset is also a subset of the Cartesian product.

- Can a relation be infinite?

Yes: If any of domain set is infinite, a subset of the Cartesian product may be infinite.

Relation

Definition 3 defines n -ary relation for $n = 0, 1, 2, \dots$,

Relation

Definition 3 defines n -ary relation for $n = 0, 1, 2, \dots$,

- $n = 0$: the trivial **nullary** relation $\{()\}$ containing only a single empty tuple.

Relation

Definition 3 defines n -ary relation for $n = 0, 1, 2, \dots$,

- $n = 0$: the trivial **nullary** relation $\{()\}$ containing only a single empty tuple.
- $n = 1$: **unary** relations, equivalent to sets of non-tuple objects
 - E.g.: $R = \{(0), (1), (2)\} = \{0, 1, 2\}$.

Relation

Definition 3 defines n -ary relation for $n = 0, 1, 2, \dots$,

- $n = 0$: the trivial **nullary** relation $\{()\}$ containing only a single empty tuple.
- $n = 1$: **unary** relations, equivalent to sets of non-tuple objects
 - E.g.: $R = \{(0), (1), (2)\} = \{0, 1, 2\}$.
- $n = 2$: **binary** relations, i.e., sets of pairs \rightsquigarrow **focus of this chapter**.
 - A binary relation $R \subseteq A \times B$ is also called a **binary relation from A to B** .
 - If $A = B$, then $R \subseteq A \times A$ is also called a **(binary) relation over A**

Relation

Definition 3 defines n -ary relation for $n = 0, 1, 2, \dots$,

- $n = 0$: the trivial **nullary** relation $\{()\}$ containing only a single empty tuple.
- $n = 1$: **unary** relations, equivalent to sets of non-tuple objects
 - E.g.: $R = \{(0), (1), (2)\} = \{0, 1, 2\}$.
- $n = 2$: **binary** relations, i.e., sets of pairs \rightsquigarrow **focus of this chapter**.
 - A binary relation $R \subseteq A \times B$ is also called a **binary relation from A to B** .
 - If $A = B$, then $R \subseteq A \times A$ is also called a **(binary) relation over A**
- $n = 3$: **ternary** relations, i.e., sets of triples.

Relation

Definition 3 defines n -ary relation for $n = 0, 1, 2, \dots$,

- $n = 0$: the trivial **nullary** relation $\{()\}$ containing only a single empty tuple.
- $n = 1$: **unary** relations, equivalent to sets of non-tuple objects
 - E.g.: $R = \{(0), (1), (2)\} = \{0, 1, 2\}$.
- $n = 2$: **binary** relations, i.e., sets of pairs \rightsquigarrow **focus of this chapter**.
 - A binary relation $R \subseteq A \times B$ is also called a **binary relation from A to B** .
 - If $A = B$, then $R \subseteq A \times A$ is also called a **(binary) relation over A**
- $n = 3$: **ternary** relations, i.e., sets of triples.
- $n > 3$: sets of n -tuples.

Relation

Definition 3 defines n -ary relation for $n = 0, 1, 2, \dots$,

- $n = 0$: the trivial **nullary** relation $\{()\}$ containing only a single empty tuple.
- $n = 1$: **unary** relations, equivalent to sets of non-tuple objects
 - E.g.: $R = \{(0), (1), (2)\} = \{0, 1, 2\}$.
- $n = 2$: **binary** relations, i.e., sets of pairs \rightsquigarrow **focus of this chapter**.
 - A binary relation $R \subseteq A \times B$ is also called a **binary relation from A to B** .
 - If $A = B$, then $R \subseteq A \times A$ is also called a **(binary) relation over A**
- $n = 3$: **ternary** relations, i.e., sets of triples.
- $n > 3$: sets of n -tuples.
- General n -ary relations are used in relational databases studied in a later course.

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

Give examples of pair in $A \times B$ that do not belong to R .

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

- A contains cities, e.g.,

Give examples of pair in $A \times B$ that do not belong to R .

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

- A contains 34 cities, e.g., Cilegon, Jakarta Pusat, Depok, Bandung, Semarang, Magelang, Surabaya, Malang, etc.

Give examples of pair in $A \times B$ that do not belong to R .

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

- A contains 34 cities, e.g., Cilegon, Jakarta Pusat, Depok, Bandung, Semarang, Magelang, Surabaya, Malang, etc.
- B contains provinces, namely

Give examples of pair in $A \times B$ that do not belong to R .

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

- A contains 34 cities, e.g., Cilegon, Jakarta Pusat, Depok, Bandung, Semarang, Magelang, Surabaya, Malang, etc.
- B contains 6 provinces, namely Banten, DKI Jakarta, Jawa Barat, Jawa Tengah, DI Yogyakarta, and Jawa Timur.

Give examples of pair in $A \times B$ that do not belong to R .

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

- A contains 34 cities, e.g., Cilegon, Jakarta Pusat, Depok, Bandung, Semarang, Magelang, Surabaya, Malang, etc.
- B contains 6 provinces, namely Banten, DKI Jakarta, Jawa Barat, Jawa Tengah, DI Yogyakarta, and Jawa Timur.
- $A \times B$ contains pairs.

Give examples of pair in $A \times B$ that do not belong to R .

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

- A contains 34 cities, e.g., Cilegon, Jakarta Pusat, Depok, Bandung, Semarang, Magelang, Surabaya, Malang, etc.
- B contains 6 provinces, namely Banten, DKI Jakarta, Jawa Barat, Jawa Tengah, DI Yogyakarta, and Jawa Timur.
- $A \times B$ contains $34 \times 6 = 204$ pairs.

Give examples of pair in $A \times B$ that do not belong to R .

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

- A contains 34 cities, e.g., Cilegon, Jakarta Pusat, Depok, Bandung, Semarang, Magelang, Surabaya, Malang, etc.
- B contains 6 provinces, namely Banten, DKI Jakarta, Jawa Barat, Jawa Tengah, DI Yogyakarta, and Jawa Timur.
- $A \times B$ contains $34 \times 6 = 204$ pairs.
- R contains pairs e.g.,

Give examples of pair in $A \times B$ that do not belong to R .

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

- A contains 34 cities, e.g., Cilegon, Jakarta Pusat, Depok, Bandung, Semarang, Magelang, Surabaya, Malang, etc.
- B contains 6 provinces, namely Banten, DKI Jakarta, Jawa Barat, Jawa Tengah, DI Yogyakarta, and Jawa Timur.
- $A \times B$ contains $34 \times 6 = 204$ pairs.
- R contains 34 pairs e.g., (Cilegon, Banten), (Cirebon, Jawa Barat), (Tegal, Jawa Tengah), (Yogyakarta, DI Yogyakarta), (Malang, Jawa Timur), etc.

Give examples of pair in $A \times B$ that do not belong to R .

Let A be the set of cities/town in Java headed by a major and B the set of Indonesian provinces in Java. Define the relation R such that $(a, b) \in R$ if city/town a is located in the province b .

- A contains 34 cities, e.g., Cilegon, Jakarta Pusat, Depok, Bandung, Semarang, Magelang, Surabaya, Malang, etc.
- B contains 6 provinces, namely Banten, DKI Jakarta, Jawa Barat, Jawa Tengah, DI Yogyakarta, and Jawa Timur.
- $A \times B$ contains $34 \times 6 = 204$ pairs.
- R contains 34 pairs e.g., (Cilegon, Banten), (Cirebon, Jawa Barat), (Tegal, Jawa Tengah), (Yogyakarta, DI Yogyakarta), (Malang, Jawa Timur), etc.

Give examples of pair in $A \times B$ that do not belong to R .

- (Bandung, Jawa Tengah), (Madiun, Jawa Barat), etc.

Exercise

- ① Let $A = \{0, 1, 2\}$ and $B = \{a, b\}$. Give two examples of binary relation R from A to B . How many such relations are possible?
- ② Check if **all** of $(1, 1), (1, 2), (2, 1), (1, -1), (2, 2)$ are in the relations below.
 - $R_1 = \{(a, b) \in \mathbb{Z}^2 \mid a \leq b\}$
 - $R_2 = \{(a, b) \in \mathbb{Z}^2 \mid a > b\}$
 - $R_3 = \{(a, b) \in \mathbb{Z}^2 \mid a = b \text{ or } a = -b\}$
 - $R_4 = \{(a, b) \in \mathbb{Z}^2 \mid a = b + 1\}$
 - $R_5 = \{(a, b) \in \mathbb{Z}^2 \mid a + b \leq 3\}$
- ③ For each of the following ternary relations, write its degree and domains, and give two triples that belong to it and two that do not belong to it.
 - $R_1 = \{(a, b, c) \in \mathbb{N}^3 \mid a < b < c\}$
 - $R_2 = \{(a, b, c) \in \mathbb{Z}^3 \mid b = a + k \text{ and } c = a + 2k \text{ for some } k \in \mathbb{Z}\}$
 - $R_3 = \{(a, b, c) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}^+ \mid a \equiv b \pmod{c}\}$

Let $A = \{0, 1, 2\}$ and $B = \{a, b\}$. Give two examples of binary relation R from A to B . How many such relations are possible?

Check if **all** of $(1, 1), (1, 2), (2, 1), (1, -1), (2, 2)$ are in the relations below.

$$R_1 = \{(a, b) \in \mathbb{Z}^2 \mid a \leq b\}$$

☐ Yes

☐ No

$$R_2 = \{(a, b) \in \mathbb{Z}^2 \mid a > b\}$$

☐ Yes

☐ No

$$R_3 = \{(a, b) \in \mathbb{Z}^2 \mid a = b \text{ or } a = -b\}$$

☐ Yes

☐ No

$$R_4 = \{(a, b) \in \mathbb{Z}^2 \mid a = b + 1\}$$

☐ Yes

☐ No

$$R_5 = \{(a, b) \in \mathbb{Z}^2 \mid a + b \leq 3\}$$

☐ Yes

☐ No

For each of the following ternary relations, write its domains, and give two triples that belong to it and two that do not belong to it.

$$R_1 = \{(a, b, c) \in \mathbb{N}^3 \mid a < b < c\}$$

$$R_2 = \{(a, b, c) \in \mathbb{Z}^3 \mid b = a + k \text{ and } c = a + 2k \text{ for some } k \in \mathbb{Z}\}$$

$$R_3 = \{(a, b, c) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}^+ \mid a \equiv b \pmod{c}\}$$