

677 Homework March 8

Albert Ding

March 7, 2019

1. There is insufficient evidence to show that the distribution is not uniform given the p-value of 0.3501.

```
maybe_uni<-read.table("maybe_uniform.txt", header = FALSE)
```

```
## Warning in read.table("maybe_uniform.txt", header = FALSE): incomplete
## final line found by readTableHeader on 'maybe_uniform.txt'
```

```
ks.test(maybe_uni, "punif")
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data: maybe_uni
## D = 0.18, p-value = 0.3501
## alternative hypothesis: two-sided
```

2. At alpha level 0.05 we can reject the null value that the distribution is drawn from the normal distribution.

```
maybe_norm<-read.table("maybe_normal.txt", header = FALSE)
ks.test(maybe_norm, "pnorm", 26, 4)
```

```
##
## One-sample Kolmogorov-Smirnov test
##
## data: maybe_norm
## D = 0.19311, p-value = 0.04157
## alternative hypothesis: two-sided
```

3. There is insufficient evidence to reject the null that the samples are drawn from the same distribution.

```
maybe1<-c(0.61, 0.29, 0.06, 0.59, -1.73, -0.74, 0.51, -0.56, -0.39, 1.64, 0.05, -0.06, 0.64, -0.82, 0.3
maybe2<-c(2.20, 1.66, 1.38, 0.20, 0.36, 0.00, 0.96, 1.56, 0.44, 1.50, -0.30, 0.66, 2.31, 3.29, -0.27, -
ks.test(maybe1, maybe2)
```

```
##
## Two-sample Kolmogorov-Smirnov test
##
## data: maybe1 and maybe2
## D = 0.27, p-value = 0.3357
## alternative hypothesis: two-sided
```

4. There is insufficient evidence to reject the null that the samples are drawn from the same distribution. D is equal to 0.177.

```
normsamp<-readRDS("norm_sample.Rdata")  
ks.test(normsamp, "pnorm",0,1)
```

```
##  
## One-sample Kolmogorov-Smirnov test  
##  
## data:  normsamp  
## D = 0.17724, p-value = 0.3683  
## alternative hypothesis: two-sided
```

```
ecdf(normsamp)
```

```
## Empirical CDF  
## Call: ecdf(normsamp)  
## x[1:25] = -2.46, -2.11, -1.23, ..., 1.64, 1.76
```

```
pnorm(0,1)
```

```
## [1] 0.1586553
```

5. Not sure about this one.