


AUTHOR QUERY FORM

 ELSEVIER	Book: Advances in Quantum Chemistry Article Number: 132	Please e-mail your responses and any corrections to: E-mail: d.mansingh@elsevier.com
---	--	--

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

We were unable to process your file(s) fully electronically and have proceeded by

☐

Scanning (parts
of) your article

☐

Rekeying (parts
of) your article

☐

Scanning the
artwork

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the ‘[Q](#)’ link to go to the location in the proof.

Location in Chapter	Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof		
Q1	<p>We have not received any alternate running heading for this chapter. Hence we have retained the existing running head “Collision processes in atoms and molecules” as it is now. Please check and suggest an alternate running head for this chapter.</p> <table border="1" data-bbox="413 1503 1076 1580"> <tr> <td data-bbox="413 1503 1009 1580">Please check this box or indicate your approval if you have no corrections to make to the PDF file</td><td data-bbox="1009 1503 1076 1580"></td></tr> </table>	Please check this box or indicate your approval if you have no corrections to make to the PDF file	
Please check this box or indicate your approval if you have no corrections to make to the PDF file			

Thank you for your assistance.

C0660

Collision processes in atoms and molecules using effective potentials



Alejandra M.P. Mendez*, Darío M. Mitnik, Jorge E. Miraglia

Instituto de Astronomía y Física del Espacio, Universidad de Buenos Aires – Consejo Nacional de Investigaciones Científicas y Técnicas, Buenos Aires, Argentina

*Corresponding author: e-mail address: alemendez@iafe.uba.ar

Contents

1. Outline	2
2. Theory	3
2.1 Pseudopotential approximation	3
2.2 Depurated inversion method potentials	7
3. Collisional processes in atoms	8
3.1 Proton-impact excitation	8
3.2 Proton-impact ionization	10
3.3 Proton-impact charge exchange	11
3.4 Photoionization	12
3.5 DIM Photoionization of many-electron atoms	13
4. Depurated inversion method for molecules	13
4.1 Theory	14
4.2 Example: Methane	15
4.3 Collisional processes	17
5. Concluding remarks	19
Acknowledgments	19
References	20

Abstract

We investigate the feasibility of using pseudopotentials to generate the bound and continuum orbitals needed in collisional calculations. By examination of several inelastic processes in the first Born approximation, we demonstrate the inconveniences of this approach. Instead, we advocate use of effective potentials obtained with the depurated inversion method (DIM). In this contribution, we extend this method to molecular systems. Calculations of single first-order photoionization and proton-impact ionization using the DIM show fair agreement with experimental results for both atoms and molecules.



1. Outline

s0010

p0010

Inelastic transition calculations require the representation of the bound and continuum states involved in the collisional processes. The hypothetical existence of an effective one-electron local potential accounting for these states would allow more direct generation of the orthogonal wavefunctions for the interacting particles. This approach should include individual nl -orbital potentials, a feature missing from most of the standard density functional methods. The idea of replacing a many-body, nonlocal interaction by an effective one-electron equation opens up the possibility of studying extremely complex systems with high accuracy.

p0015

In this context, one promising idea emerges from the pseudopotential approximation (PPA), in which all the complexity of the wavefunctions near the core—that usually consumes a huge numerical effort—is avoided. For instance, density functional theory codes using pseudopotentials, such as the PARSEC, for example,^{1,2} permit the use of an equally spaced grid involving a relatively small number of points. Otherwise, the use of realistic potentials describing the nucleus Coulomb potential requires a high density of points concentrated at the origin to describe what the pseudopotentials cast aside. Thus, if PPA was applicable in the field of collision theory, one would save an enormous amount of computational resources.

p0020

Another interesting approach is the depurated inversion method (DIM),^{3–5} which allows accurate, effective potentials to be obtained by substituting the coupled multielectron equations into a Kohn–Sham-type equation. In the first step, the potential is obtained through inversion of the one-electron equation. Next, a careful optimization of the potential is carried out, eliminating poles, and imposing the appropriate boundary conditions analytically. In that way, the DIM potentials are parametrized in simple analytical expressions.

p0025

In the present work, we explore the possibility of implementing an effective potential approximation in the atomic collision theory to describe inelastic processes. In particular, we examine several collisional processes involving a single electron transition: photoionization, excitation, ionization, and electron capture. A wide variety of *ab initio* methods have been implemented to compute scattering cross sections for atomic targets, from the early implementations of the first Born approximation (FBA),^{6,7} to more sophisticated fully quantum mechanical methods, e.g., Refs. 8–11. Whether for atoms or molecules, we shall present cross sections and compare with some

experimental data. We do not wish, here, to present a detailed comparison with existing calculations. The main purpose is to illustrate the effective use of the DIM in collision applications. To this end, several simplifications are made (1) The calculations are constrained to Hamiltonians describing only the moving projectile, the target, and the active electron; (2) The transition-matrix elements are only considered in first perturbative order. If the first-order fails, it would not make any sense to extend the calculation to higher terms of the series. For simplicity, we will restrict our calculations only to the FBA framework, which is known to give reasonable agreement with the experimental cross section in the intermediate—high projectile energy range. Moreover, within this energy range and approximation order, the Hartree–Fock orbitals are known to provide the correct high energy limit.

p0030 We examine the above mentioned inelastic processes for two atoms with a single outer electron: hydrogen and lithium. In this context, we inspect the influence of the target description in the cross sections when the PPA and DIM approaches are implemented. Furthermore, these effects have been previously studied in other perturbative approaches, i.e., the continuum distorted wave eikonal-initial-state (CDW-EIS), for various targets (for example, see Refs. 12, 13). The DIM approach is further tested in the case of many-electron atoms by comparing photoionization cross sections with experimental measurements.

p0035 On the other hand, the description of molecular systems constitutes a real challenge due to their nonspherical symmetry and multicenter character. Many ab initio and semiempirical theoretical approximations^{14–16} have been developed to this end over the last century. In this work, we present an extension of the DIM method for simple molecular systems, providing a new parametric expression for the potentials. The target description is once again tested by examination of its performance in first-order collisional processes, and the methane molecule being taken as an example.

s0015

2. Theory

s0020

2.1 Pseudopotential approximation

p0045 The pseudopotential approximation consists in replacing the Coulomb potential in the many-electron system Hamiltonian with a smooth function so that the electron wavefunctions oscillating rapidly in the core region are replaced by nodeless pseudo-orbitals having the right energy and the same

outer range properties. In general, the pseudopotentials V_{PP} can be defined through a pseudocharge Z_{PP} as

$$V_{PP}(r) = -\frac{Z_{PP}(r)}{r}, \quad (1)$$


$$Z_{PP}(r) = \begin{cases} f(r), & r \leq r_c \\ 1, & r > r_c \end{cases}, \quad (2)$$

where r_c is a cutoff radius that separates the core, $r \leq r_c$, from the valence region, $r > r_c$, of the target and $f(r)$ is a continuous function with a constant value at the origin. Fig. 1 illustrates a pseudopotential (solid line) and its corresponding pseudo-wavefunction for the 3s orbital of argon. Notice that the pseudopotential behaves as $-r^{-1}$ (dot-dash line) in the valence region, as defined in Eqs. (1) and (2). The pseudo-wavefunction agrees with the one-electron Hartree–Fock (HF) orbital (dashed line) in the outer region, losing all information about the atomic structure close to the origin.

p0050 In Section 3, we analyse the feasibility of implementing pseudopotentials in collisional processes calculations for two simple atomic targets: hydrogen and lithium. For each atom, the following pseudopotentials are examined

Name	Source	Type	Refs.
<i>A</i>	ABINIT	GGA	17, 18
<i>P</i>	PARSEC	Troullier Martins	1, 2

The hydrogen atom has only one electron, and the corresponding pseudopotential is not essential. However, the hydrogen pseudopotentials from (3) reproduce with high accuracy the main features of the wavefunctions, even for excited states.



f0010 Fig. 1 (A) Pseudopotential, (B) pseudo-wavefunction and HF orbital for the 3s orbital of argon.

p0055 We will now proceed to examine the pseudocharges and its one-electron solutions for the lithium atom closely. First, we study the spatial and momentum representation of the pseudocharges. The momentum-space equivalent of $Z(r)$ is given by the Fourier transform

$$\tilde{Z}(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} Z(r) e^{-ikr} dr. \quad (4)$$


The pseudocharges from (3) for the $2s$ orbital of lithium are illustrated in Fig. 2. For comparison, we include the potential obtained from implementing the depurated inversion method described in Section 2.2. The pseudocharges vanish at the origin, avoiding the divergence of the Coulomb potential. However, this feature comes at a price: the pseudocharges in the spatial representation are repulsive around $r=1$ a.u., and their momentum picture fails to represent the target for high k , showing an incorrect oscillatory behavior for values greater than $k_c = (2\pi r_c)^{-1} \sim 0.7$ a.u..

p0060 Secondly, we inspect the behavior of the bound pseudo-orbitals obtained from solving the one-electron Schrödinger equation with a pseudopotential. As usual, the bound state wavefunctions can be written as

$$\psi_{nlm}(\mathbf{r}) = \frac{u_{nl}(r)}{r} Y_l^m(\hat{r}), \quad (5)$$

where $u_{nl}(r)$ are the reduced radial wavefunctions, and $Y_l^m(\hat{r})$ are the spherical harmonics. Similarly, the Fourier transform of these functions is given by

$$\tilde{\psi}_{nlm}(\mathbf{k}) = \frac{\chi_{nl}(k)}{k} Y_l^m(\hat{k}). \quad (6)$$



f0015 **Fig. 2** Pseudo and DIM charges for the $2s$ orbital of lithium. (A) Spatial and (B) momentum representation.




Fig. 3 Pseudo and DIM bound state wavefunction for the 2s orbital of lithium in (A) spatial and (B) momentum representation.

The spatial and momentum representations of the 2s radial pseudo-wavefunctions of lithium corresponding to the pseudocharges from (3) are displayed in Fig. 3. Although the pseudo-orbitals are very different from the DIM 2s wavefunction, the transformed $\chi(k)$ seems to have similar characteristics. However, a closer inspection of the tail region of these functions (see the inset of the figure) shows the existence of several nodes. We will see later that these discrepancies have significant consequences in the cross sections for most of the collisional processes examined.

Finally, the pseudopotential approach not only affects the representation of the bound orbitals but also determines the form of the continuum wavefunctions. For large r , the free state orbitals of an electron in the presence of a Coulomb potential can be written as

$$u_{kl}(r) \rightarrow \sin\left(kr - l\frac{\pi}{2} - \eta \ln 2kr + \sigma_l + \delta_l\right), \quad (7)$$

where k is the particle wave number, η is Sommerfeld's parameter, σ_l is the Coulomb phase shift, and δ_l is the wave phase shift with respect to the Coulomb wave.

Comparisons between the DIM (solid line) and the pseudo (dashed) continuum ks wavefunctions for lithium are shown in Fig. 4, close to the origin (left) and asymptotically (right). The pseudo and DIM wavefunctions behave similarly away from the nucleus. The asymptotic phase shift Δ accounts for the differences between the potentials. As the energy of the free electron increases, Δ diminishes. However, the orbitals in the core region are different even with increasing energy; the first maximum of the DIM wavefunctions is consistently smaller than of the pseudo-orbitals, which is understood since the Coulomb-type attraction of the nuclei is stronger than the pseudopotential in that region.




Fig. 4 Continuum ks wavefunctions with energies E near the origin (left) and in the asymptotic region (right), calculated with the DIM potential (solid line) and the ABINIT pseudopotential (dashed line).

2.2 Depurated inversion method potentials

The depurated inversion method^{3–5} consists of assuming that the many-electron atom orbitals can be represented by the solution of Kohn–Sham-type equations, in which the nl effective potentials are given by

$$V_{nl}(r) = \frac{1}{2} \frac{1}{u_{nl}(r)} \frac{d^2 u_{nl}(r)}{dr^2} - \frac{l(l+1)}{2r^2} + \varepsilon_{nl}, \quad (8)$$

where u_{nl} and ε_{nl} are the nl orbital wavefunctions and energies, respectively. In this work, the atomic structure is approximated with the Hartree–Fock method, which is computed with the HF codes by C. Froese Fischer¹⁹ and the NRHF code by Johnson.²⁰ The computation of Eq. (8) poses various numerical problems. The nodes and asymptotic decay of the wavefunctions $u_{nl}(r)$ introduce significant numerical errors in the inversion procedure (see Ref. 5 for further details). The nodes of the orbitals produce huge unphysical poles, while the rapid asymptotic decay of the internal wavefunctions generates large divergences in the tail region of the potentials. The depuration method is implemented to tackle these unphysical features. An effective potential with a Coulomb-type shape $V_r(r) = -Z_r(r)/r$ is defined, and we enforce the correct boundary conditions fitting the inverted potential with the following analytical expression

$$Z_r(r) = \sum_{j=1}^n z_j e^{-\alpha_j r} (1 + \beta_j r) + 1 \rightarrow \begin{cases} Z_N, & r \rightarrow 0 \\ 1, & r \rightarrow \infty \end{cases} \quad (9)$$

where $\sum z_j = Z_N - 1$ (Z_N here stands for the nuclear charge). The parameters α_j and β_j are optimized to reproduce the HF values accurately.

s0030

p0090

3. Collisional processes in atoms


The most significant advantage of the pseudopotential method is its simplicity. However, it is worth determining the validity of this approach when used for computing collisional processes. In this section, we perform a thorough examination of the pseudopotentials for hydrogen and lithium by comparing the cross sections of four inelastic processes: proton-impact excitation, proton-impact ionization, charge exchange, and photoionization. The initial and final states of the targets are obtained by solving the corresponding Schrödinger equation. For the hydrogen atom, we compare the pseudopotential results with the exact analytical solutions. Furthermore, in order to assess the applicability of the depurated inversion method, we compute the photoionization of more complex many-electron atoms and compare our findings with experimental data.

s0035

3.1 Proton-impact excitation

p0095

The proton-impact excitation of target X is defined as



The excitation cross section σ of the target from the initial bound state ψ_i to the excited state ψ_f may be written as

$$\sigma = \frac{\mu^2}{4\pi^2} \frac{k_f}{k_i} \int |T_{fi}|^2 d\Omega, \quad (11)$$

where μ is the reduced mass of the proton-atom system, \mathbf{k}_i and \mathbf{k}_f are the initial and final relative momenta, and

$$T_{fi} = \langle \psi_f | V | \psi_i \rangle \quad (12)$$

is the transition-matrix or T-matrix. If the initial and final states of the transition are described by the Hartree-Fock method, the orbitals will give the correct high energy limit in the first-order approximation (this is not the case for the charge exchange process). Hence, we will concentrate our computing efforts in the first perturbative order of the transition-matrix element through the FBA, given by

$$T_{fi}^{\text{FBA}} = \tilde{V}(\mathbf{p}) F_{fi}(\mathbf{p}). \quad (13)$$

The term $F_{fi}(\mathbf{p})$ is the form factor

$$F_{fi}(\mathbf{p}) = \frac{1}{(2\pi)^{3/2}} \int \tilde{\psi}_f^*(\mathbf{k}) \tilde{\psi}_i(\mathbf{k} + \mathbf{p}) d\mathbf{k}, \quad (14)$$

where \mathbf{p} is the momentum transfer vector

$$\mathbf{p} = p_{\min} \hat{\mathbf{v}} + \boldsymbol{\eta}, \quad (15)$$

$$p_{\min} = \frac{\varepsilon_f - \varepsilon_i}{v} \rightarrow \begin{cases} \infty, & v \rightarrow 0 \\ 0, & v \rightarrow \infty \end{cases}, \quad (16)$$

$\hat{\mathbf{v}}$ is the ion velocity, $\boldsymbol{\eta}$ is the transversal momentum transfer, so that $\hat{\mathbf{v}} \cdot \boldsymbol{\eta} = 0$, whereas ε_i and ε_f are the binding energies corresponding to the initial and final states. A more comprehensive formulation of the FBA can be found, for instance, in Ref. 21.

The first Born proton-impact excitation cross sections of hydrogen and lithium from the ground states are shown in Fig. 5. The pseudopotential results for the $f_1 = 2s, 2p$ and $f_2 = 3s, 3p, 3d$ final states of hydrogen agree excellently with the analytical expression. For lithium, the pseudopotential cross sections agree in a broad velocity range with the DIM calculations, except for low proton-impact velocities. This disagreement arises from the form factor. For low impact velocities, the momentum transfer vector is large (16). As discussed earlier, in this region the bound momentum orbital $\tilde{\psi}(\mathbf{k} + \mathbf{p})$ is not described adequately by the pseudopotentials. An alternative expression for the form factor can be considered by implementing the peaking approximation

$$F_{fi}(\mathbf{p}) \sim \tilde{\psi}_i(\mathbf{p}) \tilde{\psi}_f^*(0) + \tilde{\psi}_f(\mathbf{p}) \tilde{\psi}_i^*(0). \quad (17)$$




Fig. 5 Proton-impact excitation cross section from the ground state for (A) hydrogen and (B) lithium.

Therefore, in order to have the correct form factor, it is necessary to obtain an accurate description of the initial bound state at large momentum values, which is not the case for the pseudostates (see Fig. 3B) and hence their failure when used in the cross section calculation.

s0040 3.2 Proton-impact ionization

p0105 The transition matrix (12) for the proton-impact ionization of X,


$$H^+ + X \rightarrow H^+ + X^+ + e^-, \quad (18)$$

can also be written in terms of the first-order Born approximation. In this case, the final state ψ_f in Eq. (14) is an outgoing continuum wavefunction $\psi_{\mathbf{k}_f}^-$, while $\epsilon_f = k_f^2/2$ is the energy of the ionized electron.

p0110 The single-differential proton-impact ionization cross sections $d\sigma/d\epsilon_f$ of hydrogen and lithium at a proton velocity of $v_p = 1$ a.u. are shown in Fig. 6. In the case of hydrogen, the pseudopotential and analytical results agree for all the electron energy range, except at very high values. On the other hand, for lithium, the cross sections computed with pseudopotentials only agree at low energies. Once again, assuming that $\psi_{\mathbf{k}_f}^-(\mathbf{k})$ can be approximated by a plane wave, the form factor is reduced to the Fourier transform of the initial bound state

$$F_{fi}(\mathbf{p}) \sim \tilde{\psi}_i(\mathbf{p} - \mathbf{k}_f). \quad (19)$$

Then, as k_f increases, so does p_{\min} , and the form factor is not well represented by the pseudopotentials. The significant discrepancies shown in Fig. 6 provide another demonstration of how a wrong description of the momentum space wavefunction may produce huge errors in collisional processes calculations.



f0035 **Fig. 6** Single differential proton-impact ionization cross section from the ground state of (A) hydrogen and (B) lithium at $v_p = 1$ a.u.

s0045 3.3 Proton-impact charge exchange

p0115 The proton-impact charge exchange of target X is defined as



The charge transfer cross section by the collision of a proton (electron capture) is computed with the first-order Brinkman–Kramers approximation.²² Accordingly, the matrix element is defined by

$$T_{fi}^{\text{BK}} = \tilde{\psi}_f^*(\mathbf{W}_f) \left[\varepsilon_f - \frac{W_f^2}{2} \right] \tilde{\psi}_i(\mathbf{W}_i),$$
 (21)


where \mathbf{W}_i and \mathbf{W}_f are the momentum transfer vectors

$$\mathbf{W}_i = W_{i0} \hat{\mathbf{v}} + \boldsymbol{\eta}, \quad W_{i0} = \frac{\nu}{2} - p_{\min}$$
 (22)

$$\mathbf{W}_f = W_{f0} \hat{\mathbf{v}} + \boldsymbol{\eta}, \quad W_{f0} = \frac{\nu}{2} + p_{\min},$$
 (23)

and they satisfy the condition $\mathbf{W}_i + \mathbf{W}_f = \mathbf{v}$, and p_{\min} is defined in Eq. (16).

p0120 The charge exchange cross sections of hydrogen and lithium in the ground state are illustrated in Fig. 7. The cross section of hydrogen is described with high accuracy by the pseudopotential approach for a wide range of proton velocities. However, this process constitutes a symmetrical resonance, i.e., $\varepsilon_f = \varepsilon_i$, and the agreement may be misleading. For the lithium case, the pseudopotentials fail utterly to describe the electron capture correctly at low and high velocities. For low and high ν_p values, the momentum transfer vector becomes large, and therefore, the cross sections calculated with pseudopotentials disagree completely for most of the energy values.



f0040 **Fig. 7** Proton-impact electron capture cross section for (A) hydrogen and (B) lithium in the ground state.

s0050 3.4 Photoionization

p0125 The single photoionization is defined as

$$\hbar\omega + X \rightarrow X^+ + e. \quad (24)$$

Considering a perturbative photon field, the initial bound ψ_i and final continuum $\psi_{\mathbf{k}_f}^-$ states of the target are not significantly distorted; therefore, the relevant matrix element of the photoionization process is given by


$$T_{\mathbf{k}}^{\text{Ph}} = \int \psi_{\mathbf{k}_f}^-(\mathbf{r}) [-i\hat{\epsilon}_\lambda \cdot \nabla_{\mathbf{r}}] \psi_i(\mathbf{r}), \quad (25)$$

where $\hat{\epsilon}_\lambda$ is the polarization versor and $\mathbf{k}_f = \sqrt{2(\omega + \epsilon_i)}$, as imposed by energy conservation.

p0130 The first-order photoionization cross sections of hydrogen and lithium are shown in Fig. 8. The pseudopotentials results for the hydrogen atom agree with the exact analytical expression results only for low photon energies, failing at larger values. These discrepancies can be understood considering the continuum wavefunction $\psi_{\mathbf{k}_f}^-(\mathbf{r})$ as a plane wave. Consequently, the matrix element $T_{\mathbf{k}}^{\text{Ph}}$ is reduced to

$$T_{\mathbf{k}}^{\text{Ph}} \sim -(\hat{\epsilon}_\lambda \cdot \mathbf{k}_f) \tilde{\psi}_i(\mathbf{k}_f), \quad (26)$$

and it is determined entirely by the behavior of the bound target pseudostate in the momentum representation. For hydrogen, the pseudo-orbital from PARSEC in the Fourier space coincides with the exact analytical solution for the entire range of k , which explains the excellent agreement in the cross section results. For lithium, the pseudopotential cross sections disagree with



f0045 **Fig. 8** Single photoionization cross section for (A) hydrogen and (B) lithium.

the DIM results for all energy values. The large oscillations in the cross sections are originated by the spurious oscillatory structure of the bound state for large k values (see inset of Fig. 3B).

s0055 3.5 DIM Photoionization of many-electron atoms


p0135 In order to assess the applicability of the depurated inversion method for atoms with a more complex structure, we compute the photoionization of many-electron targets with the DIM potentials⁴ and compare our results with experimental values. The first-order photoionization cross section of nitrogen and neon are shown in Fig. 9. Experimental data from^{23–26} is illustrated with hollow symbols. The DIM photoionization cross sections of these atoms agree excellently with the experimental values for low, medium and high photon energies. For neon, discrepancies start to be noticeable for low and intermediate energy. An accurate photoionization description of heavier atoms requires the inclusion of many-body effects that can be relevant, such as orbital relaxation due to the creation of a hole, collective response of inner shell electrons²⁷ and correlation effects.

s0060

4. Depurated inversion method for molecules

p0140

The depurated inversion method described above is extended here to determine effective potentials for molecules; methane is taken as an example. Furthermore, the molecular description of CH₄ given by DIM is tested by computing two collisional processes within the FBA.



f0050 **Fig. 9** Single photoionization cross section for (A) nitrogen and (B) neon.

s0065 4.1 Theory

p0145 Without loss of generality, we will present the DIM theoretical grounds for hydride compounds. The Hamiltonian of an N -electron XH_n molecule within the Born–Oppenheimer approximation is given by

$$\mathcal{H} = -\sum_{i=1}^N \frac{1}{2} \nabla_{\mathbf{r}_i}^2 - \sum_{i=1}^N \frac{Z_N}{r_i} + \sum_{i=1}^N V_H(r_i) + \sum_{i < j}^N \frac{1}{r_{ij}}, \quad (27)$$

$$V_H(r_i) = -\sum_{j=1}^n \frac{1}{|\mathbf{r}_i - \mathbf{R}_{H_j}|}, \quad (28)$$

where Z_N is the nuclear charge of the heavier atom, and \mathbf{R}_{H_j} are the coordinates of the hydrogens respect to the X atom. The corresponding Schrödinger equation $\mathcal{H}\Psi = E\Psi$ is solved and the orbitals are expressed as in Eq. (5) considering the single-center expansion (SCE). The orbitals and energies are found by solving the Hartree–Fock equations. The computation of these equations generally relies on the use of finite basis sets for the representation of the molecular orbitals (MOs). Usually, the MOs are expressed as a linear combination of atomic orbitals (LCAO),

$$\Psi_i = \sum_j c_{ji} \phi_j, \quad (29)$$

which can be constructed with Gaussian-type orbitals (GTO) basis sets.

p0150 The inverted molecular potential expression, analogous to Eq. (8), obtained from GTO basis sets present more difficulties than the atomic case. In addition to the asymptotic divergences and the poles, large unphysical oscillations arise.^{28–31} These prominent oscillations originate from undulations present in the MOs due to the finite number of the basis set. The second derivative, necessary to evaluate the inversion formula, amplifies these features.^{28,31} In some cases, the oscillations are huge, e.g., near an electronegative atom like Cl. The appearance of these oscillations in the inverted potentials forces us to incorporate further actions in the depuration scheme. To illustrate this procedure, we consider the $1s$ orbital of the carbon atom. We solved the Hartree–Fock equations using the 6-311G basis set with GAMESS code^{32,33} and obtained inverted potentials by implementing Eq. (8). The resulting Z_{1s}^{6-311G} charge is shown in Fig. 10A with a dot-dashed line. The charge oscillates significantly at low distances and diverges for higher r values. The same calculation was repeated using the universal Gaussian basis set (UGBS), which has a more significant amount of primitives. The corresponding inverted charge Z_{1s}^{UGBS} is exhibited in the figure




Fig. 10 (A) Effective charges for the 1s orbital of carbon. (B) Basis-set oscillation profiles.

with a dashed line. Although the charge still diverges around $r \approx 1$ a.u., the oscillations are now circumscribed near the nucleus. Finally, the differential Hartree–Fock equations for the carbon atom were solved using the finite-differences (FD) method. The 1s inverted charge obtained with this procedure, Z_{nl}^{FD} (solid line) shows no oscillations since no basis sets have been used to construct the orbital; however, the charge still diverges for $r > 1$ a.u., as it usually does for all HF calculations.

The oscillations pattern will vary for each basis set used in the calculations. We may define oscillation profiles as

$$p_{nl}^{BS} = Z_{nl}^{BS} - Z_{nl}^{FD}, \quad (30)$$

where Z_{nl}^{BS} is the inverted charge of the atom using a particular basis set “BS” and Z_{nl}^{FD} is the effective charge obtained from the inversion of the finite-difference wavefunctions. In the previous example, the basis set considered for calculating the 1s orbital of carbon were 6-311G and UGBS. The oscillation profiles for the 1s orbital, using Eq. (30) for these basis sets, are shown in Fig. 10B. Since the orbital profiles for each atomic basis set are distinctive, once they are determined for the atomic case, they can be removed in further molecular calculations. An example of this procedure is given in the following section.

4.2 Example: Methane

In order to illustrate the implementation of the DIM for molecules, we considered CH_4 , which is highly symmetric, and therefore, can be described with an angular averaged potential.³⁴ We computed the HF molecular orbitals and energies of CH_4 employing the UGBS basis sets of carbon and hydrogen, which considers angular momenta up to $L = 1$. Methane calculations with this basis set should include polarization functions




Fig. 11 (A) $1a_1$, (B) $2a_2$, and (C) $2t_1$ effective charges of CH_4 ; direct inversion (*dashed line*) and depurated inverted (*solid line*).

(at least d-functions) to increase the accuracy of the molecular energies.^{35,36} However, to isolate the effects of the basis set, we computed the atomic oscillation profiles and the molecular orbitals on the same footing. The charges obtained by direct inversion are given in Fig. 11 with dashed lines. Since the molecular orbitals are given by LCAO of carbon and hydrogen, the oscillations of the inverted charges are a consequence of the finite basis set of these atoms. To remove the most critical oscillations, first, we must determine the oscillation profiles produced by the atomic carbon basis set. We use Eq. (30) to determine the p_{1s}^{UGBS} , p_{2s}^{UGBS} , and p_{2p}^{UGBS} profiles of carbon. Then, we remove the oscillations by subtracting the carbon p_{nl}^{UGBS} profiles from the corresponding inverted charges Z_i^{UGBS} of CH_4 . The oscillations are removed for all orbitals except for the $2a_2$, which presents small oscillatory residues from the hydrogen basis set. Since the residual fluctuations are minimum and near the nucleus, we proceeded to implement the depuration scheme as described in Section 2.2. We define a new parametric DIM charge equation,

$$Z(r) = \sum_j Z_j e^{-\alpha_j r} + Z_{\text{H}} e^{-(\ln r - \ln \beta)^2 / (2\gamma)} + 1. \quad (31)$$

Table 1 Energies and fitting parameters for the DIM effective charges (Eq. (31)), for CH₄.

nl	E	Z	α	β	γ
1a ₁	−11.1949	1.925280	0.641982		
		0.953120	5.571510		
		2.121600	1.500440		
2a ₂	−0.9204	2.912200	3.149990		
		2.087800	0.771371		
		1.23640		2.329570	0.053420
2t ₁	−0.5042	0.901953	2.895140		
		1.112030	0.388649		
		2.986017	2.931210		
		1.30182		2.169850	0.012616

In contrast to the approximation proposed for atoms (9), a second term has been added to the formula to account for the presence of the hydrogens. This expression allows us to conveniently adjust both the location and width of the screened hydrogenic potential without affecting the correct charge value at the origin. The optimized parameters for the methane molecule are given in Table 1, and the corresponding DIM charges are shown in Fig. 11, with solid lines. The orbital energies obtained with these charges are also given in the table.

4.3 Collisional processes

The orientation of the molecular targets is important for determining the cross sections of collisional processes. However, it is generally not pre-established in the experiments. Thus, the spherically averaged description of the system assumed by the DIM potential makes sense. In the following, we examine two collisional processes in the first-order approximation: proton-impact ionization and single photoionization.

Proton-impact ionization

Results for the proton-impact ionization cross section for CH₄, calculated under the first Born approximation, are given in Fig. 12. The initial bound and the final continuum states of the molecule needed for the T-matrix computation (Eq. (12)) were calculated with the DIM potentials from




Fig. 12 Proton-impact ionization cross section for CH_4 . Solid line: first-order DIM theoretical calculations. Symbols: experiments from Refs. 37 and 38.




Fig. 13 Single photoionization cross section of CH_4 . Solid line: first-order DIM theoretical calculations. Symbols: experiments from Refs. 39–41.

Section 4.2. The ionization cross section for high and intermediate energies shows good agreement with the experimental results. The failure at low energies is ascribed to the validity of the first Born approximation and not to our DIM approach.

Photoionization

The photoionization cross section for CH_4 , calculated with the DIM potentials in a first-order approximation, is shown in Fig. 13 (solid lines). Good agreement with the experimental results (symbols) is found for high energy values and at the threshold. The curve between ~ 15 and ~ 300 eV shows the photoionization from the outer $n = 2$ shell, while the discontinuity

at 300 eV corresponds to the threshold of the $1a_1$ inner shell orbital. For low and intermediate photon energies, the agreement between our calculations and the experimental values from Refs. 39–41 is not that good. Phenomena such as molecular orbital relaxation, possible collective contributions, and correlation effects must be considered in further calculations. On the other hand, for the $1a_1$ inner shell photoionization, these effects are not significant, and we obtain a perfect agreement with the experimental results.

s0090

p0180

5. Concluding remarks

In this work, we explored the possibility of using pseudopotentials within the single electron model to calculate inelastic transitions. The first Born approximation was used to calculate proton-impact excitation, ionization, electron capture, and photoionization. Two simple atoms were studied, having a single electron in the outer shell. For hydrogen, we found excellent agreement for all the collisional processes, for low and intermediate energies. In the case of lithium, the only process that can be calculated with reasonable accuracy is the proton-impact excitation. We concluded that the range of validity is restrained to minimal momentum transfers. The depurated inversion method, on the other hand, accurately reproduces photoionization experimental results for many-electron atoms.

p0185

We extended the DIM for molecular systems. In this case, the inversion procedure produces huge oscillations due to the finite size of the basis sets involved in the Hartree–Fock orbital calculations. An additional step is included during the depuration scheme. In order to determine the oscillation profile for a particular basis set, we computed the inverted atomic charges in a finite-differences framework. By subtracting the charges, it is possible to isolate the oscillations corresponding to this particular basis set. We used the DIM method to determine the effective potentials for CH_4 . These potentials are implemented in first-order proton-impact ionization and photoionization cross sections calculations. For both processes, we found good agreement with the experimental results. The main discrepancies can be attributed to the fact that only first-order is considered in the perturbation theory.

Acknowledgments

st0105

p0190

The authors thank the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Universidad de Buenos Aires (UBA) and Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) for the grants that supported this work.

References

1. PARSEC Home Page. <https://parsec.ices.utexas.edu/styled-2/> (Accessed Jan 18, 2019).
2. Chelikowsky, J. R.; Troullier, N.; Saad, Y. Finite-Difference-Pseudopotential Method: Electronic Structure Calculations Without a Basis. *Phys. Rev. Lett.* **1994**, 72(8), 1240–1243. <https://doi.org/10.1103/PhysRevLett.72.1240>.
3. Mendez, A. M. P. *Método de Inversión Depurada para Potenciales Locales en Átomos y Moléculas*. 2015. Tesis de Licenciatura, Universidad Nacional de Salta.
4. Mendez, A. M. P.; Mitnik, D. M.; Miraglia, J. E. Depurated Inversion Method for Orbital-Specific Exchange Potentials. *Int. J. Quantum Chem.* **2016**, 116(24), 1882–1890. <https://doi.org/10.1002/qua.25295>.
5. Mendez, A. M. P.; Mitnik, D. M.; Miraglia, J. E. Local Effective Hartree-Fock Potentials Obtained by the Depurated Inversion Method. In *Nov. Electron. Struct. Theory Gen. Innov. Strongly Correl. Syst.*; Hoggan, P. E. Ed.; Advances in Quantum Chemistry; Vol. 76, Academic Press, 2018; pp 117–132. <https://doi.org/10.1016/bs.aiq.2017.07.004>.
6. Bates, D. R. Theoretical Treatment of Collisions Between Atomic Systems. *Pure Appl. Phys. Elsevier* **1962**, 13, 549–621. In *At. Mol. Process.*; Bates, D. R. Ed.
7. McDowell, M. R. C.; Peach, G. Ionization of Lithium by Fast Protons and Electrons. *Phys. Rev.* **1961**, 121, 1383–1387.
8. Pindzola, M. S.; Robicheaux, F.; Loch, S. D.; Berengut, J. C.; Topcu, T.; Colgan, J.; Foster, M.; Griffin, D. C.; Ballance, C. P.; Schultz, D. R.; Minami, T.; Badnell, N. R.; Witthoef, M. C.; Plante, D. R.; Mitnik, D. M.; Ludlow, J. A.; Kleiman, U. The Time-Dependent Close-Coupling Method for Atomic and Molecular Collision Processes. *J. Phys. B* **2007**, 40, R39–R60.
9. Burke, P. G. *R-Matrix Theory of Atomic Collisions*; Springer-Verlag: Berlin, Heidelberg, 2011.
10. Bray, I.; Abdurakhmanov, I. B.; Bailey, J. J.; Bray, A. W.; Fursa, D. V.; Kadyrov, A. S.; Rawlins, C. M.; Savage, J. S.; Stelbovics, A. T.; Zammit, M. C. Convergent Close-Coupling Approach to Light and Heavy Projectile Scattering on Atomic and Molecular Hydrogen. *J. Phys. B* **2017**, 50, 202001.
11. Pindzola, M. S.; Colgan, J.; Robicheaux, F.; Lee, T. G.; Ciappina, M. F.; Foster, M.; Ludlow, J. A.; Abdel-Naby, S. A. Time-Dependent Close-Coupling Calculations for Ion-Impact Ionization of Atoms and Molecules. *Adv. At. Mol. Opt. Phys.* **2016**, 65, 291–319.
12. Kirchner, T.; Gulyás, L.; Lüdde, H. J.; Engel, E.; Dreizler, R. M. Influence of Electronic Exchange on Single and Multiple Processes in Collisions Between Bare Ions and Noble-Gas Atoms. *Phys. Rev. A* **1998**, 58, 2063–2076.
13. Fiori, M. R.; Jalbert, G.; Bielschowsky, C. E.; Cravero, W. Ionization of Lithium by Impact of Fast Ions. *Phys. Rev. A* **2001**, 64, 012705.
14. Szabo, A.; Ostlund, N. S. *Modern Quantum Chemistry: Introduction to Advanced Electronic Structure Theory*; Dover Publications, Inc.: Mineola, NY, 1996.
15. Helgaker, T.; JØrgensen, P.; Olsen, J. *Molecular Electronic-Structure Theory*. John Wiley & Sons, Ltd: Chichester, UK, 2000.
16. Schaefer, H. F. *Quantum Chemistry: The Development of Ab Initio Methods in Molecular Electronic Structure Theory*; Dover Publications, Inc: Mineola, NY, 2004.
17. ABINIT Home Page. <https://www.abinit.org/psp-tables> (Accessed Jan 18, 2019).
18. Hamann, D. R.; Schlüter, M.; Chiang, C. Norm-Conserving Pseudopotentials. *Phys. Rev. Lett.* **1979**, 43(20), 1494–1497. <https://doi.org/10.1103/PhysRevLett.43.1494>.
19. Froese Fischer, C.; Brage, T.; Jönsson, P. *Computational Atomic Structure: An MCHF Approach*. Institute of Physics Publishing: Bristol, UK, 1997; p 279.
20. Johnson, W. R. *Atomic Structure Theory : Lectures on Atomic Physics*. Springer: Berlin, Heidelberg, 2007; p 312.

21. McDowell, M. R. C.; Coleman, J. P. *Introduction to the Theory of Ion-Atom Collisions*; North-Holland Publishing Company: Amsterdam, 1970.
22. C, B. H.; Kramers, H. A. *Physica.-Zur Theorie der Einfangung von Elektronen durch u. Teilchen. Proc. K. Akad. van Wet.* **1930**, *33*, 973–984. <http://www.dwc.knaw.nl/DL/publications/PU00015982.pdf>.
23. Henke, B. L.; Gullikson, E. M.; Davis, J. C. X-Ray Interactions: Photoabsorption, Scattering, Transmission, and Reflection at $E = 50\text{--}30,000$ eV, $Z = 1\text{--}92$. *At. Data Nucl. Data Tables* **1993**, *54*(2), 181–342. <https://doi.org/10.1006/ADND.1993.1013>.
24. Samson, J. A. R.; Angel, G. C. *Single- and double-photoionization cross sections of atomic nitrogen from threshold to 31 Å*. *Phys. Rev. A* **1990**, *42*(3), 1307–1312. <https://doi.org/10.1103/PhysRevA.42.1307>. <https://link.aps.org/doi/10.1103/PhysRevA.42.1307>.
25. Samson, J. A. R.; Stolte, W. C. Precision Measurements of the Total Photoionization Cross-Sections of He, Ne, Ar, Kr, and Xe. *J. Electron Spectros. Relat. Phenomena* **2002**, *123*(2–3), 265–276. [https://doi.org/10.1016/S0368-2048\(02\)00026-9](https://doi.org/10.1016/S0368-2048(02)00026-9).
26. Stolte, W. C.; Jonauskas, V.; Lindle, D. W.; Sant’Anna, M. M.; Savin, D. W. *Inner-Shell Photoionization Studies of Neutral Atomic Nitrogen. Astrophys. J.* **2016**, *818*(2), 149. <https://doi.org/10.3847/0004-637X/818/2/149>. <http://stacks.iop.org/0004-637X/818/i=2/a=149?key=crossref.2b86a13e417a21fe226f2d7fdc09b8c8>.
27. Ederer, D. L. Photoionization of the 4d Electrons in Xenon. *Phys. Rev. Lett.* **1964**, *13*(25), 760–762. <https://doi.org/10.1103/PhysRevLett.13.760>.
28. Schipper, P. R. T.; Gritsenko, O. V.; Baerends, E. J. Kohn-Sham potentials corresponding to Slater and Gaussian basis set densities. *Theor. Chem. Accounts Theory, Comput. Model. (Theoretica Chim. Acta)* **1997**, *98*(1), 16–24. <https://doi.org/10.1007/s002140050273>.
29. Mura, M. E.; Knowles, P. J.; Reynolds, C. A. Accurate Numerical Determination of Kohn-Sham Potentials From Electronic Densities: I. Two-Electron Systems. *J. Chem. Phys.* **1997**, *106*(23), 9659–9667. <https://doi.org/10.1063/1.473838>.
30. Jacob, C. R. Unambiguous Optimization of Effective Potentials in Finite Basis Sets. *J. Chem. Phys.* **2011**, *135*(24), 244102. <https://doi.org/10.1063/1.3670414>.
31. Gaiduk, A. P.; Ryabinkin, I. G.; Staroverov, V. N. *Removal of Basis-Set Artifacts in KohnSham Potentials Recovered From Electron Densities. J. Chem. Theory Comput.* **2013**, *9*(9), 3959–3964. <https://doi.org/10.1021/ct4004146>. <http://www.ncbi.nlm.nih.gov/pubmed/26592391>.
32. Schmidt, M. W.; Baldrige, K. K.; Boatz, J. A.; Elbert, S. T.; Gordon, M. S.; Jensen, J. H.; Koseki, S.; Matsunaga, N.; Nguyen, K. A.; Su, S.; Windus, T. L.; Dupuis, M.; Montgomery, J. A. General Atomic and Molecular Electronic Structure System. *J. Comput. Chem.* **1993**, *14*(11), 1347–1363. <https://doi.org/10.1002/jcc.540141112>.
33. Gordon, M. S.; Schmidt, M. W. Advances in Electronic Structure Theory: GAMESS a Decade Later. In *Theory Appl. Comput. Chem.*; Dykstra, C. E., Frenking, G., Kim, K. S., Scuseria, G. E., Eds.; Elsevier: Amsterdam, 2005; pp 1167–1189. chap. 41. <https://doi.org/10.1016/B978-044451719-7/50084-6>.
34. Granados-Castro, C. M. Application of Generalized Sturmian Basis Functions to Molecular Systems. Ph.D. thesis, Université de Lorraine, Metz, France and Universidad Nacional del Sur, Bahía Blanca, Argentina 2016.
35. Rothenberg, S.; Schaefer, H. F. Methane as a Numerical Experiment for Polarization Basis Function Selection. *J. Chem. Phys.* **1971**, *54*, 2764–2766.
36. Hariharan, P. C.; Pople, J. A. The Effect of D-Functions on Molecular Orbital Energies for Hydrocarbons. *Chem. Phys. Lett.* **1972**, *16*, 217–219.

37. Rudd, M. E.; DuBois, R. D.; Toburen, L. H.; Ratcliffe, C. A.; Goffe, T. V. Cross Sections for Ionization of Gases by 5–4000-keV Protons and for Electron Capture by 5–150-keV Protons. *Phys. Rev. A* **1983**, 28(6), 3244–3257. <https://doi.org/10.1103/PhysRevA.28.3244>.
38. Rudd, M. E.; Kim, Y. K.; Madison, D. H.; Gallagher, J. W. Electron Production in Proton Collisions: Total Cross Sections. *Rev. Mod. Phys.* **1985**, 57(4), 965–994. <https://doi.org/10.1103/RevModPhys.57.965>.
39. Lukirskii, A. P.; Brytov, I. A.; Zimkina, T. M. *Optika I Spekt.* Vol. 17, 1964234. www.scopus.com.
40. Henke, B. L.; Lee, P.; Tanaka, T. J.; Shimabukuro, R. L.; Fujikawa, B. K. Low-energy X-ray Interaction Coefficients: Photoabsorption, Scattering, and Reflection: $E = 1002000$ eV $Z = 194$. *At. Data Nud. Data Tables* **1982**, 27(1), 1–144. [https://doi.org/10.1016/0092-640X\(82\)90002-X](https://doi.org/10.1016/0092-640X(82)90002-X).
41. Samson, J. A. R.; Haddad, G. N.; Masuoka, T.; Pareek, P. N.; Kilcoyne, D. A. L. Ionization Yields, Total Absorption, and Dissociative Photoionization Cross Sections of CH₄ from 110 to 950 Å. *J. Chem. Phys.* **1989**, 90(12), 6925–6932. <https://doi.org/10.1063/1.456267>.

CORRECTED PROOF

Non-Print Items

Keywords: Atomic effective potential, Molecular effective potential, Depurated inversion method, Inverted Hartree–Fock wave functions, Pseudopotentials, Inelastic Collisions

CORRECTED PROOF