

Index

Note: Page numbers followed by “*f*” indicate figures, “*t*” indicate tables, and “*b*” indicate boxes.

A

Ab initio Frenkel–Davydov exciton model (AIFDEM)
2-methylene-2H-indene, 276–283
protocol, 270–271
tetracene dimer, 271–276
Addition theorems (AdTs), 100–101
Alkali hydrides, 222. *See also* NaH molecule
Analysis, multiresolution (MRA), 7
adaptiveness, 12–13
applications, 40–43, 41*f*
arithmetic operations, 17–24
differentiation, 18–19
Haar scaling function, 8, 9–10*f*, 14–15
inner product, 17–18
integral operator, 20–24
low-rank tensor approximations, 25–26
nuclear singularities, regularization of, 29–32
singular value decomposition (SVD), 25–26, 47
tree structure, 12–13
wavelet formalism, 7–11, 13–17, 24–25
Analytic continuation method, 326
Analytic energy gradient expressions, 167
Angular momentum theory, 68–69
Angular energy, 147–148, 147*t*
Askey scheme, 67–70
Asymptotic theory of functional analysis, 67–68, 72
Atom-centered basis functions, 5–6
Atomic effective potential, 186–191

B

Basis set superposition error (BSSE), 5–6
Benchmarks (for metal-catalyst systems), 313–314
Beryllium hydride, cusplless and
cusp-corrected HF wave functions
of, 124–126, 125–126*f*
Bessel functions (BFs), 98–100

Bohr atom, 64–65
Bond dissociation, 314, 321
Born–Oppenheimer approximation, 192
Boron anion, 140–147*t*, 146–151, 149–151*t*
Bound-state, 22–23, 34–35
Brinkman–Kramers approximation, 189

C

Carbon K-edge X-ray absorption, 244, 246*f*
CASINO code, 312, 317
Casula T-moves, 319–320
Central ionsphere–central ionsphere
integral, 104–105
Central ionsphere–noncentral ionsphere
integral, 105
Charge density, one-body (1-CD), 291–294
Charge transfer (CT) states, 264–265
Chlorophyll-a, MRA computation of, 40–41, 41*f*
11-Cis-retinal, MRA computation of, 40–41, 41*f*
Clebsch–Gordan coefficient, 106–109
Cluster ansatz, 4, 30, 38–40, 83
Cluster wavefunction normalization.
See Overlap integrals of interstitial
region
Collisions (inelastic, of electrons with atoms
and molecules), 180, 186–197
Complete active space self-consistent field
(CASSCF) wave functions, 266–267
Complete basis set (CBS) limit, 248–249
Configuration interaction (CI) method, for
boron anion, 136–148, 140–146*t*
SOS-CI (scaled opposite spin), 269
Configuration interaction singles (CIS),
40–41
Confinement (H-atom), 324–325, 328–330,
332–334
Continuous and discrete expansions, 70–73
Core excitations, 242, 244, 246–247, 257
Core-level spectroscopy, 242

Core–valence separation (CVS), 243
Coulomb Sturmian functions (CSF), 84–85
Coulomb three-body Schrödinger equation, 80–84
Coupled cluster linear response (CC-LR), 242–243
Coupled cluster (CC) method, 36–38, 222–223
Coupled cluster singles and doubles (CCSD), 170
Coupled cluster singles, doubles and triples (CCSDT), 243
ionization energies, 253–257, 254*t*, 256*t*
Coupled perturbed Hartree–Fock (CPHF) equations, 34–35
Cusp-corrected orbitals, 116–117, 119
Cuspless Gaussian basis functions, 114–115, 121–122

D

Density-based descriptors, 291–293, 298–299
Density functional theory (DFT), 28–29, 202
Density matrix
attachment A, 291–292
detachment D, 291–292
Depurated inversion method (DIM), 180, 185
many-electron atoms, 191
molecules, 191–197
Descriptors, density-based, 291–293, 298–299
Detachment and attachment one-body reduced density matrices, 290–294, 295*t*
Diffusion Monte Carlo (DMC) method, 317, 320
Disequilibrium function, 326, 331–332, 331*f*
Double electron affinity (DEA), 222

E

Eckart–Sayvetz conditions, 171–172
Effective Hamiltonian operator, 223–225
Electron affinity (EA), 222, 225
boron atom, 149, 149–150*t*
definition, 136, 148–149

Electron–electron potential energy, 156–157
Electronic coupling
2-methylene-2H-indene, 281–283, 281*t*
tetracene pair, 275, 275*t*
Electron–nucleus cusp correction algorithm, 119, 120*b*
Electron pairing approach, 161–163
Encapsulated atoms
applications, 323–324
bound state properties, 324
Equation-of-motion coupled cluster (EOM-CC) method, 222–223, 225, 231, 242–243
Equilibrium geometries, 170
Euler–Lagrange equation, 202–203
Exchange–correlation potential, 207–208
CASSCF wavefunctions, 212–213, 213*f*
HF wavefunctions, 211–212, 211–212*f*
unified, 204, 209, 215
Excited-state electronic structure, 290
boron anion, 149–151
helium, 83–84, 84*f*
2-methylene-2H-indene, 277–278
NaH, 226, 226*t*
phenylpyrrole molecules, 301–304, 302*f*, 303*t*
tetracene pair, 272–274, 275*t*
Exponential Shannon information entropy, 332, 332*f*

F

Favard’s theorem, 56
Fermi potential, 203, 205–207, 210
ab initio wavefunctions, 210–214, 211*t*
algorithm, 209–210
CASSCF wavefunctions, 212–213, 213*f*
HF wavefunctions, 211–212, 211–212*f*
Fermi’s golden rule, 266–267
First Born approximation (FBA), 180–181
First principle quantum chemical methods, 222–223, 233–234
Fisher information measure, 327, 331–332, 331*f*
Fisher–Shannon complexity (FS) index, 327
Fock matrix, 117–120
Fock space coupled cluster (FS-CC) method, 222–225

Fock space (FS) theory, 222–223
Full configuration interaction (FCI),
164–165, 164*f*

G

Galerkin method, 86–88
Gauge-invariant atomic orbitals (GIAO),
33–34
Gaunt coefficients, 100–101
Gaussian basis functions, 5–6
Generalized Sturmian functions (GSF),
80–82
Geometry optimization, 166–174
Geminals (strongly orthogonal), 158

H

Hahn polynomials, 66–67
Harmonic vibrational frequencies, 170–174
Hartree–Fock (HF) approximation, 157
Hartree–Fock equations, 28–29, 192–193
Hartree–Fock (HF) wavefunctions, 210
Helium atom
 cusplless and cusp-corrected HF wave
 functions, 122–124, 123–124*f*, 124*t*
 scattering states, 79–80, 83–84, 84*f*
Hellmann–Feynman theorem, 32–33
Hermite polynomials, 62–64
Hessian (expression, matrix), 168–172
Hohenberg–Kohn theorem, 156, 202
Hubbard model, 163–165
Hydrogen atom
 confinement, 324–325
 complexity measures, 326
 ionization probability, 325–326, 330,
 330*f*
 LMC shape complexity measure,
 332–334, 333*f*
 orbital energy *vs.* confinement radius,
 327–330, 327–329*f*
 reduced radial functions, 328–329, 328*f*
 Shannon entropy, 332, 332*f*
 cusplless and cusp-corrected HF wave
 functions, 121–122, 121*f*, 122*t*
 physisorption, 313, 319–320
Hydrogen-like wavefunctions, 62, 64–65
Hylleraas–configuration interaction (Hy-CI)
 calculations, 137–139, 152
Hypergeometric functions, 56–59, 65, 68

Hypergeometric polynomials, 56–57,
67–68, 70
Hyperquantization algorithm, 73
Hyperspherical harmonics, 59–62, 71

I

Inelastic collisions, 180–181, 186, 197
Information theoretical tools, 331
Intermediate Hamiltonian (IH) formulation,
225
Interstitial, 98–100, 102–111
Inverted Hartree–Fock equations, 192–193
Ionization probability, 324–325, 330, 330*f*
Ionsphere, 98–99

J

Jacobi ladder, 67–70
Jacobi polynomials, 65
Jastrow factor, 114–116, 316–317

K

Kepler–Coulomb problem, 64–65, 70–71
Kinetic energy integrals, 128
Kohn–Sham density-functional scheme,
202–204, 207–209
Kohn–Sham orbitals, 315
Kravchuk polynomials, 68, 72
Kronecker delta, 116–117

L

Laguerre polynomials, 64–65
Legendre polynomials, 8–10, 106–109
Levy–Perdew–Sahni formalism, 203–204
Light-induced electronic structure
 reorganization, 290
Linear combination of atomic orbitals
 (LCAO), 5–6, 33–34, 38–43, 45, 48
Löwdin-like population analysis, 293–294

M

Many-electron basis functions (MEBFs),
266–269, 274–275, 278–281
Mass weighted Cartesian coordinates
 (MWC), 170–171
Metal surface, 312–313, 319
Methane, DIM for, 193–195
Modified Sternheimer equation, 34–35

Moeller-Plesset perturbative (MP2), 36–38,
170, 172, 173*f*
Molecular effective potential, 191–197
Molecular nano-containers, 323–324
Muffin-tin approximation, 98
Mulliken population analysis, 293
Multireference CI (MRCI) approach,
230–231
Multiresolution analysis (MRA). *See*
Analysis, multiresolution (MRA)

N

NaH⁺ molecular ion
potential energy curves, 231–233, 232*f*
spectroscopic constants, 228*t*, 233
NaH molecule
energies of electronic states, 226, 226*t*
potential energy curves, 229, 229–230*f*
spectroscopic constants, 226, 227*t*
Natural orbital functional (NOF), 157–160
Natural orbitals (NOs), 157
Nitrogen K-edge X-ray absorption, 244,
248*f*, 251, 252*f*
Noncentral integrals c.f. central case,
105–110
Nonorthogonal configuration interaction
(NOCI), 266–267
2-methylene-2H-indene, 276–283
tetracene dimer, 271–276
N-representability, 157–161, 163–164
Nuclear attraction integrals, 129
Numerical integration on Cartesian grids
(NI grid), 293, 295–296, 298

O

Occupation numbers (ONs), 157, 159–160
One-particle reduced density matrix
(1RDM), 156–157, 205–207
Orbital population analysis, 293–295
Orthogonal polynomials, 56–63, 65–72
Overlap integrals, 127–128
Overlap integrals of interstitial region
central ionsphere–central ionsphere
integral, 104–105
central ionsphere–noncentral ionsphere
integral, 105

noncentral ionsphere–different
noncentral ionsphere integral,
109–110
noncentral ionsphere–same noncentral
ionsphere integral, 105–109
Watson–central ionsphere integral,
102–103
Watson–noncentral ionsphere integral,
103–104
Watson–Watson integral, 102
Oxygen K-edge X-ray absorption, 244,
245*f*, 247*f*, 249, 250*f*

P

Pauli potential, 203–204, 207–208
CASSCF wavefunctions, 212–213, 213*f*
HF wavefunctions, 211–212, 211–212*f*
Penetrable repulsive spherically symmetric
model for confinement, 324
Photoelectron spectroscopy, 242
Photoionization, 190–191, 197
Piris natural orbital functional (PNOF),
160–163
independent pair model PNOF5, 158,
161–163, 165–168, 170
inter-pair electron correlation model
PNOF7, 163, 165–168, 170,
172–174
Plane waves, 6
Polarizable continuum model, 42
Population analysis, 293–295
Potential energy curves (PECs), 222–223
Proton-impact charge exchange, 189
Proton-impact excitation, 186–188
Proton-impact ionization, 188, 195–196
Pseudopotential approximation (PPA),
180–184
Pseudo-potentials, 317–319

Q

Quadrature mirror coefficients (QMF),
15–16
Quantum Monte Carlo (QMC) method,
312–313
fixed-node error, 312–313, 319
heterogeneous systems, 312, 315
metal surface, 312–314, 319

R

- Racah coefficients, 66–67
- Racah polynomials, 66–67
- Real-space grid methods, 6
- Reduced density matrix (RDM)
 - one-particle, 156–157
 - two-particle, 156–157
- Restricted active space with two spin flips (RAS-2SF)
 - 2-methylene-2H-indene, 276–283
 - protocol, 271
 - tetracene dimer, 271–276
- Restricted Hartree–Fock (RHF) reference function, 222
- Rydberg excited states, 151, 251–253

S

- Scale up methods, 114
- Scattered wavefunction method, 98
- Schrödinger equation, 26–28, 99–100, 137–138, 312
- Self-consistent dressing (SCD), 118–119
- Shannon information entropy, 326–327
- Shape LMC complexity index, 326, 332–334, 333*f*
- Shockley–Queisser limit, 264
- Singlet fission (SF), 264–267, 269, 276–278
- Slater basis functions, 115–116, 121–122
- Slater–Jastrow wave function, 316
- Slater type orbitals (STOs), 137–139, 149, 151
- Spatial confinement, 323–324
- Specific reaction parameter density functional theory (SRP-DFT), 313
- Spectral method, 80–82
- Spherical harmonic (SH) function, 99–101
- Spherical harmonics, 59–62
- Spin-free exact 2C theory, 243
- Standard basis, 249, 254*t*
- Sturmian functions (SF)

- one-dimensional, 81, 84–87, 85–86*f*, 89–93, 91*t*, 91*f*
- two-dimensional, 87–93, 91*f*, 91*t*
- Sturmian orbitals, 64–65, 70
- Sturm–Liouville theory, 56
- Sudden approximation, 324
- Supercomputer (architecture), 114

T

- Tamm–Dancoff approximation (TDA), 42–43
- Time-dependent density functional theory (TD-DFT), 242
- Time-dependent Hartree–Fock (TDHF) calculations, 297
- Transition-state (TS) geometry, 312–313
- Tridiagonal and symmetric matrices, 56
- Two-particle reduced density matrix (2RDM), 156–157, 205–207, 209

U

- Unrestricted Hartree–Fock (UHF) reference function, 222

V

- Variational Monte Carlo (VMC), 316–317, 320
- Vibrational frequencies, 170–174

W

- Water molecule
 - molecular properties, 42–43, 42*t*
 - solvatochromatic shifts, 42, 42*t*
 - X-ray absorption, 244, 245*f*
- Watson–central ionsphere integral, 102–103
- Watson–noncentral ionsphere integral, 103–104
- Watson sphere, 98–99
- Watson–Watson integral, 102
- Wigner matrix elements, 68–70