10_2Q4

David Beauchemin
30 novembre 2017

10.2Q.4

a)

i)

```
\begin{split} F_{X_1,X_2}(X_1,X_2) &= P\left(X_1 \leq x_1, X_2 \leq x_2\right) \\ &= P\left(F_{X_1}^{-1}(U_1) \leq x_1, F_{X_2}^{-1}(U_2) \leq x_2\right) \\ &= P\left(U_1 \leq F_{X_1}(x_1), U_2 \leq F_{X_2}(x_2)\right) \\ &= C_{\alpha}^{EFGM}\left(F_{X_1}(x_1), F_{X_2}(x_2)\right) \end{split}
```

ii)

```
alpha <- 1
densityCopule <- function(u1, u2) {
u1 * u2 + alpha * u1 * u2 * (1 - u1) * (1 - u2)
}
FXX <- function(x1, x2) {
densityCopule(pexp(x1, 1 / 100), plnorm(x2, log(100) - 0.32, 0.8))
}
FXX(100, 100)</pre>
```

[1] 0.4668243 FXX(200, 100)

[1] 0.5931482

FXX(100, 300)

[1] 0.6165623

b)

$$\begin{split} F_{X_1,X_2}(X_1,X_2) &= P\left(X_1 \leq x_1, X_2 \leq x_2\right) \\ &= P\left(F_{X_1}^{-1}(U_1) \leq x_1, F_{X_2}^{-1}(1-U_2) \leq x_2\right) \\ &= P\left(U_1 \leq F_{X_1}(x_1), 1-U_2 \leq F_{X_2}(x_2)\right) \\ &= P\left(U_1 \leq F_{X_1}(x_1), U_2 > 1-F_{X_2}(x_2)\right) \\ &= P\left(U_1 \leq F_{X_1}(x_1), U_2 > \overline{F}_{X_2}(x_2)\right) \\ &= 1 - \overline{F}_{X_2}(x_2) - C_{\alpha}^{EFGM}\left(F_{X_1}(x_1), \overline{F}_{X_2}(x_2)\right) \end{split}$$

ii)

```
FXX <- function(x1, x2) {</pre>
     1 - pexp(x1, 1 / 100, lower.tail = FALSE) - densityCopule(pexp(x1, 1 / 100),
      1 - plnorm(x2, log(100) - 0.32, 0.8))
FXX(100, 100)
## [1] 0.3617868
FXX(200, 100)
## [1] 0.5402919
FXX(100, 300)
## [1] 0.5995209
c)
                        F_{X_1,X_2}(X_1,X_2) = P(X_1 \le x_1, X_2 \le x_2)
                                         = P(F_{X_1}^{-1}(1 - U_1) \le x_1, F_{X_2}^{-1}(U_2) \le x_2)
                                         = P(1 - U_1 \le F_{X_1}(x_1), U_2 \le F_{X_2}(x_2))
                                         = P(U_1 > 1 - F_{X_1}(x_1), U_2 \le F_{X_2}(x_2))
                                         = P(U_1 > \overline{F}_{X_1}(x_1), U_2 \le F_{X_2}(x_2))
                                         = 1 - \overline{F}_{X_1}(x_1) - C_{\alpha}^{EFGM}(\overline{F}_{X_1}(x_1), F_{X_2}(x_2))
ii)
FXX <- function(x1, x2) {</pre>
      1 - (1 - plnorm(x2, log(100) - 0.32, 0.8)) - densityCopule(pexp(x1, 1 /
      100, lower.tail = FALSE),
      plnorm(x2, log(100) - 0.32, 0.8))
}
FXX(100, 100)
## [1] 0.3617868
FXX(200, 100)
## [1] 0.5402919
FXX(100, 300)
## [1] 0.5995209
```

d)

```
\begin{split} F_{X_1,X_2}(X_1,X_2) &= P\big(X_1 \leq x_1,X_2 \leq x_2\big) \\ &= P\big(F_{X_1}^{-1}(1-U_1) \leq x_1,F_{X_2}^{-1}(U_2) \leq x_2\big) \\ &= P\big(1-U_1 \leq F_{X_1}(x_1),1-U_2 \leq F_{X_2}(x_2)\big) \\ &= P\big(U_1 > 1-F_{X_1}(x_1),U_2 > 1-F_{X_2}(x_2)\big) \\ &= P\big(U_1 > \overline{F}_{X_1}(x_1),U_2 > \overline{F}_{X_2}(x_2)\big) \\ &= 1 - \overline{F}_{X_1}(x_1)\overline{F}_{X_2}(x_2) + C_{\alpha}^{EFGM}\big(\overline{F}_{X_1}(x_1),\overline{F}_{X_2}(x_2)\big) \end{split}
```

ii)

```
FXX <- function(x1, x2) {
    1 - (1 - plnorm(x2, log(100) - 0.32, 0.8)) - pexp(x1, 1 / 100, lower.tail = FALSE) +
    densityCopule(pexp(x1, 1 /
    100, lower.tail = FALSE),
    plnorm(x2, log(100) - 0.32, 0.8))
}
FXX(100, 100)
## [1] 0.5811772
FXX(200, 100)
## [1] 0.6352163
FXX(100, 300)</pre>
```

[1] 0.956415