


$$D_\mu \text{ vs } \langle \cdot, \cdot \rangle_\mu^\pm$$

Equivalence between two notions of viscosity solutions in the Wasserstein space

An abstract, colorful, multi-lobed shape on the left side of the slide. It features a gradient from purple at the bottom to yellow at the top, with blue and green accents on the sides.

Averil Prost (LMI, INSA Rouen Normandie)
Hasnaa Zidani (LMI, INSA Rouen Normandie)

March 22, 2024
ANR COSS Meeting

INSA



anr®

Aim of the talk

We consider a first-order Hamilton-Jacobi equation of the form

$$H(\mu, D_\mu V(\mu)) = 0 \quad \mu \in \Omega, \quad V(\mu) = \mathfrak{J}(\mu) \quad \mu \in \partial\Omega. \quad (1)$$

Aim of the talk

We consider a first-order Hamilton-Jacobi equation of the form

$$H(\mu, D_\mu V(\mu)) = 0 \quad \mu \in \Omega, \quad V(\mu) = \mathfrak{J}(\mu) \quad \mu \in \partial\Omega. \quad (1)$$

Here

- μ is a measure, Ω an open set of the Wasserstein space $\mathcal{P}_2(\mathbb{R}^d)$.

Aim of the talk

We consider a first-order Hamilton-Jacobi equation of the form

$$H(\mu, D_\mu V(\mu)) = 0 \quad \mu \in \Omega, \quad V(\mu) = \mathfrak{J}(\mu) \quad \mu \in \partial\Omega. \quad (1)$$

Here

- μ is a measure, Ω an open set of the Wasserstein space $\mathcal{P}_2(\mathbb{R}^d)$.
- $D_\mu V(\mu)$ is the application of the directional derivatives.

Aim of the talk

We consider a first-order Hamilton-Jacobi equation of the form

$$H(\mu, D_\mu V(\mu)) = 0 \quad \mu \in \Omega, \quad V(\mu) = \mathfrak{J}(\mu) \quad \mu \in \partial\Omega. \quad (1)$$

Here

- μ is a measure, Ω an open set of the Wasserstein space $\mathcal{P}_2(\mathbb{R}^d)$.
- $D_\mu V(\mu)$ is the application of the directional derivatives.

Our aim Compare two notions of viscosity solutions for (1).

Table of Contents

First definitions

Geometric tangent space

Generalized sub and superdifferentials

Definitions of viscosity solutions


The equivalence result

Viscosity solutions

In \mathbb{R}^d , viscosity solutions of $H(x, \nabla_x u) = 0$ are equivalently defined using

- smooth test functions:

u is a subsolution if it is u.s.c, satisfies $u \leq \mathfrak{J}$, and if whenever $\varphi \in \mathcal{C}^1$ is such that $u - \varphi$ reaches a maximum at x ,



there holds $H(x, \nabla \varphi(x)) \leq 0$.


- sub and superdifferentials:

Viscosity solutions

In \mathbb{R}^d , viscosity solutions of $H(x, \nabla_x u) = 0$ are equivalently defined using

- smooth test functions:

u is a supersolution if it is l.s.c, satisfies $u \geq \mathfrak{J}$, and if whenever $\varphi \in \mathcal{C}^1$ is such that $u - \varphi$ reaches a minimum at x ,



there holds $H(x, \nabla \varphi(x)) \geq 0$.


- sub and superdifferentials:

Viscosity solutions

In \mathbb{R}^d , viscosity solutions of $H(x, \nabla_x u) = 0$ are equivalently defined using

- smooth test functions:


u is a supersolution if it is l.s.c, satisfies $u \geq \mathfrak{J}$, and if whenever $\varphi \in \mathcal{C}^1$ is such that $u - \varphi$ reaches a minimum at x ,



there holds $H(x, \nabla \varphi(x)) \geq 0$.

- sub and superdifferentials:

u is a subsolution if it is u.s.c, satisfies $u \leq \mathfrak{J}$, and if whenever a vector v belongs to the superdifferential of u at x ,




there holds $H(x, v) \leq 0$.

Viscosity solutions

In \mathbb{R}^d , viscosity solutions of $H(x, \nabla_x u) = 0$ are equivalently defined using

- smooth test functions:


u is a supersolution if it is l.s.c, satisfies $u \geq \mathfrak{J}$, and if whenever $\varphi \in \mathcal{C}^1$ is such that $u - \varphi$ reaches a minimum at x ,



there holds $H(x, \nabla \varphi(x)) \geq 0$.

- sub and superdifferentials:

u is a supersolution if it is l.s.c, satisfies $u \geq \mathfrak{J}$, and if whenever a vector v belongs to the subdifferential of u at x ,




there holds $H(x, v) \geq 0$.

Viscosity solutions

In \mathbb{R}^d , viscosity solutions of $H(x, \nabla_x u) = 0$ are equivalently defined using

- smooth test functions:


u is a supersolution if it is l.s.c, satisfies $u \geq \mathfrak{J}$, and if whenever $\varphi \in \mathcal{C}^1$ is such that $u - \varphi$ reaches a minimum at x ,



there holds $H(x, \nabla \varphi(x)) \geq 0$.

- sub and superdifferentials:

u is a supersolution if it is l.s.c, satisfies $u \geq \mathfrak{J}$, and if whenever a vector v belongs to the subdifferential of u at x ,



there holds $H(x, v) \geq 0$.

Both are linked by $\nabla \varphi(x) = v$. Extension to viscosity in measure spaces?

Notations

Let μ, ν be two probability measures on \mathbb{R}^d . If $f : \mathbb{R}^d \rightarrow Y$ is measurable, the pushforward $f\#\mu$ is a measure on Y given by $(f\#\mu)(A) = \mu(f^{-1}(A))$ for any measurable $A \subset Y$.

Notations

Let μ, ν be two probability measures on \mathbb{R}^d . If $f : \mathbb{R}^d \rightarrow Y$ is measurable, the pushforward $f\#\mu$ is a measure on Y given by $(f\#\mu)(A) = \mu(f^{-1}(A))$ for any measurable $A \subset Y$. Denote

$$\Gamma(\mu, \nu) := \left\{ \eta = \eta(x, y) \in \mathcal{P}_2(\mathbb{R}^d \times \mathbb{R}^d) \mid \pi_x\#\eta = \mu, \pi_y\#\eta = \nu \right\}.$$

the possible transport plans between μ and ν . The 2-Wasserstein distance is defined as

$$d_{\mathcal{W}}^2(\mu, \nu) := \inf_{\eta \in \Gamma(\mu, \nu)} \int_{(x, y) \in (\mathbb{R}^d)^2} |x - y|^2 d\eta(x, y).$$

Notations

Let μ, ν be two probability measures on \mathbb{R}^d . If $f : \mathbb{R}^d \rightarrow Y$ is measurable, the pushforward $f\#\mu$ is a measure on Y given by $(f\#\mu)(A) = \mu(f^{-1}(A))$ for any measurable $A \subset Y$. Denote

$$\Gamma(\mu, \nu) := \left\{ \eta = \eta(x, y) \in \mathcal{P}_2(\mathbb{R}^d \times \mathbb{R}^d) \mid \pi_x\#\eta = \mu, \pi_y\#\eta = \nu \right\}.$$

the possible transport plans between μ and ν . The 2-Wasserstein distance is defined as

$$d_{\mathcal{W}}^2(\mu, \nu) := \inf_{\eta \in \Gamma(\mu, \nu)} \int_{(x, y) \in (\mathbb{R}^d)^2} |x - y|^2 d\eta(x, y).$$

Def – Wasserstein space The Wasserstein space $\mathcal{P}_2(\mathbb{R}^d)$ is the set of measures μ such that $d_{\mathcal{W}}^2(\mu, \delta_0)$ is finite, endowed with the Wasserstein distance.

Viscosity solutions in $\mathcal{P}_2(\mathbb{R}^d)$

Using **Lions differentiability**, introduced in [Lio07].

- Represent any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ as the law of a set of random variables $X \in L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d)$, that is, $\mu = X \# \mathbb{P}$. Then any function $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ can be *lifted* in

$$U : L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d), \quad U(X) := u(X \# \mathbb{P}).$$

Viscosity solutions in $\mathcal{P}_2(\mathbb{R}^d)$

Using **Lions differentiability**, introduced in [Lio07].

- Represent any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ as the law of a set of random variables $X \in L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d)$, that is, $\mu = X \# \mathbb{P}$. Then any function $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ can be *lifted* in

$$U : L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d), \quad U(X) := u(X \# \mathbb{P}).$$

The gradient of U at X in the Hilbert space $L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d)$ is shown to be of the form $p \circ X$ for some $p \in L^2_{\mu}(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d)$ that depends only on $\mu = X \# \mathbb{P}$.

Viscosity solutions in $\mathcal{P}_2(\mathbb{R}^d)$

Using **Lions differentiability**, introduced in [Lio07].

- Represent any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ as the law of a set of random variables $X \in L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d)$, that is, $\mu = X \# \mathbb{P}$. Then any function $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ can be *lifted* in

$$U : L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d), \quad U(X) := u(X \# \mathbb{P}).$$

The gradient of U at X in the Hilbert space $L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d)$ is shown to be of the form $p \circ X$ for some $p \in L^2_{\mu}(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d)$ that depends only on $\mu = X \# \mathbb{P}$.

- Another equivalent formulation [CD18]: define first *linear derivative* along curves $h \mapsto (1 - h)\mu + h\nu$, then *functional derivative* as the gradient of the linear derivative.

Viscosity solutions in $\mathcal{P}_2(\mathbb{R}^d)$

Using **Lions differentiability**, introduced in [Lio07].

- Represent any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ as the law of a set of random variables $X \in L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d)$, that is, $\mu = X \# \mathbb{P}$. Then any function $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ can be *lifted* in

$$U : L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d), \quad U(X) := u(X \# \mathbb{P}).$$

The gradient of U at X in the Hilbert space $L^2_{\mathbb{P}}(E, \mathcal{E}; \mathbb{R}^d)$ is shown to be of the form $p \circ X$ for some $p \in L^2_{\mu}(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d)$ that depends only on $\mu = X \# \mathbb{P}$.

- Another equivalent formulation [CD18]: define first *linear derivative* along curves $h \mapsto (1 - h)\mu + h\nu$, then *functional derivative* as the gradient of the linear derivative.
- Provides a definition of \mathcal{C}^1 functions and higher derivatives [Sal23], used to obtain existence of “regular” solutions to mean-field games [CDLL19, CP20, MZ22] and viscosity solutions [PW18, BY19, DJS23]...

Viscosity solutions in $\mathcal{P}_2(\mathbb{R}^d)$

Using **semidifferentials** in a well-chosen tangent space [AGS05]. Usually taken as

$$\text{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) := \overline{\{\nabla \varphi \mid \varphi \in \mathcal{C}_c^\infty(\mathbb{R}^d)\}}^{L_\mu^2(\mathbb{R}^d; \mathbb{R}^d)}.$$

Viscosity solutions in $\mathcal{P}_2(\mathbb{R}^d)$

Using **semidifferentials** in a well-chosen tangent space [AGS05]. Usually taken as

$$\text{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) := \overline{\{\nabla \varphi \mid \varphi \in \mathcal{C}_c^\infty(\mathbb{R}^d)\}}^{L_\mu^2(\mathbb{R}^d; \mathbb{R}^d)}.$$

- Possible to define sub/superdifferential and a corresponding notion of viscosity solutions [CQ08], variations in [MQ18, JMQ20, JMQ22, Jim23] with δ -differentials.

Viscosity solutions in $\mathcal{P}_2(\mathbb{R}^d)$

Using **semidifferentials** in a well-chosen tangent space [AGS05]. Usually taken as

$$\text{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) := \overline{\{\nabla \varphi \mid \varphi \in \mathcal{C}_c^\infty(\mathbb{R}^d)\}}^{L_\mu^2(\mathbb{R}^d; \mathbb{R}^d)}.$$

- Possible to define sub/superdifferential and a corresponding notion of viscosity solutions [CQ08], variations in [MQ18, JMQ20, JMQ22, Jim23] with δ -differentials.
- Geometric definition of the **Wasserstein gradient** as the intersection of the sub and superdifferential in [GNT08, GŚ14], shown to be equivalent to the Lions differentiability in [GT19].

Viscosity solutions in $\mathcal{P}_2(\mathbb{R}^d)$

Using **semidifferentials** in a well-chosen tangent space [AGS05]. Usually taken as

$$\text{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) := \overline{\{\nabla \varphi \mid \varphi \in \mathcal{C}_c^\infty(\mathbb{R}^d)\}}^{L_\mu^2(\mathbb{R}^d; \mathbb{R}^d)}.$$

- Possible to define sub/superdifferential and a corresponding notion of viscosity solutions [CQ08], variations in [MQ18, JMQ20, JMQ22, Jim23] with δ -differentials.
- Geometric definition of the **Wasserstein gradient** as the intersection of the sub and superdifferential in [GNT08, GŚ14], shown to be equivalent to the Lions differentiability in [GT19].

Other ideas: applying metric viscosity [AF14, GŚ15], linear derivatives [FN12, BIRS19], pathwise solutions [WZ20, CGK⁺23], directional derivatives [Jer22, JPZ23].

Using directional derivatives

Idea: define the Hamiltonian over a set of functions, as

$$H : \mathbb{T} \rightarrow \mathbb{R},$$

where \mathbb{T} is a set of pairs (μ, p) with $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ and $p : \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$.

Using directional derivatives

Idea: define the Hamiltonian over a set of functions, as

$$H : \mathbb{T} \rightarrow \mathbb{R},$$

where \mathbb{T} is a set of pairs (μ, p) with $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ and $p : \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$.

Typically, $p : \xi \rightarrow D_\mu \varphi(\xi)$ for some $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$, the application of directional derivatives.

For instance,

$$H(\mu, p) := \sup_{u \in U} -p(f(\mu, u)), \quad H(\mu, p) := \sup_{\xi \in \mathbf{Tan}_\mu, \|\xi\|_\mu=1} |p(\xi)|.$$

Using directional derivatives

Idea: define the Hamiltonian over a set of functions, as

$$H : \mathbb{T} \rightarrow \mathbb{R},$$

where \mathbb{T} is a set of pairs (μ, p) with $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ and $p : \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$.

Typically, $p : \xi \rightarrow D_\mu \varphi(\xi)$ for some $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$, the application of directional derivatives. For instance,

$$H(\mu, p) := \sup_{u \in U} -p(f(\mu, u)), \quad H(\mu, p) := \sup_{\xi \in \mathbf{Tan}_\mu, \|\xi\|_\mu=1} |p(\xi)|.$$

- Line opened in [JJZ], developped in [Jer22, JPZ23].

Using directional derivatives

Idea: define the Hamiltonian over a set of functions, as

$$H : \mathbb{T} \rightarrow \mathbb{R},$$

where \mathbb{T} is a set of pairs (μ, p) with $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ and $p : \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$.

Typically, $p : \xi \rightarrow D_\mu \varphi(\xi)$ for some $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$, the application of directional derivatives. For instance,

$$H(\mu, p) := \sup_{u \in U} -p(f(\mu, u)), \quad H(\mu, p) := \sup_{\xi \in \mathbf{Tan}_\mu, \|\xi\|_\mu=1} |p(\xi)|.$$

- Line opened in [JJZ], developped in [Jer22, JPZ23].
- ? Is it possible to reformulate using semidifferentials? ?

Table of Contents

First definitions

Geometric tangent space

Generalized sub and superdifferentials

Definitions of viscosity solutions

The equivalence result


Velocities

Let $T\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times T_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$.

Velocities

Let $T\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times T_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(T\mathbb{R}^d)$ may be endowed with various operations:


$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}),$$



Velocities

Let $T\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times T_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(T\mathbb{R}^d)$ may be endowed with various operations:


$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}),$$



Velocities

Let $\mathbb{T}\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times \mathbb{T}_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(\mathbb{T}\mathbb{R}^d)$ may be endowed with various operations:


$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}), \quad \exp_\mu(h \cdot \xi) := (\pi_x + h \pi_v) \# \xi \quad (\text{exponential}).$$



Velocities

Let $\mathbb{T}\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times \mathbb{T}_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(\mathbb{T}\mathbb{R}^d)$ may be endowed with various operations:



$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}), \quad \exp_\mu(h \cdot \xi) := (\pi_x + h \pi_v) \# \xi \quad (\text{exponential}).$$



Velocities

Let $\mathbb{T}\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times T_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(\mathbb{T}\mathbb{R}^d)$ may be endowed with various operations:



$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}), \quad \exp_\mu(h \cdot \xi) := (\pi_x + h \pi_v) \# \xi \quad (\text{exponential}).$$



Velocities

Let $\mathbb{T}\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times \mathbb{T}_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(\mathbb{T}\mathbb{R}^d)$ may be endowed with various operations:



$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}), \quad \exp_\mu(h \cdot \xi) := (\pi_x + h \pi_v) \# \xi \quad (\text{exponential}).$$



Velocities

Let $\mathbb{T}\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times \mathbb{T}_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(\mathbb{T}\mathbb{R}^d)$ may be endowed with various operations:


$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}), \quad \exp_\mu(h \cdot \xi) := (\pi_x + h \pi_v) \# \xi \quad (\text{exponential}).$$



Velocities

Let $\mathbb{T}\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times T_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(\mathbb{T}\mathbb{R}^d)$ may be endowed with various operations:

$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}), \quad \exp_\mu(h \cdot \xi) := (\pi_x + h \pi_v) \# \xi \quad (\text{exponential}).$$



Velocities

Let $T\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times T_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(T\mathbb{R}^d)$ may be endowed with various operations:

$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}), \quad \exp_\mu(h \cdot \xi) := (\pi_x + h \pi_v) \# \xi \quad (\text{exponential}).$$

Consider the following specific class of transport plans:

$$\Gamma_\mu(\xi, \zeta) := \left\{ \eta \in \mathcal{P}_2 \left(\bigcup_{x \in \mathbb{R}^d} \{x\} \times T_x \mathbb{R}^d \times T_x \mathbb{R}^d \right) \mid (\pi_x, \pi_v) \# \eta = \xi, (\pi_x, \pi_w) \# \eta = \zeta \right\}.$$

Velocities

Let $\mathbb{T}\mathbb{R}^d := \bigcup_{x \in \mathbb{R}^d} \{x\} \times \mathbb{T}_x \mathbb{R}^d$ be the tangent bundle, endowed with $|(x, v)|^2 := |x|^2 + |v|^2$. The set $\mathcal{P}_2(\mathbb{T}\mathbb{R}^d)$ may be endowed with various operations:

$$\lambda \cdot \xi := (\pi_x, \lambda \pi_v) \# \xi \quad (\text{rescaling}), \quad \exp_\mu(h \cdot \xi) := (\pi_x + h \pi_v) \# \xi \quad (\text{exponential}).$$

Consider the following specific class of transport plans:

$$\Gamma_\mu(\xi, \zeta) := \left\{ \eta \in \mathcal{P}_2 \left(\bigcup_{x \in \mathbb{R}^d} \{x\} \times \mathbb{T}_x \mathbb{R}^d \times \mathbb{T}_x \mathbb{R}^d \right) \mid (\pi_x, \pi_v) \# \eta = \xi, (\pi_x, \pi_w) \# \eta = \zeta \right\}.$$

Def Given $\xi, \zeta \in \mathcal{P}_2(\mathbb{T}\mathbb{R}^d)_\mu$, define $W_\mu^2(\xi, \zeta) := \inf_{\eta \in \Gamma_\mu(\xi, \zeta)} \int_{(x, v, w)} |v - w|^2 d\eta$.

Generalized tangent space

Let $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$, and denote

$$\overrightarrow{\mu\nu} := \{(\pi_x, \pi_y - \pi_x) \# \eta \mid \eta \in \Gamma_o(\mu, \nu)\}$$

the set of velocities of geodesics $(\exp_\mu^{-1}(\nu))$.

Generalized tangent space

Let $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$, and denote

$$\overrightarrow{\mu\nu} := \{(\pi_x, \pi_y - \pi_x) \# \eta \mid \eta \in \Gamma_o(\mu, \nu)\}$$

the set of velocities of geodesics $(\exp_\mu^{-1}(\nu))$.

Def The generalized tangent space $\mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d)$ to μ is the set

$$\overline{\{\lambda \cdot \overrightarrow{\mu\nu} \mid \lambda \in \mathbb{R}^+, \nu \in \mathcal{P}_2(\mathbb{R}^d)\}}^{W_\mu}.$$

Generalized tangent space

Let $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$, and denote

$$\overrightarrow{\mu\nu} := \{(\pi_x, \pi_y - \pi_x) \# \eta \mid \eta \in \Gamma_o(\mu, \nu)\}$$

the set of velocities of geodesics $(\exp_\mu^{-1}(\nu))$.

Def The generalized tangent space $\mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d)$ to μ is the set

$$\overline{\{\lambda \cdot \overrightarrow{\mu\nu} \mid \lambda \in \mathbb{R}^+, \nu \in \mathcal{P}_2(\mathbb{R}^d)\}}^{W_\mu}.$$

The set \mathbf{Tan}_μ is stable by scaling by a real factor and enjoys a well-defined projection π^μ .

Generalized tangent space

Let $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$, and denote


$$\vec{\mu\nu} := \{(\pi_x, \pi_y - \pi_x) \# \eta \mid \eta \in \Gamma_o(\mu, \nu)\}$$

the set of velocities of geodesics $(\exp_\mu^{-1}(\nu))$.

Def The generalized tangent space $\mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d)$ to μ is the set

$$\overline{\{\lambda \cdot \vec{\mu\nu} \mid \lambda \in \mathbb{R}^+, \nu \in \mathcal{P}_2(\mathbb{R}^d)\}}^{W_\mu}.$$

The set \mathbf{Tan}_μ is stable by scaling by a real factor and enjoys a well-defined projection π^μ .



Convexity property

Def – Horizontal interpolation Let $\xi_0, \xi_1 \in \mathcal{P}_2(\mathbb{T}\mathbb{R}^d)_\mu$, $\beta \in \Gamma_\mu(\xi_0, \xi_1)$ and $t \in [0, 1]$.
Then

$$\xi_t^\beta := (\pi_x, (1-t)\pi_v + t\pi_w) \# \beta \in \mathcal{P}_2(\mathbb{T}\mathbb{R}^d)_\mu.$$

Convexity property

Def – Horizontal interpolation Let $\xi_0, \xi_1 \in \mathcal{P}_2(\mathbb{T}\mathbb{R}^d)_\mu$, $\beta \in \Gamma_\mu(\xi_0, \xi_1)$ and $t \in [0, 1]$.
Then

$$\xi_t^\beta := (\pi_x, (1-t)\pi_v + t\pi_w) \# \beta \in \mathcal{P}_2(\mathbb{T}\mathbb{R}^d)_\mu.$$

By [Gig08], the set $\mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d)$ is horizontally convex.

Convexity property

Def – Horizontal interpolation Let $\xi_0, \xi_1 \in \mathcal{P}_2(\mathbb{TR}^d)_\mu$, $\beta \in \Gamma_\mu(\xi_0, \xi_1)$ and $t \in [0, 1]$. Then

$$\xi_t^\beta := (\pi_x, (1-t)\pi_v + t\pi_w) \# \beta \in \mathcal{P}_2(\mathbb{TR}^d)_\mu.$$

By [Gig08], the set $\mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d)$ is horizontally convex.

Def 1 If $A \subset \mathcal{P}_2(\mathbb{TR}^d)_\mu$, define $\overline{\text{conv}} A$ as the smallest horizontally convex that is closed with respect to W_μ and contains A .

Barycenter

Notice that $\overline{\text{conv}}\{\xi\} \neq \xi$ in general!

Barycenter

Notice that $\overline{\text{conv}}\{\xi\} \neq \xi$ in general!


Proposition – Barycenter

Let $\xi \in \mathcal{P}_2(\mathbb{R}^d)_\mu$, and $\text{Bary}(\xi) \in L^2_\mu(\mathbb{R}^d; \mathbb{R}^d)$ its barycenter, given by

$$\text{Bary}(\xi)(x) = \int_{v \in T_x \mathbb{R}^d} v d\xi_x(v).$$

Then

$$\text{Bary}(\xi) \# \mu \in \overline{\text{conv}}\{\xi\}.$$



Barycenter

Notice that $\overline{\text{conv}}\{\xi\} \neq \xi$ in general!


Proposition – Barycenter

Let $\xi \in \mathcal{P}_2(\mathbb{TR}^d)_\mu$, and $\text{Bary}(\xi) \in L^2_\mu(\mathbb{R}^d; \mathbb{TR}^d)$ its barycenter, given by

$$\text{Bary}(\xi)(x) = \int_{v \in \mathbb{T}_x \mathbb{R}^d} v d\xi_x(v).$$

Then

$$\text{Bary}(\xi) \# \mu \in \overline{\text{conv}}\{\xi\}.$$



Barycenter

Notice that $\overline{\text{conv}}\{\xi\} \neq \xi$ in general!


Proposition – Barycenter

Let $\xi \in \mathcal{P}_2(\mathbb{T}\mathbb{R}^d)_\mu$, and $\text{Bary}(\xi) \in L^2_\mu(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d)$ its barycenter, given by

$$\text{Bary}(\xi)(x) = \int_{v \in \mathbb{T}_x \mathbb{R}^d} v d\xi_x(v).$$

Then

$$\text{Bary}(\xi) \# \mu \in \overline{\text{conv}}\{\xi\}.$$



Barycenter

Notice that $\overline{\text{conv}}\{\xi\} \neq \xi$ in general!


Proposition – Barycenter

Let $\xi \in \mathcal{P}_2(\mathbb{T}\mathbb{R}^d)_\mu$, and $\text{Bary}(\xi) \in L^2_\mu(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d)$ its barycenter, given by

$$\text{Bary}(\xi)(x) = \int_{v \in \mathbb{T}_x \mathbb{R}^d} v d\xi_x(v).$$

Then

$$\text{Bary}(\xi) \# \mu \in \overline{\text{conv}}\{\xi\}.$$



Directional derivatives

For an application $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$, we denote

$$D_\mu \varphi : \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}, \quad D_\mu \varphi(\xi) := \lim_{h \searrow 0} \frac{\varphi(\exp_\mu(h \cdot \xi)) - \varphi(\mu)}{h}.$$

Directional derivatives

For an application $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$, we denote

$$D_\mu \varphi : \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}, \quad D_\mu \varphi(\xi) := \lim_{h \searrow 0} \frac{\varphi(\exp_\mu(h \cdot \xi)) - \varphi(\mu)}{h}.$$

Def – Metric cotangent bundle Let

$$\mathbb{T}_\mu := \left\{ p : \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \left| \begin{array}{l} p \text{ is positively homogeneous and} \\ \text{Lipschitz-continuous w.r.t. } W_\mu. \end{array} \right. \right\}$$

Denote $\mathbb{T} := \bigcup_{\mu \in \mathcal{P}_2(\mathbb{R}^d)} \{\mu\} \times \mathbb{T}_\mu$.

Precise definition of the Hamiltonian

The Hamiltonian is defined as an application

$$H : \mathbb{T} \rightarrow \mathbb{R}.$$

Precise definition of the Hamiltonian

The Hamiltonian is defined as an application

$$H : \mathbb{T} \rightarrow \mathbb{R}.$$

For instance,

$$H(\mu, p) := \sup_{u \in U} -p(\pi^\mu f(\mu, u)).$$

Precise definition of the Hamiltonian

The Hamiltonian is defined as an application

$$H : \mathbb{T} \rightarrow \mathbb{R}.$$

For instance,

$$H(\mu, p) := \sup_{u \in U} -p(\pi^\mu f(\mu, u)).$$

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is directionally differentiable and locally Lipschitz-continuous, then $D_\mu \varphi \in \mathbb{T}_\mu$. This gives meaning to

$$H(\mu, D_\mu \varphi) = 0.$$

Precise definition of the Hamiltonian

The Hamiltonian is defined as an application

$$H : \mathbb{T} \rightarrow \mathbb{R}.$$

For instance,

$$H(\mu, p) := \sup_{u \in U} -p(\pi^\mu f(\mu, u)).$$

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is directionally differentiable and locally Lipschitz-continuous, then $D_\mu \varphi \in \mathbb{T}_\mu$. This gives meaning to

$$H(\mu, D_\mu \varphi) = 0.$$

- Give a notion of viscosity solutions using the semidifferentials of [AF14].

Precise definition of the Hamiltonian

The Hamiltonian is defined as an application

$$H : \mathbb{T} \rightarrow \mathbb{R}.$$

For instance,

$$H(\mu, p) := \sup_{u \in U} -p(\pi^\mu f(\mu, u)).$$

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is directionally differentiable and locally Lipschitz-continuous, then $D_\mu \varphi \in \mathbb{T}_\mu$. This gives meaning to

$$H(\mu, D_\mu \varphi) = 0.$$

- Give a notion of viscosity solutions using the semidifferentials of [AF14].
- Compare it with a notion of viscosity solutions using test functions.

Table of Contents

First definitions

Geometric tangent space

Generalized sub and superdifferentials

Definitions of viscosity solutions

The equivalence result

Pseudo scalar products

Denote $0_\mu := (\pi_x, 0) \# \mu$ and $\|\xi\|_\mu := W_\mu(\xi, 0_\mu)$.

$\mathcal{P}_2(\mathrm{TR}^d)_\mu$ is the set of probabilities on TR^d with base μ , over which $W_\mu(\cdot, \cdot)$ is a distance.

Pseudo scalar products

Denote $0_\mu := (\pi_x, 0) \# \mu$ and $\|\xi\|_\mu := W_\mu(\xi, 0_\mu)$.

Def Given $\xi, \zeta \in \mathcal{P}_2(\mathbb{TR}^d)_\mu$, define

$$\langle \xi, \zeta \rangle_\mu^+ := \frac{1}{2} [\|\xi\|_\mu^2 + \|\zeta\|_\mu^2 - W_\mu^2(\xi, \zeta)] .$$

To ease notations, we also denote $\langle \xi, \zeta \rangle_\mu^- := -\langle -\xi, \zeta \rangle_\mu^+$.

$\mathcal{P}_2(\mathbb{TR}^d)_\mu$ is the set of probabilities on \mathbb{TR}^d with base μ , over which $W_\mu(\cdot, \cdot)$ is a distance.

Pseudo scalar products

Denote $0_\mu := (\pi_x, 0) \# \mu$ and $\|\xi\|_\mu := W_\mu(\xi, 0_\mu)$.

Def Given $\xi, \zeta \in \mathcal{P}_2(\mathbb{TR}^d)_\mu$, define

$$\langle \xi, \zeta \rangle_\mu^+ := \frac{1}{2} [\|\xi\|_\mu^2 + \|\zeta\|_\mu^2 - W_\mu^2(\xi, \zeta)] .$$

To ease notations, we also denote $\langle \xi, \zeta \rangle_\mu^- := -\langle -\xi, \zeta \rangle_\mu^+$.

Expanding the definition of W_μ yields

$$\langle \xi, \zeta \rangle_\mu^+ = \sup_{\eta \in \Gamma_\mu(\xi, \zeta)} \int_{(x, v, w)} \langle v, w \rangle d\eta, \quad \text{and} \quad \langle \xi, \zeta \rangle_\mu^- = \inf_{\eta \in \Gamma_\mu(\xi, \zeta)} \int_{(x, v, w)} \langle v, w \rangle d\eta.$$

$\mathcal{P}_2(\mathbb{TR}^d)_\mu$ is the set of probabilities on \mathbb{TR}^d with base μ , over which $W_\mu(\cdot, \cdot)$ is a distance.

Properties of $\langle \cdot, \cdot \rangle_\mu^\pm$

If $\xi = f \# \mu$ and $\zeta = g \# \mu$ for some $f, g \in L_\mu^2(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d)$, then

$$\langle \xi, \zeta \rangle_\mu^\pm = \int_{x \in \mathbb{R}^d} \int_{x \in \mathbb{R}^d} \langle f(x), g(x) \rangle d\mu(x) = \langle f, g \rangle_{L_\mu^2}.$$

Properties of $\langle \cdot, \cdot \rangle_\mu^\pm$

If $\xi = f \# \mu$ and $\zeta = g \# \mu$ for some $f, g \in L_\mu^2(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d)$, then

$$\langle \xi, \zeta \rangle_\mu^\pm = \int_{x \in \mathbb{R}^d} \int_{x \in \mathbb{R}^d} \langle f(x), g(x) \rangle d\mu(x) = \langle f, g \rangle_{L_\mu^2}.$$

There always holds $\langle \xi, \xi \rangle_\mu^+ = \|\xi\|_\mu^2$. However,

$$\langle \xi, \xi \rangle_\mu^- = \|\xi\|_\mu^2 \quad \Longleftrightarrow \quad \exists f \in L_\mu^2(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d) \quad \text{such that} \quad \xi = f \# \mu.$$

Properties of $\langle \cdot, \cdot \rangle_\mu^\pm$

If $\xi = f \# \mu$ and $\zeta = g \# \mu$ for some $f, g \in L_\mu^2(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d)$, then

$$\langle \xi, \zeta \rangle_\mu^\pm = \int_{x \in \mathbb{R}^d} \int_{x \in \mathbb{R}^d} \langle f(x), g(x) \rangle d\mu(x) = \langle f, g \rangle_{L_\mu^2}.$$

There always holds $\langle \xi, \xi \rangle_\mu^+ = \|\xi\|_\mu^2$. However,

$$\langle \xi, \xi \rangle_\mu^- = \|\xi\|_\mu^2 \quad \Longleftrightarrow \quad \exists f \in L_\mu^2(\mathbb{R}^d; \mathbb{T}\mathbb{R}^d) \quad \text{such that} \quad \xi = f \# \mu.$$

For instance, if $\xi = \frac{1}{2}\delta_{(0,v_0)} + \frac{1}{2}\delta_{(0,v_1)}$, then $\beta := \frac{1}{2}\delta_{(0,v_0,v_1)} + \frac{1}{2}\delta_{(0,v_1,v_0)}$ yields

$$\langle \xi, \xi \rangle_\mu^- \leq \frac{1}{2} \langle v_0, v_1 \rangle + \frac{1}{2} \langle v_1, v_0 \rangle = -1 = -\|\xi\|_\mu^2.$$



Convexity properties

Convexity properties

Proposition 1 Let $\xi_0, \xi_1 \in \mathcal{P}_2(\mathbb{R}^d)_\mu$ and $\beta \in \Gamma_\mu(\xi_0, \xi_1)$. Then for any $\zeta \in \mathcal{P}_2(\mathbb{R}^d)_\mu$,

$[0, 1] : t \mapsto \langle \zeta, \xi_t^\beta \rangle_\mu^+$ is convex, $[0, 1] : t \mapsto \langle \zeta, \xi_t^\beta \rangle_\mu^-$ is concave.

Convexity properties

Proposition 1 Let $\xi_0, \xi_1 \in \mathcal{P}_2(\mathbb{R}^d)_\mu$ and $\beta \in \Gamma_\mu(\xi_0, \xi_1)$. Then for any $\zeta \in \mathcal{P}_2(\mathbb{R}^d)_\mu$,

$$[0, 1] : t \mapsto \langle \zeta, \xi_t^\beta \rangle_\mu^+ \quad \text{is convex,} \quad [0, 1] : t \mapsto \langle \zeta, \xi_t^\beta \rangle_\mu^- \quad \text{is concave.}$$

Let $A, B \subset \mathcal{P}_2(\mathbb{R}^d)_\mu$ be nonempty, horizontally convex and bounded sets, with A compact w.r.t. the topology induced by W_μ . Then

$$\sup_{\alpha \in A} \inf_{\beta \in B} \langle \alpha, \beta \rangle_\mu^\pm = \inf_{\beta \in B} \sup_{\alpha \in A} \langle \alpha, \beta \rangle_\mu^\pm.$$

Fréchet sub and superdifferentials [AF14, Definition 4.7]

Def – Superdifferential Let $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$. An element $\xi \in \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d)$ belongs to the superdifferential of φ at μ , denoted $\partial_\mu^+ \varphi$, if for all $\nu \in \mathcal{P}_2(\mathbb{R}^d)$,

$$\varphi(\nu) - \varphi(\mu) \leq \inf_{\eta \in \overrightarrow{\mu\nu}} \langle \xi, \eta \rangle_\mu^- + o(d_{\mathcal{W}}(\mu, \nu)).$$

Fréchet sub and superdifferentials [AF14, Definition 4.7]

Def – Superdifferential Let $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$. An element $\xi \in \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d)$ belongs to the superdifferential of φ at μ , denoted $\partial_\mu^+ \varphi$, if for all $\nu \in \mathcal{P}_2(\mathbb{R}^d)$,

$$\varphi(\nu) - \varphi(\mu) \leq \inf_{\eta \in \overrightarrow{\mu\nu}} \langle \xi, \eta \rangle_\mu^- + o(d_{\mathcal{W}}(\mu, \nu)).$$

Def – Subdifferential Let $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$. An element $\xi \in \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d)$ belongs to the subdifferential of φ at μ , denoted $\partial_\mu^- \varphi$, if for all $\nu \in \mathcal{P}_2(\mathbb{R}^d)$,

$$\varphi(\nu) - \varphi(\mu) \geq \sup_{\eta \in \overrightarrow{\mu\nu}} \langle \xi, \eta \rangle_\mu^+ + o(d_{\mathcal{W}}(\mu, \nu)).$$

Example

Given $A \subset \mathcal{P}_2(\mathbb{R}^d)_\mu$, let again $\overline{\text{conv}}A$ be the smallest closed set B containing A such that

$$\forall \xi_0, \xi_1 \in B, \quad \forall \beta \in \Gamma_\mu(\xi_0, \xi_1), \quad \forall t \in [0, 1], \quad \xi_t^\beta = (\pi_x, (1-t)\pi_v + t\pi_w) \# \beta \in B.$$

Example

Given $A \subset \mathcal{P}_2(\mathbb{R}^d)_\mu$, let again $\overline{\text{conv}}A$ be the smallest closed set B containing A such that

$$\forall \xi_0, \xi_1 \in B, \quad \forall \beta \in \Gamma_\mu(\xi_0, \xi_1), \quad \forall t \in [0, 1], \quad \xi_t^\beta = (\pi_x, (1-t)\pi_v + t\pi_w) \# \beta \in B.$$

Proposition 2 Let $\varphi : \mu \mapsto d_{\mathcal{W}}^2(\mu, \sigma)$ for some fixed $\sigma \in \mathcal{P}_2(\mathbb{R}^d)$. The superdifferential of φ is everywhere nonempty and given by

$$\partial_\mu^+ \varphi = \overline{\text{conv}} \{-2 \cdot \xi \mid \xi \in \overrightarrow{\mu\sigma}\}.$$

For reference, the gradient of $x \mapsto |x - y|^2$ at x is $2(x - y) = -2(y - x)$.

Table of Contents

First definitions

Geometric tangent space

Generalized sub and superdifferentials

Definitions of viscosity solutions

The equivalence result

Test function spaces

Def – Test functions For any $\mu \in \Omega \subset \mathcal{P}_2(\mathbb{R}^d)$, define

$$\mathcal{T}_{+,\mu} := \left\{ \varphi : \Omega \rightarrow \mathbb{R} \left| \begin{array}{l} \varphi \text{ is lower semicontinuous, directionally differentiable at } \mu, \\ \partial_{\mu}^{+} \varphi \text{ is nonempty, bounded and } D_{\mu} \varphi(\mu)(\cdot) = \inf_{\zeta \in \partial_{\mu}^{+} \varphi} \langle \cdot, \zeta \rangle_{\mu}^{-} . \end{array} \right. \right\}.$$

Test function spaces

Def – Test functions For any $\mu \in \Omega \subset \mathcal{P}_2(\mathbb{R}^d)$, define

$$\mathcal{T}_{+,\mu} := \left\{ \varphi : \Omega \rightarrow \mathbb{R} \left| \begin{array}{l} \varphi \text{ is lower semicontinuous, directionally differentiable at } \mu, \\ \partial_{\mu}^{+} \varphi \text{ is nonempty, bounded and } D_{\mu} \varphi(\mu)(\cdot) = \inf_{\zeta \in \partial_{\mu}^{+} \varphi} \langle \cdot, \zeta \rangle_{\mu}^{-} . \end{array} \right. \right\}.$$

Similarly, $\mathcal{T}_{-,\mu} := -\mathcal{T}_{+,\mu}$.

Test function spaces

Def – Test functions For any $\mu \in \Omega \subset \mathcal{P}_2(\mathbb{R}^d)$, define

$$\mathcal{T}_{+,\mu} := \left\{ \varphi : \Omega \rightarrow \mathbb{R} \left| \begin{array}{l} \varphi \text{ is lower semicontinuous, directionally differentiable at } \mu, \\ \partial_{\mu}^{+} \varphi \text{ is nonempty, bounded and } D_{\mu} \varphi(\mu)(\cdot) = \inf_{\zeta \in \partial_{\mu}^{+} \varphi} \langle \cdot, \zeta \rangle_{\mu}^{-} \end{array} \right. \right\}.$$

Similarly, $\mathcal{T}_{-,\mu} := -\mathcal{T}_{+,\mu}$.

- Does not appeal to the theory of Wasserstein gradient.

Test function spaces

Def – Test functions For any $\mu \in \Omega \subset \mathcal{P}_2(\mathbb{R}^d)$, define

$$\mathcal{T}_{+,\mu} := \left\{ \varphi : \Omega \rightarrow \mathbb{R} \left| \begin{array}{l} \varphi \text{ is lower semicontinuous, directionally differentiable at } \mu, \\ \partial_\mu^+ \varphi \text{ is nonempty, bounded and } D_\mu \varphi(\mu)(\cdot) = \inf_{\zeta \in \partial_\mu^+ \varphi} \langle \cdot, \zeta \rangle_\mu^- \end{array} \right. \right\}.$$

Similarly, $\mathcal{T}_{-,\mu} := -\mathcal{T}_{+,\mu}$.

- Does not appeal to the theory of Wasserstein gradient.
- Retains a link between directional derivatives and semidifferentials.

Examples of test functions

Proposition 3 Let $\sigma \in \mathcal{P}_2(\mathbb{R}^d)$ be fixed. Then the function $d_{\mathcal{W}}^2(\cdot, \sigma)$ belongs to $\mathcal{T}_{+, \mu}$ for any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$.

Examples of test functions

Proposition 3 Let $\sigma \in \mathcal{P}_2(\mathbb{R}^d)$ be fixed. Then the function $d_{\mathcal{W}}^2(\cdot, \sigma)$ belongs to $\mathcal{T}_{+, \mu}$ for any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$.

From [Gig08, Proposition 4.10], there holds that

$$D_{\mu} d_{\mathcal{W}}^2(\cdot, \sigma)(\xi) = \inf_{\eta \in -2 \cdot \overrightarrow{\mu \sigma}} \langle \xi, \eta \rangle_{\mu}^{-}.$$

Examples of test functions

Proposition 3 Let $\sigma \in \mathcal{P}_2(\mathbb{R}^d)$ be fixed. Then the function $d_{\mathcal{W}}^2(\cdot, \sigma)$ belongs to $\mathcal{T}_{+, \mu}$ for any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$.

From [Gig08, Proposition 4.10], there holds that

$$D_{\mu} d_{\mathcal{W}}^2(\cdot, \sigma)(\xi) = \inf_{\eta \in -2 \cdot \vec{\mu} \vec{\sigma}} \langle \xi, \eta \rangle_{\mu}^{-}.$$

Proposition 4 Let $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ and $\zeta \in \mathcal{P}_2(\mathbb{TR}^d)_{\mu}$ be fixed. Then the function $\varphi : \nu \mapsto \inf_{\eta \in \vec{\mu} \vec{\nu}} \langle \eta, \zeta \rangle_{\mu}^{-}$ belongs to $\mathcal{T}_{+, \mu}$, and there holds

$$D_{\mu} \varphi(\xi) = \langle \xi, \zeta \rangle_{\mu}^{-}.$$

The notion of viscosity

Consider the HJB equation

$$H(\mu, D_\mu u(\mu)) = 0 \quad \mu \in \Omega, \quad u(\mu) = \mathfrak{J}(\mu) \quad \mu \in \partial\Omega. \quad (2)$$

Def – Using test functions

A map $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is a subsolution of (2) if it is **u.s.c.**, if $u \leq \mathfrak{J}$ over $\partial\Omega$, and if for any μ and $\varphi \in \mathcal{T}_{+,\mu}$ such that $u - \varphi$ reaches a **maximum** at μ ,

$$H(\mu, D_\mu \varphi) \leq 0.$$

Def – Using semidifferentials

A map $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is a subsolution of (2) if it is **u.s.c.**, if $u \leq \mathfrak{J}$ over $\partial\Omega$, and if for any element $\xi \in \partial_\mu^+ u$,

$$H(\mu, \langle \xi, \cdot \rangle_\mu^-) \leq 0.$$

The notion of viscosity

Consider the HJB equation

$$H(\mu, D_\mu u(\mu)) = 0 \quad \mu \in \Omega, \quad u(\mu) = \mathfrak{J}(\mu) \quad \mu \in \partial\Omega. \quad (2)$$

Def – Using test functions

A map $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is a subsolution of (2) if it is **I.s.c.**, if $u \geq \mathfrak{J}$ over $\partial\Omega$, and if for any μ and $\varphi \in \mathcal{T}_{-, \mu}$ such that $u - \varphi$ reaches a **minimum** at μ ,

$$H(\mu, D_\mu \varphi) \geq 0.$$

Def – Using semidifferentials

A map $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is a subsolution of (2) if it is **I.s.c.**, if $u \geq \mathfrak{J}$ over $\partial\Omega$, and if for any element $\xi \in \partial_\mu^- u$,

$$H(\mu, \langle \xi, \cdot \rangle_\mu^+) \geq 0.$$

Table of Contents

First definitions

Geometric tangent space

Generalized sub and superdifferentials

Definitions of viscosity solutions

The equivalence result

Statement

Assume that $H : \mathbb{T} \rightarrow \mathbb{R}$ satisfies

$$\begin{aligned}
 \forall \varphi \in \mathcal{I}_{+,\mu}, \quad & H(\mu, D_\mu \varphi) \leq \sup_{\xi \in \partial_\mu^+ \varphi} H\left(\mu, \langle \xi, \cdot \rangle_\mu^-\right), \\
 \forall \varphi \in \mathcal{I}_{-,\mu}, \quad & H(\mu, D_\mu \varphi) \geq \inf_{\xi \in \partial_\mu^- \varphi} H\left(\mu, \langle \xi, \cdot \rangle_\mu^+\right).
 \end{aligned}
 \tag{hyp-H}$$

Statement

Assume that $H : \mathbb{T} \rightarrow \mathbb{R}$ satisfies

$$\begin{aligned} \forall \varphi \in \mathcal{I}_{+,\mu}, \quad H(\mu, D_\mu \varphi) &\leq \sup_{\xi \in \partial_\mu^+ \varphi} H(\mu, \langle \xi, \cdot \rangle_\mu^-), \\ \forall \varphi \in \mathcal{I}_{-,\mu}, \quad H(\mu, D_\mu \varphi) &\geq \inf_{\xi \in \partial_\mu^- \varphi} H(\mu, \langle \xi, \cdot \rangle_\mu^+). \end{aligned} \tag{hyp-H}$$

Theorem Assume that (hyp-H) is satisfied. Then a map $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is a viscosity subsolution (resp. supersolution) in the sense of test functions if and only if it is a viscosity subsolution (resp. supersolution) in the sense of semidifferentials.

Statement

Assume that $H : \mathbb{T} \rightarrow \mathbb{R}$ satisfies

$$\begin{aligned} \forall \varphi \in \mathcal{I}_{+,\mu}, \quad H(\mu, D_\mu \varphi) &\leq \sup_{\xi \in \partial_\mu^+ \varphi} H(\mu, \langle \xi, \cdot \rangle_\mu^-), \\ \forall \varphi \in \mathcal{I}_{-,\mu}, \quad H(\mu, D_\mu \varphi) &\geq \inf_{\xi \in \partial_\mu^- \varphi} H(\mu, \langle \xi, \cdot \rangle_\mu^+). \end{aligned} \tag{hyp-H}$$

Theorem Assume that (hyp-H) is satisfied. Then a map $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is a viscosity subsolution (resp. supersolution) in the sense of test functions if and only if it is a viscosity subsolution (resp. supersolution) in the sense of semidifferentials.

- Given an element $\zeta \in \partial_\mu^+ u$, build a test function φ such that $D_\mu \varphi(\xi) = \langle \xi, \zeta \rangle_\mu^-$.

Statement

Assume that $H : \mathbb{T} \rightarrow \mathbb{R}$ satisfies

$$\begin{aligned} \forall \varphi \in \mathcal{I}_{+, \mu}, \quad H(\mu, D_\mu \varphi) &\leq \sup_{\xi \in \partial_\mu^+ \varphi} H(\mu, \langle \xi, \cdot \rangle_\mu^-), \\ \forall \varphi \in \mathcal{I}_{-, \mu}, \quad H(\mu, D_\mu \varphi) &\geq \inf_{\xi \in \partial_\mu^- \varphi} H(\mu, \langle \xi, \cdot \rangle_\mu^+). \end{aligned} \tag{hyp-H}$$

Theorem Assume that (hyp-H) is satisfied. Then a map $u : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is a viscosity subsolution (resp. supersolution) in the sense of test functions if and only if it is a viscosity subsolution (resp. supersolution) in the sense of semidifferentials.

- Given an element $\zeta \in \partial_\mu^+ u$, build a test function φ such that $D_\mu \varphi(\xi) = \langle \xi, \zeta \rangle_\mu^-$.
- Given a test function, use the representation of $D_\mu \varphi$ and (hyp-H).

Examples of applications

- **Eikonal-type Hamiltonians** Let $\kappa : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be nondecreasing.

$$H : \mathbb{T} \rightarrow \mathbb{R}, \quad H(\mu, p) := \sup_{\xi \in \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d), \|\xi\|_\mu=1} \kappa(|p(\xi)|).$$

Examples of applications

- **Eikonal-type Hamiltonians** Let $\kappa : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be nondecreasing.

$$H : \mathbb{T} \rightarrow \mathbb{R}, \quad H(\mu, p) := \sup_{\xi \in \mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d), \|\xi\|_\mu=1} \kappa(|p(\xi)|).$$

- **“Concave-convex” Hamiltonians** Let F_1 and $F_2 : \mathcal{P}_2(\mathbb{R}^d) \rightrightarrows \mathcal{P}_2(\mathbb{R}^d)$ be set-valued maps such that for any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ and $i \in \{1, 2\}$, $F_i[\mu]$ is a nonempty, horizontally convex and compact subset of $\mathbf{Tan}_\mu \mathcal{P}_2(\mathbb{R}^d)$ endowed with W_μ .

$$H : \mathbb{T} \rightarrow \mathbb{R}, \quad H(\mu, p) := \sup_{\xi_1 \in F_1[\mu]} -p(\xi_1) + \inf_{\xi_2 \in F_2[\mu]} -p(\xi_2).$$

Conclusion and perspectives

Conclusion

- Possibility to use explicit test functions built from the squared Wasserstein distance or pseudo scalar products.

Conclusion and perspectives

Conclusion

- Possibility to use explicit test functions built from the squared Wasserstein distance or pseudo scalar products.
- Equivalence between two notions of viscosity solutions under an explicit condition over the Hamiltonian.

Conclusion and perspectives

Conclusion

- Possibility to use explicit test functions built from the squared Wasserstein distance or pseudo scalar products.
- Equivalence between two notions of viscosity solutions under an explicit condition over the Hamiltonian.

Perspectives

- Extension over $\mathcal{P}_2(\mathcal{N})$, where \mathcal{N} is not Hilbertian (network structure).

Conclusion and perspectives

Conclusion

- Possibility to use explicit test functions built from the squared Wasserstein distance or pseudo scalar products.
- Equivalence between two notions of viscosity solutions under an explicit condition over the Hamiltonian.

Perspectives

- Extension over $\mathcal{P}_2(\mathcal{N})$, where \mathcal{N} is not Hilbertian (network structure).
- Link with Lions differentiability?

Thank you!

- [AF14] Luigi Ambrosio and Jin Feng.
On a class of first order Hamilton–Jacobi equations in metric spaces.
Journal of Differential Equations, 256(7):2194–2245, April 2014.
- [AGS05] Luigi Ambrosio, Nicola Gigli, and Guiseppe Savaré.
Gradient Flows.
Lectures in Mathematics ETH Zürich. Birkhäuser-Verlag, Basel, 2005.
- [BIRS19] Matteo Burzoni, Vincenzo Ignazio, A. Max Reppen, and H. Mete Soner.
Viscosity Solutions for Controlled McKean-Vlasov Jump-Diffusions.
2019.
- [BY19] Alain Bensoussan and Sheung Chi Phillip Yam.
Control problem on space of random variables and master equation.
ESAIM: Control, Optimisation and Calculus of Variations, 25:10, 2019.
- [CD18] René Carmona and François Delarue.
Probabilistic Theory of Mean Field Games with Applications I, volume 83 of *Probability Theory and Stochastic Modelling*.
Springer International Publishing, 2018.

- [CDLL19] Pierre Cardaliaguet, François Delarue, Jean-Michel Lasry, and Pierre-Louis Lions.
The Master Equation and the Convergence Problem in Mean Field Games.
 Number 201 in Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2019.

- [CGK⁺23] Andrea Cosso, Fausto Gozzi, Idris Kharroubi, Huyền Pham, and Mauro Rosestolato.
 Optimal control of path-dependent McKean–Vlasov SDEs in infinite-dimension.
The Annals of Applied Probability, 33(4):2863–2918, August 2023.

- [CP20] Pierre Cardaliaguet and Alessio Porretta.
 An Introduction to Mean Field Game Theory.
 In *Mean Field Games: Cetraro, Italy 2019*, Lecture Notes in Mathematics, pages 1–158. Springer International Publishing, 2020.

- [CQ08] P. Cardaliaguet and M. Quincampoix.
 Deterministic differential games under probability knowledge of initial condition.
International Game Theory Review, 10(01):1–16, March 2008.

- [DJS23] Samuel Daudin, Joe Jackson, and Benjamin Seeger.
 Well-posedness of Hamilton-Jacobi equations in the Wasserstein space: Non-convex Hamiltonians and common noise, December 2023.
 Preprint (arXiv:2312.02324).

- [FN12] Jin Feng and Truyen Nguyen.
Hamilton–Jacobi equations in space of measures associated with a system of conservation laws.
Journal de Mathématiques Pures et Appliquées, 97(4):318–390, April 2012.
- [Gig08] Nicola Gigli.
On the Geometry of the Space of Probability Measures Endowed with the Quadratic Optimal Transport Distance.
PhD thesis, Scuola Normale Superiore di Pisa, Pisa, 2008.
- [GNT08] Wilfrid Gangbo, Truyen Nguyen, and Adrian Tudorascu.
Hamilton–Jacobi Equations in the Wasserstein Space.
Methods and Applications of Analysis, 15(2):155–184, 2008.
- [GŚ14] Wilfrid Gangbo and Andrzej Świech.
Optimal transport and large number of particles.
Discrete and Continuous Dynamical Systems, 34(4):1397–1441, 2014.
- [GŚ15] Wilfrid Gangbo and Andrzej Świech.
Existence of a solution to an equation arising from the theory of Mean Field Games.
Journal of Differential Equations, 259(11):6573–6643, December 2015.
- [GT19] Wilfrid Gangbo and Adrian Tudorascu.
On differentiability in the Wasserstein space and well-posedness for Hamilton–Jacobi equations.
Journal de Mathématiques Pures et Appliquées, 125:119–174, May 2019.

- [Jer22] Othmane Jerhaoui.
Viscosity Theory of First Order Hamilton Jacobi Equations in Some Metric Spaces.
 PhD thesis, Institut Polytechnique de Paris, Paris, 2022.
- [Jim23] Chloé Jimenez.
 Equivalence between strict viscosity solution and viscosity solution in the space of Wasserstein and regular extension of the Hamiltonian in $L^2_{\mathbb{P}}$.
 November 2023.
- [JJZ] Frédéric Jean, Othmane Jerhaoui, and Hasnaa Zidani.
 Deterministic optimal control on Riemannian manifolds under probability knowledge of the initial condition.
 Preprint, available at <https://ensta-paris.hal.science/hal-03564787>.
- [JMQ20] Chloé Jimenez, Antonio Marigonda, and Marc Quincampoix.
 Optimal control of multiagent systems in the Wasserstein space.
Calculus of Variations and Partial Differential Equations, 59, March 2020.
- [JMQ22] Chloé Jimenez, Antonio Marigonda, and Marc Quincampoix.
 Dynamical systems and Hamilton-Jacobi-Bellman equations on the Wasserstein space and their L^2 representations.
Journal of Mathematical Analysis (SIMA), 2022.
 (In press).

- [JPZ23] Othmane Jerhaoui, Averil Prost, and Hasnaa Zidani.
Viscosity solutions of centralized control problems in measure spaces, 2023.
Preprint, available at <https://hal.science/hal-04335852>.
- [Lio07] Pierre-Louis Lions.
Jeux à champ moyen, 2006/2007.
Conférences au Collège de France.
- [MQ18] Antonio Marigonda and Marc Quincampoix.
Mayer control problem with probabilistic uncertainty on initial positions.
Journal of Differential Equations, 264(5):3212–3252, March 2018.
- [MZ22] Chenchen Mou and Jianfeng Zhang.
Wellposedness of Second Order Master Equations for Mean Field Games with Nonsmooth Data, 2022.
Preprint (arXiv:1903.09907).
- [PW18] Huyên Pham and Xiaoli Wei.
Bellman equation and viscosity solutions for mean-field stochastic control problem.
ESAIM: Control, Optimisation and Calculus of Variations, 24(1):437–461, January 2018.
- [Sal23] William Salkeld.
Higher order Lions-Taylor expansions, March 2023.
Preprint (arXiv:2303.17571).

- [WZ20] Cong Wu and Jianfeng Zhang.
Viscosity Solutions to Parabolic Master Equations and McKean-Vlasov SDEs with Closed-loop Controls.
The Annals of Applied Probability, 30(2):936–986, 2020.