

Tropical heat

The eikonal equation as a $(\max,+)$ version of the Poisson equation

Averil Prost

January 30, 2024

LMI-LMRS Doctoral seminar

INSA



anr®

Table of Contents

The $(\max, +)$ semialgebra

Link with linear algebra

An application: finite-dimensional $(\max, +)$ system

The heat equation

The $(\max, +)$ idempotent calculus

Let $\Omega := \mathbb{R} \cup \{-\infty\}$.

Def 1 – Operations For $a, b \in \Omega$, define

$$a \oplus b := \max(a, b), \quad a \otimes b := a + b.$$

The $(\max, +)$ idempotent calculus

Let $\Omega := \mathbb{R} \cup \{-\infty\}$.

Def 1 – Operations For $a, b \in \Omega$, define

$$a \oplus b := \max(a, b), \quad a \otimes b := a + b.$$

Both operations are commutative and associative, and

$$a \otimes (b \oplus c)$$

The $(\max, +)$ idempotent calculus

Let $\Omega := \mathbb{R} \cup \{-\infty\}$.

Def 1 – Operations For $a, b \in \Omega$, define

$$a \oplus b := \max(a, b), \quad a \otimes b := a + b.$$

Both operations are commutative and associative, and

$$a \otimes (b \oplus c) = a + \max(b, c)$$

The $(\max, +)$ idempotent calculus

Let $\Omega := \mathbb{R} \cup \{-\infty\}$.

Def 1 – Operations For $a, b \in \Omega$, define

$$a \oplus b := \max(a, b), \quad a \otimes b := a + b.$$

Both operations are commutative and associative, and

$$a \otimes (b \oplus c) = a + \max(b, c) = \max(a + b, a + c)$$

The $(\max, +)$ idempotent calculus

Let $\Omega := \mathbb{R} \cup \{-\infty\}$.

Def 1 – Operations For $a, b \in \Omega$, define

$$a \oplus b := \max(a, b), \quad a \otimes b := a + b.$$

Both operations are commutative and associative, and

$$a \otimes (b \oplus c) = a + \max(b, c) = \max(a + b, a + c) = (a \otimes b) \oplus (a \otimes c).$$

"Idempotent" since $a \oplus a = a$.

The $(\max, +)$ idempotent calculus

Let $\Omega := \mathbb{R} \cup \{-\infty\}$.

Def 1 – Operations For $a, b \in \Omega$, define

$$a \oplus b := \max(a, b), \quad a \otimes b := a + b.$$

Both operations are commutative and associative, and

$$a \otimes (b \oplus c) = a + \max(b, c) = \max(a + b, a + c) = (a \otimes b) \oplus (a \otimes c).$$

"Idempotent" since $a \oplus a = a$. Define $\mathbb{0} := -\infty$ and $\mathbb{1} := 0$. Then

$$\mathbb{0} \oplus a = \max(-\infty, a) = a,$$

The $(\max, +)$ idempotent calculus

Let $\Omega := \mathbb{R} \cup \{-\infty\}$.

Def 1 – Operations For $a, b \in \Omega$, define

$$a \oplus b := \max(a, b), \quad a \otimes b := a + b.$$

Both operations are commutative and associative, and

$$a \otimes (b \oplus c) = a + \max(b, c) = \max(a + b, a + c) = (a \otimes b) \oplus (a \otimes c).$$

"Idempotent" since $a \oplus a = a$. Define $\mathbb{0} := -\infty$ and $\mathbb{1} := 0$. Then

$$\mathbb{0} \oplus a = \max(-\infty, a) = a, \quad \mathbb{0} \otimes a = -\infty + a = \mathbb{0},$$

The $(\max, +)$ idempotent calculus

Let $\Omega := \mathbb{R} \cup \{-\infty\}$.

Def 1 – Operations For $a, b \in \Omega$, define

$$a \oplus b := \max(a, b), \quad a \otimes b := a + b.$$

Both operations are commutative and associative, and

$$a \otimes (b \oplus c) = a + \max(b, c) = \max(a + b, a + c) = (a \otimes b) \oplus (a \otimes c).$$

"Idempotent" since $a \oplus a = a$. Define $\mathbb{0} := -\infty$ and $\mathbb{1} := 0$. Then

$$\mathbb{0} \oplus a = \max(-\infty, a) = a, \quad \mathbb{0} \otimes a = -\infty + a = \mathbb{0}, \quad \mathbb{1} \otimes a = a + 0 = a.$$

The $(\max, +)$ idempotent calculus

Let $\Omega := \mathbb{R} \cup \{-\infty\}$.

Def 1 – Operations For $a, b \in \Omega$, define

$$a \oplus b := \max(a, b), \quad a \otimes b := a + b.$$

Both operations are commutative and associative, and

$$a \otimes (b \oplus c) = a + \max(b, c) = \max(a + b, a + c) = (a \otimes b) \oplus (a \otimes c).$$

"Idempotent" since $a \oplus a = a$. Define $\mathbb{0} := -\infty$ and $\mathbb{1} := 0$. Then

$$\mathbb{0} \oplus a = \max(-\infty, a) = a, \quad \mathbb{0} \otimes a = -\infty + a = \mathbb{0}, \quad \mathbb{1} \otimes a = a + 0 = a.$$

Then $(\Omega, \oplus, \otimes, \mathbb{0}, \mathbb{1})$ is a semiring (ring without additive inverse).

Integrals

Endow Ω with the application $\mathfrak{d}(a, b) := |e^a - e^b|$, and let $f : \Omega \rightarrow \Omega$ be continuous. Then

$$\sum_{i \in \mathbb{Z} \cup \{-\infty\}}^{\oplus} f(hi) = \max_{i \in \mathbb{Z} \cup \{-\infty\}} f(hi) \xrightarrow{h \searrow 0} \sup_{x \in \Omega = \mathbb{R} \cup \{-\infty\}} f(x).$$

Integrals

Endow Ω with the application $\mathfrak{d}(a, b) := |e^a - e^b|$, and let $f : \Omega \rightarrow \Omega$ be continuous. Then

$$\sum_{i \in \mathbb{Z} \cup \{-\infty\}}^{\oplus} f(hi) = \max_{i \in \mathbb{Z} \cup \{-\infty\}} f(hi) \xrightarrow{h \searrow 0} \sup_{x \in \Omega = \mathbb{R} \cup \{-\infty\}} f(x).$$

Def 2 – Integral Define

$$\int_{x \in \Omega}^{\oplus} f(x) := \sup_{x \in \Omega} f(x).$$

Integrals

Endow Ω with the application $\mathsf{d}(a, b) := |e^a - e^b|$, and let $f : \Omega \rightarrow \Omega$ be continuous. Then

$$\sum_{i \in \mathbb{Z} \cup \{-\infty\}}^{\oplus} f(hi) = \max_{i \in \mathbb{Z} \cup \{-\infty\}} f(hi) \xrightarrow{h \searrow 0} \sup_{x \in \Omega = \mathbb{R} \cup \{-\infty\}} f(x).$$

Def 2 – Integral Define

$$\int_{x \in \Omega}^{\oplus} f(x) := \sup_{x \in \Omega} f(x).$$

In particular, the scalar product becomes

$$\langle f, g \rangle_{\oplus} = \int_{x \in \Omega}^{\oplus} f(x) \otimes g(x) = \sup_{x \in \Omega} f(x) + g(x).$$

People and motivation

- Several names: tropical analysis, idempotent or $(\max, +)$ algebra.

People and motivation

- Several names: tropical analysis, idempotent or $(\max, +)$ algebra.
- 1997: book from Kolokolstov and Maslov [KM97]

People and motivation

- Several names: tropical analysis, idempotent or $(\max, +)$ algebra.
- 1997: book from Kolokolstov and Maslov [KM97]
- 2006: numerical book from McEneaney [McE06]

People and motivation

- Several names: tropical analysis, idempotent or $(\max, +)$ algebra.
- 1997: book from Kolokolstov and Maslov [KM97]
- 2006: numerical book from McEneaney [McE06]
- In France: Marianne Akian and Stéphane Gaubert ([Aki07], [numerical talk](#))

People and motivation

- Several names: tropical analysis, idempotent or $(\max, +)$ algebra.
- 1997: book from Kolokolstov and Maslov [KM97]
- 2006: numerical book from McEneaney [McE06]
- In France: Marianne Akian and Stéphane Gaubert ([Aki07], [numerical talk](#))
- Currently a topic in image processing (morphological operations, [Ang15, Ang22]) and physics ([de]quantization, [Lit12]).

People and motivation

- Several names: tropical analysis, idempotent or $(\max, +)$ algebra.
- 1997: book from Kolokolstov and Maslov [KM97]
- 2006: numerical book from McEneaney [McE06]
- In France: Marianne Akian and Stéphane Gaubert ([Aki07], [numerical talk](#))
- Currently a topic in image processing (morphological operations, [Ang15, Ang22]) and physics ([de]quantization, [Lit12]).

Motivation Ease the study of **optimization problems** by directly working with the "natural" operations of (here) **maximization and sum** of gains.

Table of Contents

The $(\max, +)$ semialgebra

Link with linear algebra

An application: finite-dimensional $(\max, +)$ system

The heat equation

An elementary result

Proposition –  Let $f > g$. Then

$$\lim_{h \searrow 0} h \log \left(e^{f/h} + e^{g/h} \right) = \lim_{h \searrow 0} h \log \left(e^{f/h} - e^{g/h} \right) = f.$$

An elementary result

Proposition –  Let $f > g$. Then

$$\lim_{h \searrow 0} h \log \left(e^{f/h} + e^{g/h} \right) = \lim_{h \searrow 0} h \log \left(e^{f/h} - e^{g/h} \right) = f.$$

Readily visible by noticing that

$$e^{f/h} \pm e^{g/h} = e^{f/h} \left(1 \pm e^{(g-f)/h} \right).$$

An elementary result

Proposition –  Let $f > g$. Then

$$\lim_{h \searrow 0} h \log \left(e^{f/h} + e^{g/h} \right) = \lim_{h \searrow 0} h \log \left(e^{f/h} - e^{g/h} \right) = f.$$

Readily visible by noticing that

$$e^{f/h} \pm e^{g/h} = e^{f/h} \left(1 \pm e^{(g-f)/h} \right).$$

Going further, for any upper bounded and continuous f , there holds

$$\lim_{h \searrow 0} h \log \left(\int_{x \in \mathbb{R}^d} \exp \left(\frac{f(x)}{h} \right) dx \right) = \sup_{x \in \mathbb{R}^d} f(x).$$

Logarithm transform

Def 3 – Logarithm trick For $h > 0$, consider the operations

$$a \otimes_h b := h \log \left(e^{a/h} \cdot e^{b/h} \right), \quad a \oplus_h b := h \log(e^{a/h} + e^{b/h}).$$

Logarithm transform

Def 3 – Logarithm trick For $h > 0$, consider the operations

$$a \otimes_h b := h \log \left(e^{a/h} \cdot e^{b/h} \right), \quad a \oplus_h b := h \log(e^{a/h} + e^{b/h}).$$

Then

$$a \otimes_h b = a + b, \quad a \oplus_h b \xrightarrow[h \searrow 0]{} \max(a, b).$$

Logarithm transform

Def 3 – Logarithm trick For $h > 0$, consider the operations

$$a \otimes_h b := h \log \left(e^{a/h} \cdot e^{b/h} \right), \quad a \oplus_h b := h \log(e^{a/h} + e^{b/h}).$$

Then

$$a \otimes_h b = a + b, \quad a \oplus_h b \xrightarrow[h \searrow 0]{} \max(a, b).$$

We could go further with this game, for instance with

$$a \wedge^\oplus b := \lim_{h \searrow 0} h \log (\exp(a/h) \wedge \exp(b/h))$$

Logarithm transform

Def 3 – Logarithm trick For $h > 0$, consider the operations

$$a \otimes_h b := h \log \left(e^{a/h} \cdot e^{b/h} \right), \quad a \oplus_h b := h \log(e^{a/h} + e^{b/h}).$$

Then

$$a \otimes_h b = a + b, \quad a \oplus_h b \xrightarrow[h \searrow 0]{} \max(a, b).$$

We could go further with this game, for instance with

$$a \hat{\oplus} b := \lim_{h \searrow 0} h \log (\exp(a/h) \wedge \exp(b/h)) = \begin{cases} a & b = 0 \\ \mathbb{I} = 0 & b < 0 \\ +\infty & b > 0. \end{cases}$$

Dialog between $(+, \cdot)$ and $(\max, +)$

$(+, \cdot)$

Classical
operations

Dialog between $(+, \cdot)$ and $(\max, +)$


$(+, \cdot)$

Classical
operations


(\oplus_h, \otimes_h)

Intermediate
operations


Dialog between $(+, \cdot)$ and $(\max, +)$



Dialog between $(+, \cdot)$ and $(\max, +)$



Dialog between $(+, \cdot)$ and $(\max, +)$



Linearity

Let $I : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be linear, and denote

$$\Phi_h(x) := h \log \circ I \circ \exp(x/h).$$

Linearity

Let $I : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be linear, and denote

$$\Phi_h(x) := h \log \circ I \circ \exp(x/h).$$

Here all the operations are understood coordinate by coordinate. Then

$$\Phi_h(a \oplus_h b)$$

Linearity

Let $I : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be linear, and denote

$$\Phi_h(x) := h \log \circ I \circ \exp(x/h).$$

Here all the operations are understood coordinate by coordinate. Then

$$\Phi_h(a \oplus_h b) = h \log \circ I \circ e^{\frac{a \oplus_h b}{h}}$$

Linearity

Let $I : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be linear, and denote

$$\Phi_h(x) := h \log \circ I \circ \exp(x/h).$$

Here all the operations are understood coordinate by coordinate. Then

$$\Phi_h(a \oplus_h b) = h \log \circ I \circ e^{\frac{a \oplus_h b}{h}} = h \log \circ I \left(e^{a/h} + e^{b/h} \right)$$

Linearity

Let $I : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be linear, and denote

$$\Phi_h(x) := h \log \circ I \circ \exp(x/h).$$

Here all the operations are understood coordinate by coordinate. Then

$$\Phi_h(a \oplus_h b) = h \log \circ I \circ e^{\frac{a \oplus_h b}{h}} = h \log \circ I \left(e^{a/h} + e^{b/h} \right) = h \log \left(I e^{a/h} + I e^{b/h} \right)$$

Linearity

Let $I : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be linear, and denote

$$\Phi_h(x) := h \log \circ I \circ \exp(x/h).$$

Here all the operations are understood coordinate by coordinate. Then

$$\begin{aligned}\Phi_h(a \oplus_h b) &= h \log \circ I \circ e^{\frac{a \oplus_h b}{h}} = h \log \circ I \left(e^{a/h} + e^{b/h} \right) = h \log \left(I e^{a/h} + I e^{b/h} \right) \\ &= h \log \left(e^{\frac{\Phi_h(a)}{h}} + e^{\frac{\Phi_h(b)}{h}} \right)\end{aligned}$$

Linearity

Let $I : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be linear, and denote

$$\Phi_h(x) := h \log \circ I \circ \exp(x/h).$$

Here all the operations are understood coordinate by coordinate. Then

$$\begin{aligned}\Phi_h(a \oplus_h b) &= h \log \circ I \circ e^{\frac{a \oplus_h b}{h}} = h \log \circ I \left(e^{a/h} + e^{b/h} \right) = h \log \left(I e^{a/h} + I e^{b/h} \right) \\ &= h \log \left(e^{\frac{\Phi_h(a)}{h}} + e^{\frac{\Phi_h(b)}{h}} \right) = \Phi_h(a) \oplus_h \Phi_h(b).\end{aligned}$$

Linearity

Let $I : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be linear, and denote

$$\Phi_h(x) := h \log \circ I \circ \exp(x/h).$$

Here all the operations are understood coordinate by coordinate. Then

$$\begin{aligned}\Phi_h(a \oplus_h b) &= h \log \circ I \circ e^{\frac{a \oplus_h b}{h}} = h \log \circ I \left(e^{a/h} + e^{b/h} \right) = h \log \left(I e^{a/h} + I e^{b/h} \right) \\ &= h \log \left(e^{\frac{\Phi_h(a)}{h}} + e^{\frac{\Phi_h(b)}{h}} \right) = \Phi_h(a) \oplus_h \Phi_h(b).\end{aligned}$$

Similarly,

$$\Phi_h(a \otimes_h b) = a \otimes_h \Phi_h(b) = \Phi_h(a) \otimes_h b = a \otimes b \otimes \Phi_h(\mathbb{I}).$$

Linearity

Let $I : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be linear, and denote

$$\Phi_h(x) := h \log \circ I \circ \exp(x/h).$$

Here all the operations are understood coordinate by coordinate. Then

$$\begin{aligned}\Phi_h(a \oplus_h b) &= h \log \circ I \circ e^{\frac{a \oplus_h b}{h}} = h \log \circ I \left(e^{a/h} + e^{b/h} \right) = h \log \left(I e^{a/h} + I e^{b/h} \right) \\ &= h \log \left(e^{\frac{\Phi_h(a)}{h}} + e^{\frac{\Phi_h(b)}{h}} \right) = \Phi_h(a) \oplus_h \Phi_h(b).\end{aligned}$$

Similarly,

$$\Phi_h(a \otimes_h b) = a \otimes_h \Phi_h(b) = \Phi_h(a) \otimes_h b = a \otimes b \otimes \Phi_h(\mathbb{I}).$$

The limit operator $\Phi := \lim_{h \searrow 0} \Phi_h$ is $(\max, +)$ -linear.

Example

Let $d = 2$ and $I(x) = Ax$, where $A := \begin{pmatrix} 1 & 1 \\ 2 & 0 \end{pmatrix}$. Then

$$\Phi_h(x) = h \log \circ I \circ \exp(x/h)$$

Example

Let $d = 2$ and $I(x) = Ax$, where $A := \begin{pmatrix} 1 & 1 \\ 2 & 0 \end{pmatrix}$. Then

$$\Phi_h(x) = h \log \circ I \circ \exp(x/h) = h \log \begin{pmatrix} \exp(x_1/h) + \exp(x_2/h) \\ 2 \exp(x_1/h) \end{pmatrix}$$

Example

Let $d = 2$ and $I(x) = Ax$, where $A := \begin{pmatrix} 1 & 1 \\ 2 & 0 \end{pmatrix}$. Then

$$\Phi_h(x) = h \log \circ I \circ \exp(x/h) = h \log \begin{pmatrix} \exp(x_1/h) + \exp(x_2/h) \\ 2 \exp(x_1/h) \end{pmatrix} \xrightarrow[h \searrow 0]{} \begin{pmatrix} \max(x_1, x_2) \\ x_1 \end{pmatrix}.$$

Example

Let $d = 2$ and $I(x) = Ax$, where $A := \begin{pmatrix} 1 & 1 \\ 2 & 0 \end{pmatrix}$. Then

$$\Phi_h(x) = h \log \circ I \circ \exp(x/h) = h \log \begin{pmatrix} \exp(x_1/h) + \exp(x_2/h) \\ 2 \exp(x_1/h) \end{pmatrix} \xrightarrow[h \searrow 0]{} \begin{pmatrix} \max(x_1, x_2) \\ x_1 \end{pmatrix}.$$

Define $\Phi(x) := \begin{pmatrix} \max(x_1, x_2) \\ x_1 \end{pmatrix}$: then for any $\lambda \in \mathbb{R}$, (x_1, x_2) and (y_1, y_2) ,

Example

Let $d = 2$ and $I(x) = Ax$, where $A := \begin{pmatrix} 1 & 1 \\ 2 & 0 \end{pmatrix}$. Then

$$\Phi_h(x) = h \log \circ I \circ \exp(x/h) = h \log \begin{pmatrix} \exp(x_1/h) + \exp(x_2/h) \\ 2 \exp(x_1/h) \end{pmatrix} \xrightarrow[h \searrow 0]{} \begin{pmatrix} \max(x_1, x_2) \\ x_1 \end{pmatrix}.$$

Define $\Phi(x) := \begin{pmatrix} \max(x_1, x_2) \\ x_1 \end{pmatrix}$: then for any $\lambda \in \mathbb{R}$, (x_1, x_2) and (y_1, y_2) ,

$$\Phi(\lambda \otimes x \oplus y) = \Phi \begin{pmatrix} \max(\lambda + x_1, y_1) \\ \max(\lambda + x_2, y_2) \end{pmatrix}$$

Example

Let $d = 2$ and $I(x) = Ax$, where $A := \begin{pmatrix} 1 & 1 \\ 2 & 0 \end{pmatrix}$. Then

$$\Phi_h(x) = h \log \circ I \circ \exp(x/h) = h \log \begin{pmatrix} \exp(x_1/h) + \exp(x_2/h) \\ 2 \exp(x_1/h) \end{pmatrix} \xrightarrow[h \searrow 0]{} \begin{pmatrix} \max(x_1, x_2) \\ x_1 \end{pmatrix}.$$

Define $\Phi(x) := \begin{pmatrix} \max(x_1, x_2) \\ x_1 \end{pmatrix}$: then for any $\lambda \in \mathbb{R}$, (x_1, x_2) and (y_1, y_2) ,

$$\Phi(\lambda \otimes x \oplus y) = \Phi \begin{pmatrix} \max(\lambda + x_1, y_1) \\ \max(\lambda + x_2, y_2) \end{pmatrix} = \begin{pmatrix} \max(\lambda + x_1, y_1, \lambda + x_2, y_2) \\ \max(\lambda + x_1, y_1) \end{pmatrix}$$

Example

Let $d = 2$ and $I(x) = Ax$, where $A := \begin{pmatrix} 1 & 1 \\ 2 & 0 \end{pmatrix}$. Then

$$\Phi_h(x) = h \log \circ I \circ \exp(x/h) = h \log \begin{pmatrix} \exp(x_1/h) + \exp(x_2/h) \\ 2 \exp(x_1/h) \end{pmatrix} \xrightarrow[h \searrow 0]{} \begin{pmatrix} \max(x_1, x_2) \\ x_1 \end{pmatrix}.$$

Define $\Phi(x) := \begin{pmatrix} \max(x_1, x_2) \\ x_1 \end{pmatrix}$: then for any $\lambda \in \mathbb{R}$, (x_1, x_2) and (y_1, y_2) ,

$$\Phi(\lambda \otimes x \oplus y) = \Phi \begin{pmatrix} \max(\lambda + x_1, y_1) \\ \max(\lambda + x_2, y_2) \end{pmatrix} = \begin{pmatrix} \max(\lambda + x_1, y_1, \lambda + x_2, y_2) \\ \max(\lambda + x_1, y_1) \end{pmatrix} = \lambda \otimes \Phi(x) \oplus \varphi(y).$$

Table of Contents

The $(\max, +)$ semialgebra

Link with linear algebra

An application: finite-dimensional $(\max, +)$ system

The heat equation

Setting

Consider the $(\max, +)$ system

$$\Pi \otimes \xi = \beta, \tag{1}$$

where $\Pi \in \mathbb{M}_{2,2}$ is a matrix, $\xi, \beta \in \mathbb{R}^2$ are vectors with β given, and for each $i \in \llbracket 1, 2 \rrbracket$,

$$(\Pi \otimes \xi)_i = \sum_{j \in \llbracket 1, 2 \rrbracket}^{\oplus} \Pi_{ij} \otimes \xi_j = \max_{j \in \llbracket 1, 2 \rrbracket} \Pi_{ij} + \xi_j.$$

Setting

Consider the $(\max, +)$ system

$$\Pi \otimes \xi = \beta, \tag{1}$$

where $\Pi \in \mathbb{M}_{2,2}$ is a matrix, $\xi, \beta \in \mathbb{R}^2$ are vectors with β given, and for each $i \in \llbracket 1, 2 \rrbracket$,

$$(\Pi \otimes \xi)_i = \sum_{j \in \llbracket 1, 2 \rrbracket}^{\oplus} \Pi_{ij} \otimes \xi_j = \max_{j \in \llbracket 1, 2 \rrbracket} \Pi_{ij} + \xi_j.$$

In which cases can we get a solution to (1) by using the link with classical algebra?

Transformation into linear variables

For each $h > 0$, define $P \in \mathbb{M}_{2,2}$ and $x, b \in \mathbb{R}^2$ by

$$P_{ij}^h := \exp(\Pi_{ij}/h), \quad x_j^h := \exp(\xi_j/h), \quad b_i^h := \exp(\beta_i/h).$$

Transformation into linear variables

For each $h > 0$, define $P \in \mathbb{M}_{2,2}$ and $x, b \in \mathbb{R}^2$ by

$$P_{ij}^h := \exp(\Pi_{ij}/h), \quad x_j^h := \exp(\xi_j/h), \quad b_i^h := \exp(\beta_i/h).$$

The linear system $P^h x^h = b^h$ has a solution if $\det(P^h) \neq 0$, i.e.

$$P_{11}^h P_{22}^h - P_{12}^h P_{21}^h \neq 0 \quad \Longleftrightarrow \quad \exp\left(\frac{\Pi_{11} + \Pi_{22}}{h}\right) \neq \exp\left(\frac{\Pi_{12} + \Pi_{21}}{h}\right).$$

Transformation into linear variables

For each $h > 0$, define $P \in \mathbb{M}_{2,2}$ and $x, b \in \mathbb{R}^2$ by

$$P_{ij}^h := \exp(\Pi_{ij}/h), \quad x_j^h := \exp(\xi_j/h), \quad b_i^h := \exp(\beta_i/h).$$

The linear system $P^h x^h = b^h$ has a solution if $\det(P^h) \neq 0$, i.e.

$$P_{11}^h P_{22}^h - P_{12}^h P_{21}^h \neq 0 \quad \Longleftrightarrow \quad \exp\left(\frac{\Pi_{11} + \Pi_{22}}{h}\right) \neq \exp\left(\frac{\Pi_{12} + \Pi_{21}}{h}\right).$$

Assume that $\Pi_{11} + \Pi_{22} \neq \Pi_{12} + \Pi_{21}$, and consider $x^h = (P^h)^{-1} b^h$ solving $P^h x^h = b^h$.

Transformation into linear variables

For each $h > 0$, define $P \in \mathbb{M}_{2,2}$ and $x, b \in \mathbb{R}^2$ by

$$P_{ij}^h := \exp(\Pi_{ij}/h), \quad x_j^h := \exp(\xi_j/h), \quad b_i^h := \exp(\beta_i/h).$$

The linear system $P^h x^h = b^h$ has a solution if $\det(P^h) \neq 0$, i.e.

$$P_{11}^h P_{22}^h - P_{12}^h P_{21}^h \neq 0 \quad \Longleftrightarrow \quad \exp\left(\frac{\Pi_{11} + \Pi_{22}}{h}\right) \neq \exp\left(\frac{\Pi_{12} + \Pi_{21}}{h}\right).$$

Assume that $\Pi_{11} + \Pi_{22} \neq \Pi_{12} + \Pi_{21}$, and consider $x^h = (P^h)^{-1} b^h$ solving $P^h x^h = b^h$. Then

$$\beta = h \log(b^h)$$

Transformation into linear variables

For each $h > 0$, define $P \in \mathbb{M}_{2,2}$ and $x, b \in \mathbb{R}^2$ by

$$P_{ij}^h := \exp(\Pi_{ij}/h), \quad x_j^h := \exp(\xi_j/h), \quad b_i^h := \exp(\beta_i/h).$$

The linear system $P^h x^h = b^h$ has a solution if $\det(P^h) \neq 0$, i.e.

$$P_{11}^h P_{22}^h - P_{12}^h P_{21}^h \neq 0 \quad \Longleftrightarrow \quad \exp\left(\frac{\Pi_{11} + \Pi_{22}}{h}\right) \neq \exp\left(\frac{\Pi_{12} + \Pi_{21}}{h}\right).$$

Assume that $\Pi_{11} + \Pi_{22} \neq \Pi_{12} + \Pi_{21}$, and consider $x^h = (P^h)^{-1} b^h$ solving $P^h x^h = b^h$. Then

$$\beta = h \log(b^h) = h \log(P^h x^h)$$

Transformation into linear variables

For each $h > 0$, define $P \in \mathbb{M}_{2,2}$ and $x, b \in \mathbb{R}^2$ by

$$P_{ij}^h := \exp(\Pi_{ij}/h), \quad x_j^h := \exp(\xi_j/h), \quad b_i^h := \exp(\beta_i/h).$$

The linear system $P^h x^h = b^h$ has a solution if $\det(P^h) \neq 0$, i.e.

$$P_{11}^h P_{22}^h - P_{12}^h P_{21}^h \neq 0 \quad \Longleftrightarrow \quad \exp\left(\frac{\Pi_{11} + \Pi_{22}}{h}\right) \neq \exp\left(\frac{\Pi_{12} + \Pi_{21}}{h}\right).$$

Assume that $\Pi_{11} + \Pi_{22} \neq \Pi_{12} + \Pi_{21}$, and consider $x^h = (P^h)^{-1} b^h$ solving $P^h x^h = b^h$. Then

$$\beta = h \log(b^h) = h \log(P^h x^h) = h \log\left(\sum_{j \in \llbracket 1,2 \rrbracket} \exp\left(\frac{\Pi_{ij} + \xi_j}{h}\right)\right)$$

Transformation into linear variables

For each $h > 0$, define $P \in \mathbb{M}_{2,2}$ and $x, b \in \mathbb{R}^2$ by

$$P_{ij}^h := \exp(\Pi_{ij}/h), \quad x_j^h := \exp(\xi_j/h), \quad b_i^h := \exp(\beta_i/h).$$

The linear system $P^h x^h = b^h$ has a solution if $\det(P^h) \neq 0$, i.e.

$$P_{11}^h P_{22}^h - P_{12}^h P_{21}^h \neq 0 \quad \Longleftrightarrow \quad \exp\left(\frac{\Pi_{11} + \Pi_{22}}{h}\right) \neq \exp\left(\frac{\Pi_{12} + \Pi_{21}}{h}\right).$$

Assume that $\Pi_{11} + \Pi_{22} \neq \Pi_{12} + \Pi_{21}$, and consider $x^h = (P^h)^{-1} b^h$ solving $P^h x^h = b^h$. Then

$$\beta = h \log(b^h) = h \log\left(P^h x^h\right) = h \log\left(\sum_{j \in \llbracket 1,2 \rrbracket} \exp\left(\frac{\Pi_{ij} + \xi_j}{h}\right)\right) \xrightarrow[h \searrow 0]{} \Pi \otimes \xi.$$

Going back

Thus if we can apply $h \log(\cdot)$ to x^h and pass to the limit, we would get a solution.

Going back

Thus if we can apply $h \log(\cdot)$ to x^h and pass to the limit, we would get a solution. Assume

$$\forall i \in \llbracket 1, 2 \rrbracket, \quad x_i^h = \left((P^h)^{-1} b \right)_i \geq 0. \quad (2)$$

Going back

Thus if we can apply $h \log(\cdot)$ to x^h and pass to the limit, we would get a solution. Assume

$$\forall i \in \llbracket 1, 2 \rrbracket, \quad x_i^h = \left((P^h)^{-1} b \right)_i \geq 0. \quad (2)$$

One also obtains that $(x^h)_h$ is bounded uniformly in h . By compactness, (1) admits solutions.

Going back

Thus if we can apply $h \log(\cdot)$ to x^h and pass to the limit, we would get a solution. Assume

$$\forall i \in \llbracket 1, 2 \rrbracket, \quad x_i^h = \left((P^h)^{-1} b \right)_i \geq 0. \quad (2)$$

One also obtains that $(x^h)_h$ is bounded uniformly in h . By compactness, (1) admits solutions.

Example Consider for instance

$$\Pi = \begin{pmatrix} \mathbb{I} & \mathbb{O} \\ \mathbb{O} & \mathbb{I} \end{pmatrix} = \begin{pmatrix} 0 & -\infty \\ -\infty & 0 \end{pmatrix}$$

Going back

Thus if we can apply $h \log(\cdot)$ to x^h and pass to the limit, we would get a solution. Assume

$$\forall i \in \llbracket 1, 2 \rrbracket, \quad x_i^h = \left((P^h)^{-1} b \right)_i \geq 0. \quad (2)$$

One also obtains that $(x^h)_h$ is bounded uniformly in h . By compactness, (1) admits solutions.

Example Consider for instance

$$\Pi = \begin{pmatrix} \mathbb{I} & \mathbb{O} \\ \mathbb{O} & \mathbb{I} \end{pmatrix} = \begin{pmatrix} 0 & -\infty \\ -\infty & 0 \end{pmatrix} \quad \Longrightarrow \quad P = \mathbb{I}_d.$$

Going back

Thus if we can apply $h \log(\cdot)$ to x^h and pass to the limit, we would get a solution. Assume

$$\forall i \in \llbracket 1, 2 \rrbracket, \quad x_i^h = \left((P^h)^{-1} b \right)_i \geq 0. \quad (2)$$

One also obtains that $(x^h)_h$ is bounded uniformly in h . By compactness, (1) admits solutions.

Example Consider for instance

$$\Pi = \begin{pmatrix} \mathbb{I} & \mathbb{O} \\ \mathbb{O} & \mathbb{I} \end{pmatrix} = \begin{pmatrix} 0 & -\infty \\ -\infty & 0 \end{pmatrix} \quad \Longrightarrow \quad P = \mathbb{I}_d.$$

Then $P^{-1} = \mathbb{I}_d$. As (2) is satisfied for all $b = \exp(\beta/h)$, we obtain that $\xi = \beta$ solves (1).

Another example

Consider

$$\Pi = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix}$$

Another example

Consider

$$\Pi = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \quad \Rightarrow \quad P = \begin{pmatrix} e^{1/h} & e^{2/h} \\ e^{4/h} & e^{3/h} \end{pmatrix},$$

Another example

Consider

$$\Pi = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \quad \Rightarrow \quad P = \begin{pmatrix} e^{1/h} & e^{2/h} \\ e^{4/h} & e^{3/h} \end{pmatrix}, \quad P^{-1} = \frac{1}{e^{4/h} - e^{6/h}} \begin{pmatrix} e^{3/h} & -e^{2/h} \\ -e^{4/h} & e^{1/h} \end{pmatrix}.$$

Another example

Consider

$$\Pi = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \quad \Rightarrow \quad P = \begin{pmatrix} e^{1/h} & e^{2/h} \\ e^{4/h} & e^{3/h} \end{pmatrix}, \quad P^{-1} = \frac{1}{e^{4/h} - e^{6/h}} \begin{pmatrix} e^{3/h} & -e^{2/h} \\ -e^{4/h} & e^{1/h} \end{pmatrix}.$$

Since $\det P^h = e^{4/h} - e^{6/h} < 0$, the condition (2) over β becomes

$$e^{\frac{3+\beta_1}{h}} - e^{\frac{4+\beta_2}{h}} \leq 0, \quad \text{and} \quad -e^{\frac{2+\beta_1}{h}} + e^{\frac{1+\beta_2}{h}} \leq 0.$$

Another example

Consider

$$\Pi = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \quad \Rightarrow \quad P = \begin{pmatrix} e^{1/h} & e^{2/h} \\ e^{4/h} & e^{3/h} \end{pmatrix}, \quad P^{-1} = \frac{1}{e^{4/h} - e^{6/h}} \begin{pmatrix} e^{3/h} & -e^{2/h} \\ -e^{4/h} & e^{1/h} \end{pmatrix}.$$

Since $\det P^h = e^{4/h} - e^{6/h} < 0$, the condition (2) over β becomes

$$e^{\frac{3+\beta_1}{h}} - e^{\frac{4+\beta_2}{h}} \leq 0, \quad \text{and} \quad -e^{\frac{2+\beta_1}{h}} + e^{\frac{1+\beta_2}{h}} \leq 0. \quad \text{Equivalently,} \quad |\beta_1 - \beta_2| \leq 1.$$

Another example

Consider

$$\Pi = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \implies P = \begin{pmatrix} e^{1/h} & e^{2/h} \\ e^{4/h} & e^{3/h} \end{pmatrix}, \quad P^{-1} = \frac{1}{e^{4/h} - e^{6/h}} \begin{pmatrix} e^{3/h} & -e^{2/h} \\ -e^{4/h} & e^{1/h} \end{pmatrix}.$$

Since $\det P^h = e^{4/h} - e^{6/h} < 0$, the condition (2) over β becomes

$$e^{\frac{3+\beta_1}{h}} - e^{\frac{4+\beta_2}{h}} \leq 0, \quad \text{and} \quad -e^{\frac{2+\beta_1}{h}} + e^{\frac{1+\beta_2}{h}} \leq 0. \quad \text{Equivalently,} \quad |\beta_1 - \beta_2| \leq 1.$$

Whenever this condition is satisfied, the solution is given by

$$\begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} = \lim_{h \searrow 0} \begin{pmatrix} h \log \left(e^{\frac{4+\beta_2}{h}} - e^{\frac{3+\beta_1}{h}} \right) - h \log (e^{6/h} - e^{4/h}) \\ h \log \left(e^{\frac{2+\beta_1}{h}} - e^{\frac{1+\beta_2}{h}} \right) - h \log (e^{6/h} - e^{4/h}) \end{pmatrix}$$

Another example

Consider

$$\Pi = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \implies P = \begin{pmatrix} e^{1/h} & e^{2/h} \\ e^{4/h} & e^{3/h} \end{pmatrix}, \quad P^{-1} = \frac{1}{e^{4/h} - e^{6/h}} \begin{pmatrix} e^{3/h} & -e^{2/h} \\ -e^{4/h} & e^{1/h} \end{pmatrix}.$$

Since $\det P^h = e^{4/h} - e^{6/h} < 0$, the condition (2) over β becomes

$$e^{\frac{3+\beta_1}{h}} - e^{\frac{4+\beta_2}{h}} \leq 0, \quad \text{and} \quad -e^{\frac{2+\beta_1}{h}} + e^{\frac{1+\beta_2}{h}} \leq 0. \quad \text{Equivalently,} \quad |\beta_1 - \beta_2| \leq 1.$$

Whenever this condition is satisfied, the solution is given by

$$\begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} = \lim_{h \searrow 0} \begin{pmatrix} h \log \left(e^{\frac{4+\beta_2}{h}} - e^{\frac{3+\beta_1}{h}} \right) - h \log (e^{6/h} - e^{4/h}) \\ h \log \left(e^{\frac{2+\beta_1}{h}} - e^{\frac{1+\beta_2}{h}} \right) - h \log (e^{6/h} - e^{4/h}) \end{pmatrix} = \begin{pmatrix} 4 + \beta_2 - 6 \\ 2 + \beta_1 - 6 \end{pmatrix}$$

Another example

Consider

$$\Pi = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \implies P = \begin{pmatrix} e^{1/h} & e^{2/h} \\ e^{4/h} & e^{3/h} \end{pmatrix}, \quad P^{-1} = \frac{1}{e^{4/h} - e^{6/h}} \begin{pmatrix} e^{3/h} & -e^{2/h} \\ -e^{4/h} & e^{1/h} \end{pmatrix}.$$

Since $\det P^h = e^{4/h} - e^{6/h} < 0$, the condition (2) over β becomes

$$e^{\frac{3+\beta_1}{h}} - e^{\frac{4+\beta_2}{h}} \leq 0, \quad \text{and} \quad -e^{\frac{2+\beta_1}{h}} + e^{\frac{1+\beta_2}{h}} \leq 0. \quad \text{Equivalently,} \quad |\beta_1 - \beta_2| \leq 1.$$

Whenever this condition is satisfied, the solution is given by

$$\begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} = \lim_{h \searrow 0} \begin{pmatrix} h \log \left(e^{\frac{4+\beta_2}{h}} - e^{\frac{3+\beta_1}{h}} \right) - h \log (e^{6/h} - e^{4/h}) \\ h \log \left(e^{\frac{2+\beta_1}{h}} - e^{\frac{1+\beta_2}{h}} \right) - h \log (e^{6/h} - e^{4/h}) \end{pmatrix} = \begin{pmatrix} 4 + \beta_2 - 6 \\ 2 + \beta_1 - 6 \end{pmatrix} = \begin{pmatrix} \beta_2 - 2 \\ \beta_1 - 4 \end{pmatrix}.$$

To go further

Of course, the bulky reasoning may be refined.

To go further

Of course, the bulky reasoning may be refined.

- Algorithms to solve linear problems may be transposed in the $(\max, +)$ semialgebra (for instance, Euler scheme \Rightarrow the semi-lagrangian).

To go further


Of course, the bulky reasoning may be refined.

- Algorithms to solve linear problems may be transposed in the $(\max, +)$ semialgebra (for instance, Euler scheme \Rightarrow the semi-lagrangian). Even tropical finite elements [McE06]!

To go further

Of course, the bulky reasoning may be refined.

- Algorithms to solve linear problems may be transposed in the $(\max, +)$ semialgebra (for instance, Euler scheme \Rightarrow the semi-lagrangian). Even tropical finite elements [McE06]!



To go further

Of course, the bulky reasoning may be refined.

- Algorithms to solve linear problems may be transposed in the $(\max, +)$ semialgebra (for instance, Euler scheme \Rightarrow the semi-lagrangian). Even tropical finite elements [McE06]!




Table of Contents

The $(\max, +)$ semialgebra

Link with linear algebra

An application: finite-dimensional $(\max, +)$ system

The heat equation

What about transforming equations?

FIRST ORDER

Consider the first-order transport equation

$$\partial_t u(t, x) - \langle \nabla u(t, x), b(t, x) \rangle = 0.$$

What about transforming equations?

FIRST ORDER

Consider the first-order transport equation

$$\partial_t u(t, x) - \langle \nabla u(t, x), b(t, x) \rangle = 0.$$

Assume that $u \geq 0$, and define

$$v(t, x) = h \log(u(t, x)), \quad \text{i.e.} \quad u(t, x) = \exp(u(t, x)/h).$$

What about transforming equations?

FIRST ORDER Consider the first-order transport equation

$$\partial_t u(t, x) - \langle \nabla u(t, x), b(t, x) \rangle = 0.$$

Assume that $u \geq 0$, and define

$$v(t, x) = h \log(u(t, x)), \quad \text{i.e.} \quad u(t, x) = \exp(v(t, x)/h).$$

Then

$$\partial_t u = \frac{\exp(v/h)}{h} \partial_t v, \quad \nabla u = \frac{\exp(v/h)}{h} \nabla v, \quad (3)$$

What about transforming equations?

FIRST ORDER Consider the first-order transport equation

$$\partial_t u(t, x) - \langle \nabla u(t, x), b(t, x) \rangle = 0.$$

Assume that $u \geq 0$, and define

$$v(t, x) = h \log(u(t, x)), \quad \text{i.e.} \quad u(t, x) = \exp(v(t, x)/h).$$

Then

$$\partial_t u = \frac{\exp(v/h)}{h} \partial_t v, \quad \nabla u = \frac{\exp(v/h)}{h} \nabla v, \quad (3)$$

so that after dividing by $\frac{\exp(v/h)}{h} > 0$, we get again $\partial_t v(t, x) - \langle \nabla v(t, x), b(t, x) \rangle = 0$.

More transformations

SECOND ORDER

Consider now the heat equation

$$\partial_t u(t, x) - h \Delta u(t, x) = 0.$$

More transformations

SECOND ORDER Consider now the heat equation

$$\partial_t u(t, x) - h \Delta u(t, x) = 0.$$

Let again $v = h \log(u)$, i.e. $u = \exp(v/h)$. Then in addition to (3), there holds

$$\Delta u = \frac{\exp(v/t)}{h} \Delta v + \frac{\exp(v/t)}{h^2} |\nabla v|^2.$$

More transformations

SECOND ORDER Consider now the heat equation

$$\partial_t u(t, x) - h \Delta u(t, x) = 0.$$

Let again $v = h \log(u)$, i.e. $u = \exp(v/h)$. Then in addition to (3), there holds

$$\Delta u = \frac{\exp(v/t)}{h} \Delta v + \frac{\exp(v/t)}{h^2} |\nabla v|^2.$$

Hence

$$\frac{\exp(v/h)}{h} \partial_t v - h \left(\frac{\exp(v/t)}{h} \Delta v + \frac{\exp(v/t)}{h^2} |\nabla v|^2 \right) = 0,$$

More transformations

SECOND ORDER Consider now the heat equation

$$\partial_t u(t, x) - h \Delta u(t, x) = 0.$$

Let again $v = h \log(u)$, i.e. $u = \exp(v/h)$. Then in addition to (3), there holds

$$\Delta u = \frac{\exp(v/t)}{h} \Delta v + \frac{\exp(v/t)}{h^2} |\nabla v|^2.$$

Hence

$$\frac{\exp(v/h)}{h} \partial_t v - h \left(\frac{\exp(v/t)}{h} \Delta v + \frac{\exp(v/t)}{h^2} |\nabla v|^2 \right) = 0,$$

or after simplification,

$$\partial_t v(t, x) - h \Delta v(t, x) - |\nabla v(t, x)|^2 = 0.$$

A link with viscosity solutions

Def 4 – (Historical intuition, see [CL83]) The *vanishing viscosity* solution of

$$\partial_t v - |\nabla v|^2 = 0$$

is the limit when h goes to 0 of the (unique) solution of the equation

$$\partial_t v - h\Delta v - |\nabla v|^2 = 0.$$

A link with viscosity solutions

Def 4 – (Historical intuition, see [CL83]) The *vanishing viscosity* solution of

$$\partial_t v - |\nabla v|^2 = 0$$

is the limit when h goes to 0 of the (unique) solution of the equation

$$\partial_t v - h\Delta v - |\nabla v|^2 = 0.$$

It may be characterized by two sign inequalities that maintain the validity of the comparison principle coming from the elliptic perturbation.

The heat kernel

Def 5 – Heat kernel Let C_d be a normalizing constant, and for any $(t, x) \in \mathbb{R}^+ \times \mathbb{R}^d$, define

$$\mu_{t,x}^h := \frac{1}{\sqrt{S_d t h}} \exp \left(-\frac{|\cdot - x|^2}{2th} \right) \mathcal{L}_{\mathbb{R}^d} \in \mathcal{P}_2(\mathbb{R}^d).$$

The heat kernel

Def 5 – Heat kernel Let C_d be a normalizing constant, and for any $(t, x) \in \mathbb{R}^+ \times \mathbb{R}^d$, define

$$\mu_{t,x}^h := \frac{1}{\sqrt{S_d t h}} \exp\left(-\frac{|\cdot - x|^2}{2th}\right) \mathcal{L}_{\mathbb{R}^d} \in \mathcal{P}_2(\mathbb{R}^d).$$

In particular, $\mu_{t,x}^h \xrightarrow[t \searrow 0]{} \delta_x$ narrowly and in the Wasserstein topology.

The heat kernel

Def 5 – Heat kernel Let C_d be a normalizing constant, and for any $(t, x) \in \mathbb{R}^+ \times \mathbb{R}^d$, define

$$\mu_{t,x}^h := \frac{1}{\sqrt{S_d t h}} \exp\left(-\frac{|\cdot - x|^2}{2th}\right) \mathcal{L}_{\mathbb{R}^d} \in \mathcal{P}_2(\mathbb{R}^d).$$

In particular, $\mu_{t,x}^h \xrightarrow[t \searrow 0]{} \delta_x$ narrowly and in the Wasserstein topology.

Proposition – Kernel representation The (weak) solution of the heat equation $\partial_t u(t, x) - h \Delta u(t, x) = 0$ with initial value u_0 is given by

$$u(t, x) = \int_{y \in \mathbb{R}^d} u_0(y) d\mu_{t,x}^h(y).$$

Going through the transformation

Let $u_0 > 0$ upper bounded and continuous.

Going through the transformation

Let $u_0 > 0$ upper bounded and continuous. Denoting again $v = h \log(u)$, and $v_0 = h \log(u_0)$,

$$v(t, x) = h \log \left(\int_{y \in \mathbb{R}^d} \exp \left(\frac{v_0(y)}{h} \right) d\mu_{t,x}^h(y) \right)$$

Going through the transformation

Let $u_0 > 0$ upper bounded and continuous. Denoting again $v = h \log(u)$, and $v_0 = h \log(u_0)$,

$$\begin{aligned} v(t, x) &= h \log \left(\int_{y \in \mathbb{R}^d} \exp \left(\frac{v_0(y)}{h} \right) d\mu_{t,x}^h(y) \right) \\ &= h \log \left(\frac{1}{\sqrt{C_d t h}} \int_{y \in \mathbb{R}^d} \exp \left(\frac{v_0(y)}{h} \right) \exp \left(-\frac{|y-x|^2}{2th} \right) dy \right) \end{aligned}$$

Going through the transformation

Let $u_0 > 0$ upper bounded and continuous. Denoting again $v = h \log(u)$, and $v_0 = h \log(u_0)$,

$$\begin{aligned} v(t, x) &= h \log \left(\int_{y \in \mathbb{R}^d} \exp \left(\frac{v_0(y)}{h} \right) d\mu_{t,x}^h(y) \right) \\ &= h \log \left(\frac{1}{\sqrt{C_d t h}} \int_{y \in \mathbb{R}^d} \exp \left(\frac{v_0(y)}{h} \right) \exp \left(-\frac{|y-x|^2}{2th} \right) dy \right) \\ &= -\frac{h}{2} \log(C_d t h) + h \log \left(\int_{y \in \mathbb{R}^d} \exp \left(\frac{1}{h} \left[v_0(y) - \frac{|y-x|^2}{2t} \right] \right) dy \right) \end{aligned}$$

Going through the transformation

Let $u_0 > 0$ upper bounded and continuous. Denoting again $v = h \log(u)$, and $v_0 = h \log(u_0)$,

$$\begin{aligned} v(t, x) &= h \log \left(\int_{y \in \mathbb{R}^d} \exp \left(\frac{v_0(y)}{h} \right) d\mu_{t,x}^h(y) \right) \\ &= h \log \left(\frac{1}{\sqrt{C_d t h}} \int_{y \in \mathbb{R}^d} \exp \left(\frac{v_0(y)}{h} \right) \exp \left(-\frac{|y-x|^2}{2th} \right) dy \right) \\ &= -\frac{h}{2} \log(C_d t h) + h \log \left(\int_{y \in \mathbb{R}^d} \exp \left(\frac{1}{h} \left[v_0(y) - \frac{|y-x|^2}{2t} \right] \right) dy \right) \\ &\xrightarrow{h \searrow 0} 0 + \sup_{y \in \mathbb{R}^d} \left[v_0(y) - \frac{|y-x|^2}{2t} \right]. \end{aligned}$$

Control interpretation

Let

$$\hat{m}_{t,x} := -\frac{|\cdot - x|^2}{2t}, \quad \hat{\mu}_{t,x}(B) := \sup_{y \in B} \hat{m}_{t,x}(y) \quad \forall B \subset \mathbb{R}^d.$$

Control interpretation

Let

$$\hat{m}_{t,x} := -\frac{|\cdot - x|^2}{2t}, \quad \hat{\mu}_{t,x}(B) := \sup_{y \in B} \hat{m}_{t,x}(y) \quad \forall B \subset \mathbb{R}^d.$$

In the vocabulary of [DMD99], $\hat{m}_{t,x}$ is the *density* of the *Maslov measure* $\hat{\mu}_{t,x}$.

Control interpretation

Let

$$\hat{m}_{t,x} := -\frac{|\cdot - x|^2}{2t}, \quad \hat{\mu}_{t,x}(B) := \sup_{y \in B} \hat{m}_{t,x}(y) \quad \forall B \subset \mathbb{R}^d.$$

In the vocabulary of [DMD99], $\hat{m}_{t,x}$ is the *density* of the *Maslov measure* $\hat{\mu}_{t,x}$.

Proposition – Value function [Lio82] The function

$$V(t, x) := \int_{y \in \mathbb{R}^d}^{\oplus} v_0 \otimes \hat{\mu}_{t,x} = \sup_{y \in \mathbb{R}^d} \left[v_0(y) - \frac{|y - x|^2}{2t} \right] \quad (4)$$

is the unique viscosity solution of the Hamilton-Jacobi equation $\partial_t V - |\nabla V|^2 = 0$ such that $V(0, \cdot) = v_0$.

Control interpretation

Let

$$\hat{m}_{t,x} := -\frac{|\cdot - x|^2}{2t}, \quad \hat{\mu}_{t,x}(B) := \sup_{y \in B} \hat{m}_{t,x}(y) \quad \forall B \subset \mathbb{R}^d.$$

In the vocabulary of [DMD99], $\hat{m}_{t,x}$ is the *density* of the *Maslov measure* $\hat{\mu}_{t,x}$.

Proposition – Value function [Lio82] The function

$$V(t, x) := \int_{y \in \mathbb{R}^d}^{\oplus} v_0 \otimes \hat{\mu}_{t,x} = \sup_{y \in \mathbb{R}^d} \left[v_0(y) - \frac{|y - x|^2}{2t} \right] \quad (4)$$

is the unique viscosity solution of the Hamilton-Jacobi equation $\partial_t V - |\nabla V|^2 = 0$ such that $V(0, \cdot) = v_0$. The formula (4) is known as the Hopf-Lax formula.

Going further

- In the case of the heat equation, **Hopf-Cole** transform. But Lax-Hopf formula valid for a larger class of HJ equations of the type

$$\partial_t V(t, x) + H(x, \nabla V(t, x)) = 0,$$

provided the Hamiltonian H is *concave* in its second variable (+ regularity conditions).

Going further

- In the case of the heat equation, **Hopf-Cole** transform. But Lax-Hopf formula valid for a larger class of HJ equations of the type

$$\partial_t V(t, x) + H(x, \nabla V(t, x)) = 0,$$

provided the Hamiltonian H is *concave* in its second variable (+ regularity conditions).

- Maslov measures may be used to recast the Lax-Hopf semigroup as the conditional expectation of Maslov stochastic processes.

Going further

- In the case of the heat equation, **Hopf-Cole** transform. But Lax-Hopf formula valid for a larger class of HJ equations of the type

$$\partial_t V(t, x) + H(x, \nabla V(t, x)) = 0,$$


provided the Hamiltonian H is *concave* in its second variable (+ regularity conditions).

- Maslov measures may be used to recast the Lax-Hopf semigroup as the conditional expectation of Maslov stochastic processes.
- Using the Hopf-Lax semigroup, Maslov defined weak solution by "duality", in the spirit of

$$\langle u, \varphi \rangle_{\oplus} = \langle u_0, S_t^* \varphi \rangle_{\oplus},$$

where S_t^* is a "dual" semigroup acting on test functions φ [KM97, Definition 3.1].

Conclusion



Thank you!

- [Aki07] Marianne Akian.
Algèbre Max-plus, Applications Monotones Contractantes et Équations Aux Dérivées Partielles : Trois Approches Du Contrôle Optimal.
Habilitation à diriger les recherches, 2007.
- [Ang15] Jesus Angulo.
Morphological Scale-Space Operators for Images Supported on Point Clouds.
In Springer-Verlag Berlin Heidelberg, editor, *5th International Conference on Scale Space and Variational Methods in Computer Vision*, volume LNCS 9087 of *Proc. of SSVM'15 (5th International Conference on Scale Space and Variational Methods in Computer Vision)*, Lège-Cap Ferret, France, June 2015.
- [Ang22] Jesús Angulo.
Morphological Counterpart of Ornstein–Uhlenbeck Semigroups and PDEs.
In Étienne Baudrier, Benoît Naegel, Adrien Krähenbühl, and Mohamed Tajine, editors, *Discrete Geometry and Mathematical Morphology*, Lecture Notes in Computer Science, pages 169–181, Cham, 2022. Springer International Publishing.
- [CL83] Michael G. Crandall and Pierre-Louis Lions.
Viscosity solutions of Hamilton–Jacobi equations.
Transactions of the American Mathematical Society, 277(1):1–42, 1983.

- [DMD99] P. Del Moral and M. Doisy.
Maslov Idempotent Probability Calculus, I.
Theory of Probability & Its Applications, 43(4):562–576, January 1999.
- [KM97] Vassili N. Kolokoltsov and Victor P. Maslov.
Idempotent Analysis and Its Applications.
Springer Netherlands, Dordrecht, 1997.
- [Lio82] P. L. Lions.
Generalized Solutions of Hamilton-Jacobi Equations.
Number 69 in Research Notes in Mathematics. Pitman, Boston, 1982.
- [Lit12] Grigory L. Litvinov.
Idempotent/tropical analysis, the Hamilton-Jacobi and Bellman equations, March 2012.
- [McE06] William M. McEneaney.
Max-plus Methods for Nonlinear Control and Estimation.
Systems and Control. Birkhäuser, Boston, 2006.