

A background image of a man with dark, curly hair and glasses, smiling broadly. He is wearing a dark jacket. The image is faded and serves as a background for the text.

Ekeland

A beginner's point of view on some variational principles

Averil Prost

January 10, 2022
LMI/LMRS doctoral seminar

Table of Contents

The original principle

History

Statement

The proof

What since

History


Hilbert version: Ekeland-Lebourg

Smooth version: Borwein-Preiss


An application

History (1/2)


1974 Ivar Ekeland's *On the variational principle* [Eke74], metric, distances.



Theorem – Ekeland [Eke74] Let (X, d) be a complete metric space. Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded.




Theorem – Ekeland [Eke74] Let (X, d) be a complete metric space. Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let $x \in \text{dom}(f)$, $\delta > 0$.



Theorem – Ekeland [Eke74] Let (X, d) be a complete metric space. Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let $x \in \text{dom}(f)$, $\delta > 0$. Then $\exists y \in X$ s.t.


$$\left\{ \begin{array}{l} f(y) \leq f(x) - \delta d(x, y), \end{array} \right. \quad \begin{array}{l} (1a) \\ (1b) \end{array}$$




Theorem – Ekeland [Eke74] Let (X, d) be a complete metric space. Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let $x \in \text{dom}(f)$, $\delta > 0$. Then $\exists y \in X$ s.t.


$$\begin{cases} f(y) \leq f(x) - \delta d(x, y), \\ f(y) - \delta d(z, y) < f(z) \end{cases} \quad \forall z \in X \setminus \{y\}. \quad (1a)$$

$$(1b)$$




Theorem – Ekeland [Eke74] Let (X, d) be a complete metric space. Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let $x \in \text{dom}(f)$, $\delta > 0$. Then $\exists y \in X$ s.t.


$$\begin{cases} f(y) \leq f(x) - \delta d(x, y), \\ f(y) - \delta d(z, y) < f(z) \end{cases} \quad \begin{matrix} (1a) \\ \forall z \in X \setminus \{y\}. \end{matrix} \quad (1b)$$




 No min when it exists


Theorem – Ekeland [Eke74] Let (X, d) be a complete metric space. Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let $x \in \text{dom}(f)$, $\delta > 0$. Then $\exists y \in X$ s.t.


$$\begin{cases} f(y) \leq f(x) - \delta d(x, y), \\ f(y) - \delta d(z, y) < f(z) \end{cases} \quad \forall z \in X \setminus \{y\}. \quad (1a)$$

$$\forall z \in X \setminus \{y\}. \quad (1b)$$







 No min when it exists

 δ is arbitrary

Theorem – Ekeland [Eke74] Let (X, d) be a complete metric space. Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let $x \in \text{dom}(f)$, $\delta > 0$. Then $\exists y \in X$ s.t.

$$\begin{cases} f(y) \leq f(x) - \delta d(x, y), \\ f(y) - \delta d(z, y) < f(z) \end{cases} \quad \forall z \in X \setminus \{y\}. \quad (1a) \quad (1b)$$




-  No min when it exists
-  δ is arbitrary
-  y stays in X

Theorem – Ekeland [Eke74] Let (X, d) be a complete metric space. Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let $x \in \text{dom}(f)$, $\delta > 0$. Then $\exists y \in X$ s.t.

$$\begin{cases} f(y) \leq f(x) - \delta d(x, y), \\ f(y) - \delta d(z, y) < f(z) \end{cases} \quad \forall z \in X \setminus \{y\}. \quad (1a)$$


$$(1b)$$



- 👎 No min when it exists
- 👍 δ is arbitrary
- 👍 y stays in X
- 👍 \sim no $+\infty$ behavior

Theorem – Ekeland [Eke74] Let (X, d) be a complete metric space. Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let $x \in \text{dom}(f)$, $\delta > 0$. Then $\exists y \in X$ s.t.


$$\begin{cases} f(y) \leq f(x) - \delta d(x, y), \\ f(y) - \delta d(z, y) < f(z) \end{cases} \quad \forall z \in X \setminus \{y\}. \quad (1a) \quad (1b)$$



- 👎 No min when it exists
- 👍 δ is arbitrary
- 👍 y stays in X
- 👍 \sim no $+\infty$ behavior
- 👍 No (local) compactness


The proof (1/3)

Let $S_0 := \text{dom } f$, and $\varepsilon > 0$.




The proof (1/3)

Let $S_0 := \text{dom } f$, and $\varepsilon > 0$. Pick $x_0 \in S_0$ such that $f(x_0) \leq \inf_X f + \varepsilon$.



The proof (1/3)


Let $S_0 := \text{dom } f$, and $\varepsilon > 0$. Pick $x_0 \in S_0$ such that $f(x_0) \leq \inf_X f + \varepsilon$. Build $(x_i, S_i)_i$ as



$$S_i := \{x \in X \mid f(x) \leq f(x_{i-1}) - \delta d(x, x_{i-1})\},$$

The proof (1/3)

Let $S_0 := \text{dom } f$, and $\varepsilon > 0$. Pick $x_0 \in S_0$ such that $f(x_0) \leq \inf_X f + \varepsilon$. Build $(x_i, S_i)_i$ as




$$S_i := \{x \in X \mid f(x) \leq f(x_{i-1}) - \delta d(x, x_{i-1})\},$$

$$\delta d(x, x_{i-1}) \leq f(x_{i-1}) - f(x) \quad \text{✎}$$

The proof (1/3)

Let $S_0 := \text{dom } f$, and $\varepsilon > 0$. Pick $x_0 \in S_0$ such that $f(x_0) \leq \inf_X f + \varepsilon$. Build $(x_i, S_i)_i$ as



$$S_i := \{x \in X \mid f(x) \leq f(x_{i-1}) - \delta d(x, x_{i-1})\},$$


$$\delta d(x, x_{i-1}) \leq f(x_{i-1}) - f(x) \quad \text{✎}$$

and pick $x_i \in S_i$ such that

$$f(x_i) \leq \frac{f(x_{i-1}) + \inf_{y \in S_i} f(y)}{2}.$$

The proof (1/3)

Let $S_0 := \text{dom } f$, and $\varepsilon > 0$. Pick $x_0 \in S_0$ such that $f(x_0) \leq \inf_X f + \varepsilon$. Build $(x_i, S_i)_i$ as



$$S_i := \{x \in X \mid f(x) \leq f(x_{i-1}) - \delta d(x, x_{i-1})\},$$


$$\delta d(x, x_{i-1}) \leq f(x_{i-1}) - f(x) \quad \text{✎}$$

and pick $x_i \in S_i$ such that

$$f(x_i) \leq \frac{f(x_{i-1}) + \inf_{y \in S_i} f(y)}{2}.$$

The proof (1/3)

Let $S_0 := \text{dom } f$, and $\varepsilon > 0$. Pick $x_0 \in S_0$ such that $f(x_0) \leq \inf_X f + \varepsilon$. Build $(x_i, S_i)_i$ as



$$S_i := \{x \in X \mid f(x) \leq f(x_{i-1}) - \delta d(x, x_{i-1})\},$$


$$\delta d(x, x_{i-1}) \leq f(x_{i-1}) - f(x) \quad \text{✎}$$

and pick $x_i \in S_i$ such that

$$f(x_i) \leq \frac{f(x_{i-1}) + \inf_{y \in S_i} f(y)}{2}.$$

The proof (1/3)

Let $S_0 := \text{dom } f$, and $\varepsilon > 0$. Pick $x_0 \in S_0$ such that $f(x_0) \leq \inf_X f + \varepsilon$. Build $(x_i, S_i)_i$ as



$$S_i := \{x \in X \mid f(x) \leq f(x_{i-1}) - \delta d(x, x_{i-1})\},$$

$$\delta d(x, x_{i-1}) \leq f(x_{i-1}) - f(x) \quad \text{✎}$$

and pick $x_i \in S_i$ such that

$$f(x_i) \leq \frac{f(x_{i-1}) + \inf_{y \in S_i} f(y)}{2}.$$

S_i nonempty and closed.

Let us show that $S_{i+1} \subset S_i$, and $\text{diam } S_i \xrightarrow{i \rightarrow \infty} 0$.

The proof (2/3)

Let $x \in S_{i+1}$:

$$f(x) \underset{x \in S_{i+1}}{\leq} f(x_i) - \delta d(x, x_i)$$

The proof (2/3)

Let $x \in S_{i+1}$:

$$\begin{aligned} f(x) &\leq f(x_i) - \delta d(x, x_i) \\ &\leq f(x_{i-1}) - \delta [d(x_i, x_{i-1}) + d(x, x_i)] \end{aligned}$$

The proof (2/3)

Let $x \in S_{i+1}$:

$$\begin{aligned} f(x) &\leq f(x_i) - \delta d(x, x_i) \\ &\leq f(x_{i-1}) - \delta [d(x_i, x_{i-1}) + d(x, x_i)] \\ &\leq f(x_{i-1}) - \delta d(x, x_{i-1}), \\ \triangle \text{ ineq} \end{aligned}$$

The proof (2/3)

Let $x \in S_{i+1}$:

$$\begin{aligned} f(x) &\leq f(x_i) - \delta d(x, x_i) \\ &\leq f(x_{i-1}) - \delta [d(x_i, x_{i-1}) + d(x, x_i)] \\ &\leq f(x_{i-1}) - \delta d(x, x_{i-1}), \\ \triangle \text{ ineq} \end{aligned}$$

and $x \in S_i$, so that $S_{i+1} \subset S_i$.

The proof (2/3)

Let $x \in S_{i+1}$:

$$\begin{aligned} f(x) &\leq f(x_i) - \delta d(x, x_i) \\ &\leq f(x_{i-1}) - \delta [d(x_i, x_{i-1}) + d(x, x_i)] \\ &\leq f(x_{i-1}) - \delta d(x, x_{i-1}), \\ &\triangle \text{ ineq} \end{aligned}$$

and $x \in S_i$, so that $S_{i+1} \subset S_i$.

On the other hand, since $\inf_{S_{i+1}} f \geq \inf_{S_i} f$,

$$f(x_i) - \inf_{S_{i+1}} f \leq [f(x_{i-1}) + \inf_{S_i} f - 2 \inf_{S_{i+1}} f]/2$$

The proof (2/3)

Let $x \in S_{i+1}$:

$$\begin{aligned} f(x) &\leq f(x_i) - \delta d(x, x_i) \\ &\leq f(x_{i-1}) - \delta [d(x_i, x_{i-1}) + d(x, x_i)] \\ &\leq f(x_{i-1}) - \delta d(x, x_{i-1}), \\ &\triangle \text{ ineq} \end{aligned}$$

and $x \in S_i$, so that $S_{i+1} \subset S_i$.

On the other hand, since $\inf_{S_{i+1}} f \geq \inf_{S_i} f$,

$$\begin{aligned} f(x_i) - \inf_{S_{i+1}} f &\leq [f(x_{i-1}) + \inf_{S_i} f - 2 \inf_{S_{i+1}} f]/2 \\ &\leq [f(x_{i-1}) - \inf_{S_i} f]/2 \leq \dots \leq \frac{\varepsilon}{2^i}. \end{aligned}$$

The proof (2/3)

Let $x \in S_{i+1}$:

$$\begin{aligned}
 f(x) &\leq f(x_i) - \delta d(x, x_i) \\
 &\leq f(x_{i-1}) - \delta [d(x_i, x_{i-1}) + d(x, x_i)] \\
 &\triangle \text{ ineq} \leq f(x_{i-1}) - \delta d(x, x_{i-1}),
 \end{aligned}$$

and $x \in S_i$, so that $S_{i+1} \subset S_i$.

On the other hand, since $\inf_{S_{i+1}} f \geq \inf_{S_i} f$,

$$\begin{aligned}
 f(x_i) - \inf_{S_{i+1}} f &\leq [f(x_{i-1}) + \inf_{S_i} f - 2 \inf_{S_{i+1}} f]/2 \\
 &\leq [f(x_{i-1}) - \inf_{S_i} f]/2 \leq \dots \leq \frac{\varepsilon}{2^i}. \\
 \delta d(x, x_i) &\leq f(x_i) - f(x) \leq f(x_i) - \inf_{S_{i+1}} f \leq \frac{\varepsilon}{2^i}.
 \end{aligned}$$

The proof (2/3)

Let $x \in S_{i+1}$:

$$\begin{aligned}
 f(x) &\leq f(x_i) - \delta d(x, x_i) \\
 &\leq f(x_{i-1}) - \delta [d(x_i, x_{i-1}) + d(x, x_i)] \\
 &\leq f(x_{i-1}) - \delta d(x, x_{i-1}), \\
 \triangle \text{ ineq}
 \end{aligned}$$

and $x \in S_i$, so that $S_{i+1} \subset S_i$.

On the other hand, since $\inf_{S_{i+1}} f \geq \inf_{S_i} f$,

$$\begin{aligned}
 f(x_i) - \inf_{S_{i+1}} f &\leq [f(x_{i-1}) + \inf_{S_i} f - 2 \inf_{S_{i+1}} f]/2 \\
 &\leq [f(x_{i-1}) - \inf_{S_i} f]/2 \leq \dots \leq \frac{\varepsilon}{2^i}. \\
 \delta d(x, x_i) &\leq f(x_i) - f(x) \leq f(x_i) - \inf_{S_{i+1}} f \leq \frac{\varepsilon}{2^i}.
 \end{aligned}$$

so that $d(x, x_{i-1}) \leq \frac{\varepsilon}{\delta 2^i}$, and $\text{diam } S_i \xrightarrow{i \rightarrow \infty} 0$.

The proof (2/3)

Let $x \in S_{i+1}$:

$$\begin{aligned} f(x) &\leq f(x_i) - \delta d(x, x_i) \\ &\leq f(x_{i-1}) - \delta [d(x_i, x_{i-1}) + d(x, x_i)] \\ &\leq f(x_{i-1}) - \delta d(x, x_{i-1}), \\ \triangle \text{ ineq} \end{aligned}$$

and $x \in S_i$, so that $S_{i+1} \subset S_i$.


On the other hand, since $\inf_{S_{i+1}} f \geq \inf_{S_i} f$,

$$\begin{aligned} f(x_i) - \inf_{S_{i+1}} f &\leq [f(x_{i-1}) + \inf_{S_i} f - 2 \inf_{S_{i+1}} f]/2 \\ &\leq [f(x_{i-1}) - \inf_{S_i} f]/2 \leq \dots \leq \frac{\varepsilon}{2^i}. \\ \delta d(x, x_i) &\leq f(x_i) - f(x) \leq f(x_i) - \inf_{S_{i+1}} f \leq \frac{\varepsilon}{2^i}. \end{aligned}$$

so that $d(x, x_{i-1}) \leq \frac{\varepsilon}{\delta 2^i}$, and $\text{diam } S_i \xrightarrow{i \rightarrow \infty} 0$.


Since X is closed, by Cantor's intersection theorem, there exists an unique $y \in \bigcap_{i=0}^{\infty} S_i$.

The proof (3/3)




- Since $y \in S_1$, $f(y) \leq f(x_0) - \delta d(y, x_0)$, hence (1a).

The proof (3/3)



- Since $y \in S_1$, $f(y) \leq f(x_0) - \delta d(y, x_0)$, hence (1a).
- Let $x \neq y$, and $i \in \mathbb{N}$ s.t. $x \notin S_{i+1}$.


The proof (3/3)



- Since $y \in S_1$, $f(y) \leq f(x_0) - \delta d(y, x_0)$, hence (1a).
- Let $x \neq y$, and $i \in \mathbb{N}$ s.t. $x \notin S_{i+1}$.

$$f(x) \underset{x \notin S_{i+1}}{>} f(x_i) - \delta d(x, x_i)$$


The proof (3/3)



- Since $y \in S_1$, $f(y) \leq f(x_0) - \delta d(y, x_0)$, hence (1a).
- Let $x \neq y$, and $i \in \mathbb{N}$ s.t. $x \notin S_{i+1}$.

$$\begin{aligned}
 f(x) &>_{x \notin S_{i+1}} f(x_i) - \delta d(x, x_i) \\
 &\geq_{y \in S_{i+1}} f(y) + \delta d(y, x_i) - \delta d(x, x_i),
 \end{aligned}$$

The proof (3/3)



- Since $y \in S_1$, $f(y) \leq f(x_0) - \delta d(y, x_0)$, hence (1a).
- Let $x \neq y$, and $i \in \mathbb{N}$ s.t. $x \notin S_{i+1}$.

$$\begin{aligned}
 f(x) &>_{x \notin S_{i+1}} f(x_i) - \delta d(x, x_i) \\
 &\geq_{y \in S_{i+1}} f(y) + \delta d(y, x_i) - \delta d(x, x_i), \\
 &\geq_{\triangle \text{ ineq.}} f(y) - \delta d(y, x),
 \end{aligned}$$

hence (1b). □

Table of Contents

The original principle

History

Statement

The proof

What since

History


Hilbert version: Ekeland-Lebourg

Smooth version: Borwein-Preiss


An application

History (2/2)

1974 Ivar Ekeland's *On the variational principle* [Eke74], metric, distances.




History (2/2)




1974 Ivar Ekeland's *On the variational principle* [Eke74], metric, distances.

1976 Ekeland and Lebourg [EL76], Banach, linear functions.


History (2/2)




History (2/2)



History (2/2)




History (2/2)



The Hilbertian Ekeland principle

Theorem – Ekeland-Lebourg [EL76] Let a closed bounded $D \subset H$ real Hilbert, and $f : D \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded.




The Hilbertian Ekeland principle

Theorem – Ekeland-Lebourg [EL76] Let a closed bounded $D \subset H$ real Hilbert, and $f : D \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. $\forall \delta > 0, \exists \bar{x} \in D$ and $p \in H'$ s.t.

$$\begin{cases} |p|_{H'} < \delta, & (2a) \end{cases}$$

$$\begin{cases} x \rightarrow f(x) + \langle p, x \rangle_{H',H} \text{ admits a strict minimum over } D \text{ in } \bar{x}. & (2b) \end{cases}$$





The Hilbertian Ekeland principle

Theorem – Ekeland-Lebourg [EL76] Let a closed bounded $D \subset H$ real Hilbert, and $f : D \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. $\forall \delta > 0, \exists \bar{x} \in D$ and $p \in H'$ s.t.

$$\begin{cases} |p|_{H'} < \delta, & (2a) \end{cases}$$

$$\begin{cases} x \rightarrow f(x) + \langle p, x \rangle_{H',H} \text{ admits a strict minimum over } D \text{ in } \bar{x}. & (2b) \end{cases}$$




 Boundedness of D
really essential ($f \equiv c$)

The Hilbertian Ekeland principle

Theorem – Ekeland-Lebourg [EL76] Let a closed bounded $D \subset H$ real Hilbert, and $f : D \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. $\forall \delta > 0, \exists \bar{x} \in D$ and $p \in H'$ s.t.

$$\begin{cases} |p|_{H'} < \delta, & (2a) \end{cases}$$

$$\begin{cases} x \rightarrow f(x) + \langle p, x \rangle_{H',H} \text{ admits a strict minimum over } D \text{ in } \bar{x}. & (2b) \end{cases}$$



👎 Boundedness of D
really essential ($f \equiv c$)


👍 Very nice perturbation

The Hilbertian Ekeland principle

Theorem – Ekeland-Lebourg [EL76] Let a closed bounded $D \subset H$ real Hilbert, and $f : D \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. $\forall \delta > 0, \exists \bar{x} \in D$ and $p \in H'$ s.t.

$$\begin{cases} |p|_{H'} < \delta, & (2a) \end{cases}$$

$$\begin{cases} x \rightarrow f(x) + \langle p, x \rangle_{H',H} \text{ admits a strict minimum over } D \text{ in } \bar{x}. & (2b) \end{cases}$$



👎 Boundedness of D
really essential ($f \equiv c$)

👍 Very nice perturbation

The proof is quite different.

The smooth Ekeland principle

Let (X, d) be a complete metric space.

gauge-type functions Any lower semicontinuous $\rho : X \times X \mapsto [0, \infty]$ satisfying $\rho(x, x) = 0$ for all $x \in X$, and $\forall \varepsilon > 0, \exists \eta > 0$ such that $\rho(x, y) \leq \eta$ implies $d(x, y) \leq \varepsilon$.

The smooth Ekeland principle

Let (X, d) be a complete metric space.

gauge-type functions Any lower semicontinuous $\rho : X \times X \mapsto [0, \infty]$ satisfying $\rho(x, x) = 0$ for all $x \in X$, and $\forall \varepsilon > 0, \exists \eta > 0$ such that $\rho(x, y) \leq \eta$ implies $d(x, y) \leq \varepsilon$.

Theorem – Borwein-Preiss [BP87] Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let ρ be gauge-type, $(\delta_i)_i \subset \mathbb{R}_*^+$, and $x_0 \in X$ such that $f(x_0) \leq \inf_X f + \varepsilon$.

The smooth Ekeland principle

Let (X, d) be a complete metric space.

gauge-type functions Any lower semicontinuous $\rho : X \times X \mapsto [0, \infty]$ satisfying $\rho(x, x) = 0$ for all $x \in X$, and $\forall \varepsilon > 0, \exists \eta > 0$ such that $\rho(x, y) \leq \eta$ implies $d(x, y) \leq \varepsilon$.

Theorem – Borwein-Preiss [BP87] Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let ρ be gauge-type, $(\delta_i)_i \subset \mathbb{R}_*^+$, and $x_0 \in X$ such that $f(x_0) \leq \inf_X f + \varepsilon$. Then there exist $y \in X$ and a sequence $(x_i)_{i=0}^\infty \subset X$ such that

$$\left\{ \begin{array}{ll} \rho(x_0, y) \leq \varepsilon/\delta_0 & \text{and} \quad \rho(x_i, y) \leq \varepsilon/(2^i \delta_0) \end{array} \right. \quad \begin{array}{l} (3a) \\ (3b) \\ (3c) \end{array}$$

The smooth Ekeland principle

Let (X, d) be a complete metric space.

gauge-type functions Any lower semicontinuous $\rho : X \times X \mapsto [0, \infty]$ satisfying $\rho(x, x) = 0$ for all $x \in X$, and $\forall \varepsilon > 0, \exists \eta > 0$ such that $\rho(x, y) \leq \eta$ implies $d(x, y) \leq \varepsilon$.

Theorem – Borwein-Preiss [BP87] Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let ρ be gauge-type, $(\delta_i)_i \subset \mathbb{R}_*^+$, and $x_0 \in X$ such that $f(x_0) \leq \inf_X f + \varepsilon$. Then there exist $y \in X$ and a sequence $(x_i)_{i=0}^\infty \subset X$ such that

$$\left\{ \begin{array}{ll} \rho(x_0, y) \leq \varepsilon/\delta_0 & \text{and } \rho(x_i, y) \leq \varepsilon/(2^i \delta_0) \\ f(y) + \sum_{i=0}^\infty \delta_i \rho(y, x_i) \leq f(x_0) \end{array} \right. \quad \begin{array}{l} (3a) \\ (3b) \\ (3c) \end{array}$$

The smooth Ekeland principle

Let (X, d) be a complete metric space.

gauge-type functions Any lower semicontinuous $\rho : X \times X \mapsto [0, \infty]$ satisfying $\rho(x, x) = 0$ for all $x \in X$, and $\forall \varepsilon > 0, \exists \eta > 0$ such that $\rho(x, y) \leq \eta$ implies $d(x, y) \leq \varepsilon$.

Theorem – Borwein-Preiss [BP87] Let $f : X \mapsto \mathbb{R} \cup \{\infty\}$ be proper, lsc and lower bounded. Let ρ be gauge-type, $(\delta_i)_i \subset \mathbb{R}_*^+$, and $x_0 \in X$ such that $f(x_0) \leq \inf_X f + \varepsilon$. Then there exist $y \in X$ and a sequence $(x_i)_{i=0}^\infty \subset X$ such that

$$\left\{ \begin{array}{l} \rho(x_0, y) \leq \varepsilon/\delta_0 \quad \text{and} \quad \rho(x_i, y) \leq \varepsilon/(2^i \delta_0) \end{array} \right. \quad (3a)$$

$$\left\{ \begin{array}{l} f(y) + \sum_{i=0}^\infty \delta_i \rho(y, x_i) \leq f(x_0) \end{array} \right. \quad (3b)$$

$$\left\{ \begin{array}{l} f(x) + \sum_{i=0}^\infty \delta_i \rho(x, x_i) > f(y) + \sum_{i=0}^\infty \delta_i \rho(y, x_i) \quad \forall x \in X \setminus \{y\}. \end{array} \right. \quad (3c)$$

Illustration of Borwein-Preiss




Figure: Iterative construction with $f(x) = (1 + |x|)^{-1}$, $\delta_i = 0.01/(1 + i)^2$, $\rho(x, y) = |x - y|^2$.

Illustration of Borwein-Preiss




Figure: Iterative construction with $f(x) = (1 + |x|)^{-1}$, $\delta_i = 0.01/(1 + i)^2$, $\rho(x, y) = |x - y|^2$.

Illustration of Borwein-Preiss




Figure: Iterative construction with $f(x) = (1 + |x|)^{-1}$, $\delta_i = 0.01/(1 + i)^2$, $\rho(x, y) = |x - y|^2$.

Table of Contents

The original principle

History

Statement

The proof

What since

History

Hilbert version: Ekeland-Lebourg

Smooth version: Borwein-Preiss

An application

Viscosity (I'm sure you missed it)

The Wasserstein context

- Space of measures μ with finite *second moment* $\int_{x \in E} |x|^2 d\mu(x)$.

Viscosity (I'm sure you missed it)

The Wasserstein context

- Space of measures μ with finite *second moment* $\int_{x \in E} |x|^2 d\mu(x)$.
- Narrow topology: duality with $\mathcal{C}_b(E)$.

Viscosity (I'm sure you missed it)

The Wasserstein context

- Space of measures μ with finite *second moment* $\int_{x \in E} |x|^2 d\mu(x)$.
- Narrow topology: duality with $\mathcal{C}_b(E)$.
- Wasserstein topology: duality with $\mathcal{C}_2(E)$.

Viscosity (I'm sure you missed it)

The Wasserstein context

- Space of measures μ with finite *second moment* $\int_{x \in E} |x|^2 d\mu(x)$.
- Narrow topology: duality with $\mathcal{C}_b(E)$.
- Wasserstein topology: duality with $\mathcal{C}_2(E)$.

The problem

- Aim: minimize a (coercive, lsc, proper) function in the Wasserstein space.

Viscosity (I'm sure you missed it)

The Wasserstein context

- Space of measures μ with finite *second moment* $\int_{x \in E} |x|^2 d\mu(x)$.
- Narrow topology: duality with $\mathcal{C}_b(E)$.
- Wasserstein topology: duality with $\mathcal{C}_2(E)$.

The problem

- Aim: minimize a (coercive, lsc, proper) function in the Wasserstein space.
- Bounded sets for the Wasserstein distance are narrowly compact, but not compact in the Wasserstein topology.

Viscosity (I'm sure you missed it)

The Wasserstein context

- Space of measures μ with finite *second moment* $\int_{x \in E} |x|^2 d\mu(x)$.
- Narrow topology: duality with $\mathcal{C}_b(E)$.
- Wasserstein topology: duality with $\mathcal{C}_2(E)$.

The problem

- Aim: minimize a (coercive, lsc, proper) function in the Wasserstein space.
- Bounded sets for the Wasserstein distance are narrowly compact, but not compact in the Wasserstein topology.

A solution by Marigonda & Quincampoix [MQ18]

- Use Ekeland to obtain δ -minimizers that are exact strict minima of perturbed functions.

Viscosity (I'm sure you missed it)

The Wasserstein context

- Space of measures μ with finite *second moment* $\int_{x \in E} |x|^2 d\mu(x)$.
- Narrow topology: duality with $\mathcal{C}_b(E)$.
- Wasserstein topology: duality with $\mathcal{C}_2(E)$.

The problem

- Aim: minimize a (coercive, lsc, proper) function in the Wasserstein space.
- Bounded sets for the Wasserstein distance are narrowly compact, but not compact in the Wasserstein topology.

A solution by Marigonda & Quincampoix [MQ18]

- Use Ekeland to obtain δ -minimizers that are exact strict minima of perturbed functions.
- What with Borwein & Preiss?

Thank you!

[AE84] Jean Pierre Aubin and I. Ekeland.

Applied Nonlinear Analysis.

Pure and Applied Mathematics. Wiley, New York, 1984.

[BP87] J. M. Borwein and D. Preiss.

A smooth variational principle with applications to subdifferentiability and to differentiability of convex functions.

Transactions of the American Mathematical Society, 303(2):517–527, 1987.

[BZ05] Jonathan M. Borwein and Qiji J. Zhu.

Techniques of Variational Analysis.

CMS Books in Mathematics. Springer-Verlag, New York, 2005.

[DGZ93] R. Deville, G. Godefroy, and V. Zizler.

A Smooth Variational Principle with Applications to Hamilton-Jacobi Equations in Infinite Dimensions.

Journal of Functional Analysis, 111(1):197–212, January 1993.

[Eke74] I. Ekeland.

On the variational principle.

Journal of Mathematical Analysis and Applications, 47(2):324–353, August 1974.

[EL76] Ivar Ekeland and Gérard Lebourg.

Generic Fréchet-differentiability and perturbed optimization problems in Banach spaces.

Transactions of the American Mathematical Society, 224(2):193–216, 1976.

[MQ18] Antonio Marigonda and Marc Quincampoix.

Mayer control problem with probabilistic uncertainty on initial positions.

Journal of Differential Equations, 264(5):3212–3252, March 2018.