From the Department of Public Health Sciences Karolinska Institutet, Stockholm, Sweden

NOVEL METHODS FOR DOSE-RESPONSE META-ANALYSIS

Alessio Crippa



Stockholm 2018

All published papers reproduced with permission Published by Karolinska Institutet Printed by E-Print AB 2018

Edited in R using knitr ©Alessio Crippa, 2018 ISBN <include number>

NOVEL METHODS FOR DOSE-RESPONSE META-ANALYSIS

THESIS FOR DOCTORAL DEGREE (Ph.D.)

By

Alessio Crippa

Principal supervisor:

Associate Professor Nicola Orsini

Karolinska Institutet

Department of Public Health Sciences

Co-supervisor:

Professor Alicja Wolk Karolinska Institutet

Institute of Environmental Medicine

Professor Matteo Bottai Karolinska Institutet

Institute of Environmental Medicine

Professor Donna Spiegelman

Harvard T.H. Chan School of Public Health

Department of Epidemiology

Opponent:

Professor Christopher H. Schmid

Brown University

Center for Evidence Based Medicine

Examination board:

Associate Professor Nele Brusselaers

Karolinska Institutet

Department of Microbiology, Tumor and Cell Biology

Associate Professor Antonio Gasparrini

London School of Hygiene and Tropical Medicine

Department of Social & Environmental Health Research

Professor Paul Lambert University of Leicester

Department of Health Sciences



Abstract

My abstract:

In Paper I,

In Paper II,

In Paper III,

In Paper IV,

In Paper V,

In conclusion,

List of publications

- I. Alessio Crippa, and Nicola Orsini
 Multivariate dose-response meta-analysis: the dosresmeta R Package
 Journal of Statistical Software, Code Snippets 2016; 72(1), 1–15
- II. Andrea Discacciati, Alessio Crippa, and Nicola Orsini
 Goodness of fit tools for dose–response meta-analysis of binary outcomes
 Research Synthesis Methods 2015
- III. Alessio Crippa, Polyna Khudyakov, Molin Wang, Nicola Orsini, and Donna Spiegelman A new measure of between-studies heterogeneity in meta-analysis *Statistics in medicine* 2016; 35(21), 3661–75
- IV. Alessio Crippa, Ilias Thomas, and Nicola OrsiniA pointwise approach to dose-response meta-analysis of aggregated dataManuscript 2018
- V. Alessio Crippa, Andrea Discacciati, Matteo Bottai, Alicja Wolk, and Nicola Orsini One-stage dose–response meta-analysis for aggregated data Manuscript 2018

The articles will be referred to in the text by their Roman numerals, and are reproduced in full at the end of the thesis.

Related publications

 Alessio Crippa, Susanna C. Larsson, Andrea Discacciati, Alicja Wolk, and Nicola Orsini Red and processed meat consumption and risk of bladder cancer: a dose–response meta-analysis of epidemiological studies

European journal of nutrition 2016, 1-13

- Andrea D. Smith, Alessio Crippa, James Woodcock, and Søren Brage
 Physical activity and incident type 2 diabetes mellitus: a systematic review and dose–response meta-analysis of prospective cohort studies
 Diabetologia 2016, 1–19
- Marco Vinceti, Tommaso Filippini, Alessio Crippa, Agnès de Sesmaisons, Lauren A. Wise, and Nicola Orsini

Meta-Analysis of Potassium Intake and the Risk of Stroke Journal of the American Heart Association 2016, 5(10), e004210

- Alessio Crippa, and Nicola Orsini
 Dose–response meta-analysis of differences in means
 BMC medical research methodology 2016, 16(1), 91
- Emir Veledar, Alessio Crippa, Chukwuemeka U. Osondu, Adnan Younus, and Khurram Nasir

Letter to Editor: Ideal cardiovascular health metrics and risk of cardiovascular disease or mortality

International journal of cardiology 2016, 222, 737

- Alessio Crippa, Andrea Discacciati, Nicola Orsini, and Viktor Oskarsson
 Letter: coffee consumption and gallstone disease—a cautionary note on the assignment of exposure values in dose–response meta-analyses
 Alimentary Pharmacology & Therapeutics 2016, 43(1), 166-167
- Susanna C. Larsson, Alessio Crippa, Nicola Orsini, Alicja Wolk, and Karl Michaëlsson
 Milk consumption and mortality from all causes, cardiovascular disease, and cancer: a systematic review and meta-analysis

Nutrients 2016, 7(9), 7749-7763

- Daniela Di Giuseppe, Alessio Crippa, Nicola Orsini, and Alicja Wolk
 Fish consumption and risk of rheumatoid arthritis: a dose-response meta-analysis
 Arthritis research & therapy 2014, 16(5), 446
- Alessio Crippa, Andrea Discacciati, Susanna C. Larsson, Alicja Wolk, and Nicola Orsini
 Coffee consumption and mortality from all causes, cardiovascular disease, and cancer: a dose–response meta-analysis

American journal of epidemiology 2014, 180(8), 763-775

Contents

1	Introduction	1
2	Background	3
3	Aims of the thesis	4
4	Materials and methods	5
5	Results	6
6	Discussion	7
7	Conclusions	8
8	Future research	9
Α	Supplementary figures	10
В	Supplementary tables	11
Re	References	
Αc	Acknowledgements	

List of abbreviations

AIC Akaike Information Criterion

CI Confidence Interval df Degrees of Freedom

GLS Generalized Least Squares

GRSS Generalized Residual Sum of SquaresGTSS Generalized Total Sum of SquaresFP2 Second-degree Fractional Polynomials

HRR Hazard Rate Ratio
IR Incidence Rate

IRR Incidence Rate Ratio logRR log-Relative Risk MR Mortality Rate

MRR Mortality Rate Ratio RCS Restricted Cubic Splines R^2 Coefficient of Determination

RR Relative Risk

WLS Weighted Least Squares

Introduction

A single experiment or study can hardly provide a definitive answer to a specific scientific question. Science is oftentimes referred to as a cumulative process where results from many studies, aiming to address the common question of interest, contribute to create and update the scientific evidence. In the cumulative paradigm, meta-analysis is the statistical methodology to combine and compare the current evidence in the field. This process lies at the heart of evidence-based medicine, and plays a major role in informing policy and practice.

Many epidemiological studies assess whether the occurrance of a health outcome (e.g. mortality, incidence of a diseases) varies according to a quantitative exposure (e.g. amount of physical activity, alcohol intake). The quantitative exposure is frequently divided in intervals and the results are expressed in a tabular format as relative risks for different exposure categories. A high vs. low meta-analysis contrasts the relative risks for the highest exposure versus the lower ones. This approach, however, discards the results for intermediate categories and thus provides only a limited picture. The information of the quantitative exposure is also lost, and the estimates being combined may be associated to different exposure values.

A dose–response meta-analysis, instead, has the potential to be more informative and powerful since it uses the whole available information to estimate the dose–response association. Because the estimates depends on the same reference category, it is not possible to regress the relative risks on the assigned dose using oridinaly least square. Greenland and Longnecker described in their seminal paper in 1992 how to reconstruct the correlation within set of relative risks and incorporate it in the dose–response analysis using generalized least square regression. Since then, the number of published dose–response meta-analysis has rapidly increased in many fields of application including oncology, public health, environmental sciences, nutrition, endocrinology, and internal medicine. Additional papers refined selected aspects of the proposed methodology, mainly focusing on the implementation of flexible strategies to model the dose-response curve and incorporating the advances of multivariate meta-analysis. However, there were still several relevant questions that needed to be addressed, including how to assess the goodness-of-fit, how to quantify the impact of heterogeneity, how to deal with differences in the exposure range across studies, and how to estimate complex model without excluding relevant studies.

2 1. Introduction

This thesis aims to address these open questions by developing new strategies and ad-hoc measures. The proposed methodologies are demonstrated reanalyzing published meta-analyses and implemented in user friendly packages in the free and open source R language to bridge the gap between theory and application.

Background

Write my background with subsections.

Here an example of a figure (Figure 2.1).




Figure 2.1

Aims of the thesis

The overall aims of this thesis were to develop and implement new methods for dose–response in meta-analysis, in order to deal with the methodological aspects that have not yet been addressed.

More specifically, the aims were:

- To describe the implementation of the main aspects of a dose–response meta-analysis and the usage of the dosresmeta R package.
- To present and discuss relevant measures and graphical tools to assess the goodness-of-fit in dose–response meta-analysis of binary outcome.
- To develop a new measure of between-study heterogeneity in the broader context of metaanalysis, and assess its statistical properties as compared to other available measures.
- To explore possible advantages of a point-wise approach, especially, in case of dose–response meta-analysis where the exposure range varies substantially across the studies.
- To formalize and present an alternative one-stage random-effects method for doseresponse meta-analysis of aggregated data, formulating the meta-analytic model in terms of a general linear mixed-effect model.

Materials and methods

Write materials and methods with subsections as in the background section

Results

Write the results with subsections as in the background section

Discussion

Write the discussion with subsections as in the background section

Conclusions

Write summary of conclusions.

More specifically we conclude the following:

- <>
- <>
- <>
- <>

Future research

Based on the conclusions presented in this thesis, future research includes:

- <>
- <>
- <>

Appendix A Supplementary figures

Figures.

Appendix B Supplementary tables

Tables.

References

- Crippa A, Discacciati A, Bottai M, Spiegelman D, Orsini N (2018a). "One-stage dose–response meta-analysis for aggregated data." *Manuscript*.
- Crippa A, Khudyakov P, Wang M, Orsini N, Spiegelman D (2016). "A new measure of between-studies heterogeneity in meta-analysis." *Statistics in medicine*, **35**(21), 3661–3675.
- Crippa A, Orsini N (2016). "Multivariate dose-response meta-analysis: The dosresmeta R Package." *Journal of statistical software, Code Snippets*, **72**(1), 1–15. doi:10.18637/jss.v072. c01.
- Crippa A, Thomas I, Orsini N (2018b). "A pointwise approach to dose-response meta-analysis of aggregated data." *Submitted*.
- Discacciati A, Crippa A, Orsini N (2015). "Goodness of fit tools for dose–response meta-analysis of binary outcomes." *Research synthesis methods*.

Acknowledgements

There are many people that I would like to thank for their contributions to this thesis, and for their support and encouragement during these years.

Nicola Orsini, my main supervisor for the second half of my doctoral education.

This work was supported by **Karolinska Institutet**'s funding for doctoral students (KID-funding).