Correlated residuals in a three-trait QTL SEM: identification, a replication of Mi et al. (2010), and a bow-free estimator

27 October 2026

Abstract

Mi, Eskridge, Wang, Baenziger, Campbell, Gill, Dweikat, and Bovaird asked, in the discussion of their 2010 QTL structural-equation paper, whether trait residuals can be correlated after environment and block effects have been removed. On the three-trait pattern printed in their equations (7)–(8), a diagonal residual matrix is globally identified, each single residual covariance on that pattern collides, and each of the three bow-free patterns is a left inverse of the covariance. Those statements are Lean proofs. This paper then runs three seeded simulations. The replication of their RIL experiment recovers the ordering of their multi-trait powers and the direct QTL coefficients, and it does not reproduce their single-trait power for the third QTL on the first trait. That discrepancy sits next to an arithmetic disagreement between equation (7) and several cells of their Table 2. A bow-free Monte Carlo shows that the diagonal plug-in invents a direct edge and biases two QTL coefficients, while `recoverBow` covers the path and QTL coefficients at the delta-method rate. A third run replaces the path coefficients by a fit to three phenotypic correlations from Campbell et al. (2003). It is a realism check on raw correlations, not a replication and not a fit to the wheat sample.

Question, source, and answer

Question	Can the trait residuals in the Mi et al. (2010) QTL SEM be correlated after environment and block effects are removed? Printed page 18 (PDF page 20).
Source	Mi et al., SAGMB 9 (2010), article 38. The request is not a numbered conjecture.
Answer on the printed pattern	On equation (7), diagonal residuals are a left inverse of the covariance. Each single residual covariance collides. Each bow-free missing-edge pattern is a left inverse, and the plug-in is almost surely consistent. Theorems 246, 114, 187, and 270.
What the simulations add	A seeded replication, the bias and size distortion of the diagonal fit, and coverage of `recoverBow`. These are sampled experiments, not Lean proofs.
Still open	A fit to the wheat measurements, the general mixed-graph characterization, and a Lean central limit theorem.

The manuscript date 27 October 2026 is an internal date. It is not an external publication date. Elapsed times below are measured to that date. A year-only source is reported as a range.

Source	Earliest date used	Inspected version	Elapsed to 27 Oct 2026
Brito and Pearl, bow-free identification	2002 (year)	SEM 9, 459–474	23–24 years
Campbell et al., chromosome 3A QTLs	2003 (year)	Crop Sci. 43, 1493–1505	22–23 years
Li et al., structural QTL model	21 July 2006	PLoS Genet. 2, e114	20 years, 3 months, 6 days
Mi et al., the source question	2010 (year)	SAGMB 9, article 38	15 years 10 months to 16 years 10 months
Valente et al., recursive search	June 2010 (month)	Genetics 185, 633–644	16 years, 4 months (month precision)
Drton, Foygel, and Sullivant	arXiv 4 Mar 2010; online 13 May 2011	Ann. Statist. 39, 865–886	15 years, 5 months, 14 days from the journal date
Foygel, Draisma, and Drton, half-trek	arXiv 27 July 2011; online 3 Oct 2012	Ann. Statist. 40, 1682–1713	14 years, 24 days from the journal date
Hidalgo-Contreras et al.	28 September 2021	Agronomy 11, 1953	5 years, 29 days
This manuscript	27 October 2026	internal manuscript date	—

1 The question, as they pose it

The source is Mi et al., Statistical Applications in Genetics and Molecular Biology 9 (2010), article 38. On printed page 18 (PDF page 20) they write:

> Here, we assumed that the residual errors of the traits were independent which we believe is a reasonable assumption since we removed the effects of environment by using, for the y values in equation (2), the residuals from a main effects linear model with environment and block within environment as the main effects. However, even after removing environment main effects, the traits may still be correlated and the trait error terms can be modeled as being correlated with each other.

(taken from a cited source) The parent request is that extension. It is not a numbered conjecture. Mi's 2009 dissertation (ProQuest AAI3350451) is the obvious place a fuller design might have been written. It is restricted to a ProQuest preview. It was not consulted.

2 The restriction

Equation (7) is the only fully printed structural pattern. With the sign convention of that display, $y = -My + Gx + \varepsilon$, the matrices on PDF page 10 are

$$M = \begin{pmatrix} 0 & 1/2 & 1/4 \\ 0 & 0 & 1/2 \\ 0 & 0 & 0 \end{pmatrix}, \qquad G = \begin{pmatrix} 1/8 & 1/4 & 1/2 \\ 1/2 & 1/2 & 1/2 \\ 1/4 & 1/8 & 1/4 \end{pmatrix},$$

and equation (8) is $\Psi = \operatorname{diag}(1.6, 1.8, 2.5)$. *(taken from a cited source)* In the regression form $y = By + Gx + \varepsilon$ one has $B = -M$. *(our assumption, read off the printed minus)* The wheat Figure 5 graph has four traits and is not this matrix. The restriction is covariance identification for this three-trait recursive pattern: diagonal Ψ, one extra residual covariance, and the three ways to

drop one edge and free the residual pair that edge had used. Singular Ψ is outside the claim. The general mixed-graph characterization is not this restriction.

3 What we proved

The following are Lean proofs on the positive-definite domain. The axioms are `propext`, `Classical.choice`, and `Quot.sound`. There is no `sorry`. *(our own Lean proof)*

- `recover_sigma`: diagonal residuals and positive-definite Φ are a left inverse of the joint covariance on the equation-(7) support.
- `bow12_same_law` and its two siblings: each one-edge residual extension collides, for every exogenous Γ.
- `recover_bow`, `recover_bow01`, `recover_bow23`: each bow-free missing-edge pattern is a left inverse.
- `unitUpper_factor_unique` and `sigmaRec_injective`: in every trait and covariate dimension, a recursive pattern with positive diagonal residuals has a unique unitriangular factorization, and the joint blocks determine (B, Ψ, Γ, Φ).
- `plug_in_consistent_ae` and the three bow-free analogues `plug_in_bow_consistent_ae`, `plug_in_bow01_consistent_ae`, `plug_in_bow23_consistent_ae`: pairwise i.i.d. samples with integrable coordinate products make those plug-ins almost surely consistent.

The Gaussian delta-method formula for the diagonal plug-in is a written derivative together with Isserlis' covariance, checked numerically in `delta_method.py`. *(our own written proof, and our own simulation)* It is not a Lean central limit theorem, and it is not listed among the proved results.

Statement	Lean declaration	File and lines
Diagonal left inverse	`recover_sigma`	`Diagonal.lean`, 246–279
One-edge collision	`bow12_same_law` and two siblings	`Collision.lean`, 114–160
Bow-free left inverse	`recover_bow`	`BowFree.lean`, 187–213
Other missing edges	`recover_bow01`, `recover_bow23`	`BowMissing.lean`, 148 and 353
Triangular factor	`unitUpper_factor_unique`	`Triangular.lean`, 68–135
Recursive identification	`sigmaRec_injective`	`Triangular.lean`, 298–317
Almost-sure consistency	`plug_in_consistent_ae` and `plug_in_bow_consistent_ae`	`Law.lean`, 174–199; `BowConsistency.lean`, 270–304

The links are pinned to commit `05d69ea141adf3d5ee936d3b48061bde241250dc`, which contains these Lean files. The simulation script and the JSON outputs are commit `5cd3c9f6d2f6e38f25e32146d74d55f3778436e1`. Appendix A prints the central proof bodies.

Brito and Pearl (Structural Equation Modeling, 2002) prove that a recursive linear model is identifiable when no residual covariance joins a variable to its direct effect. Our three bow-free left inverses are the three-trait case of that pattern, proved from the printed matrices rather than cited as a substitute for the algebra. Drton, Foygel, and Sullivant (Annals of Statistics, 2011) give a global criterion for mixed graphs; the half-trek criterion of Foygel, Draisma, and Drton (Annals of Statistics, 2012) is a generic sufficient condition. Neither is re-proved here, and the 2011 characterization stays a note with no inventory id. Li et al. (PLoS Genetics, 2006) and Valente, Rosa, de los Campos, Gianola, and Silva (Genetics, 2010) are earlier structural and recursive

treatments of multiple traits. Hidalgo-Contreras, Salinas-Ruiz, Eskridge, and Baenziger (Agronomy, 2021) fit a later Nebraska wheat SEM; their lines are a 2013 F6 panel, not the 1999–2001 RICLs, and none of their estimates enter the simulations.

4 Replication design

The printed design, PDF pages 9–10, is 100 replicates of 150 RIL lines, one chromosome of 100 cM, 11 evenly spaced markers, additive QTLs at 22, 42, and 78 cM, phenotypes from equations (7)–(8), and a test every 1 cM. QTL genotypes are coded $+1$ and -1. Cofactors are all markers except the flanking markers. The printed critical values are 14.86 and 10.60. *(taken from a cited source)* Those two numbers are the chi-square 0.005 quantiles on 4 and 2 degrees of freedom (14.860 and 10.597). The same paragraph also says the likelihood ratio has p degrees of freedom for p traits. We compare with the printed numbers. *(our own calculation of the printed thresholds)*

The paper does not state the following, so each is our assumption. *(our assumption)*

1. The map function is Haldane, with no interference.
2. An RIL is an F1 selfed for 12 generations. A leftover heterozygote is coded ± 1 with equal probability. There is no segregation distortion.
3. The eleven markers are at $0, 10, \ldots, 100$ cM.
4. The analysis uses the centered covariance with divisor n, which is the fitting function (6). We did not run SAS PROC TCALIS.
5. The flanking expectation is the conditional mean of the QTL genotype under that same 12-generation law. The 10 cM window mentioned later in the paper is for the wheat scan, not for this simulation, and was not used.
6. A named QTL is detected when the maximum likelihood ratio in $[0, 32]$, $(32, 60]$, or $(60, 100]$ exceeds the threshold. Effects are read at that argmax.

Seeds are `20101027` for this study. The outputs are `study1.json`, `study1_power.csv`, and `study1_effects.csv` under `simulation/out`.

For a complete recursive order, the Gaussian likelihood of the diagonal structural model equals the likelihood of the multivariate regression on the same exogenous regressors: the sum of the successive residual log variances is the log determinant of the residual covariance. The two joint likelihood ratios are therefore the same number. In the run, the maximum absolute difference of the two mean profiles is 7×10^{-14}. *(our own written proof, confirmed by our own simulation)* Mi et al. describe the two profiles as very close (PDF page 14). Their Table 1 powers still differ by 7, 8, and 2 replicates out of 100.

5 Replication results

Table 1 compares detection percentages. Our multi-trait powers keep the printed order, third QTL above the first two, and they sit 5 to 16 points below Table 1. The single-trait column for the first trait at the first QTL matches Table 1 exactly (6%). The single-trait column for the first trait at the third QTL does not: Table 1 prints 97% and the replication gives 13%.

Table 1: Detection percentages. "Mi" is Table 1 on PDF page 12. "Ours" is 100 replicates, seed `20101027`. SEM and LS coincide in our run.

QTL	SEM Mi	SEM ours	LS Mi	ST Y_1 Mi	ST Y_1 ours	ST overall ours
1	64	59	57	6	6	24
2	77	61	69	3	0	35
3	96	86	94	97	13	34

The algebraic total effects under $B = -M$ are, by columns of G,

$$(I + M)^{-1}G = \begin{pmatrix} -1/8 & 0 & 1/4 \\ 3/8 & 7/16 & 3/8 \\ 1/4 & 1/8 & 1/4 \end{pmatrix}.$$

(our own calculation) Table 2's parameter block matches this matrix on Y_2 and Y_3 for the first and third QTLs, and it does not match it on Y_1. The printed total effect of the third QTL on Y_1 is 0.75, against 0.25 from the display. A regression of Y_1 on the true third QTL genotype, with no cofactors, has power about 0.10 at the threshold 10.60 when the total effect is 0.25 (200 further replicates, same generator). The printed 97% is the power of a substantially larger effect. We did not find a sign convention that matches both the minus in equation (7) and every parameter cell in Table 2. The simulation follows the equation.

Direct-effect means at the SEM argmax are close to G where the signal is large. For the third QTL they are 0.460 (0.175), 0.476 (0.179), and 0.242 (0.190), against the printed SEM estimates 0.479 (0.176), 0.527 (0.166), and 0.257 (0.205) and against $G = (1/2, 1/2, 1/4)$. *(our own simulation, set next to a cited source)* Position means are 22.65 (7.28), 41.26 (5.25), and 79.85 (3.76), against the printed 21.57 (2.83), 41.95 (2.58), and 78.03 (2.16). The biases are small. Our standard deviations are larger. The mean profile exceeds 14.86 only at 40 cM and on 75–82 cM. Figure 2 is described as one simulated population, with excursions on 16–28, 38–48, and 70–88 cM. An average profile is a different object, and ours is lower.

6 Correlated residuals

The bow drops the edge into trait 0 from trait 2, sets that structural coefficient to zero, and frees the residual covariance $c = \rho\sqrt{1.6 \cdot 2.5}$. The other equation-(7) coefficients stay. The exogenous variables are the three QTL genotypes on the RIL map, not a marker scan. *(our assumption, matching `recoverBow`)* The sample is the centered second-moment matrix, $n = 150$, 400 replicates, seed `20101028`.

At the population covariance the bow plug-in recovers the parameter to machine precision. The diagonal plug-in does not. Its probability limit for the missing edge is 0.8ρ: zero at $\rho = 0$, 0.20 at $\rho = 0.25$, and 0.48 at $\rho = 0.6$. The QTL coefficient $G_{11} = 1/8$ has probability limit 0.075 at $\rho = 0.25$ and 0.005 at $\rho = 0.6$. The coefficient $G_{13} = 1/2$ moves to 0.45 and 0.38. *(our own simulation)* A Wald test that the missing edge is zero, using the diagonal delta-method standard error, rejects at rate 0.058 when $\rho = 0$ and at rate 0.728, 0.877, and 1 at $\rho = 0.25$, 0.30, and 0.60. The joint nine-degree Wald test of all QTL coefficients is only mildly inflated at $\rho = 0.25$ (rejection rate 0.065) and reaches 0.297 at $\rho = 0.6$. The damage shows up first as bias in two coefficients and as a spurious edge, not as an immediate explosion of the joint QTL test.

`recoverBow` is centered on the QTL and path coefficients. At $\rho = 0.25$ the Monte Carlo means of b_{12}, b_{23}, G_{11}, and G_{13} are -0.507, -0.498, 0.127, and 0.509. Coverage of the nominal 95%

delta-method intervals, with the Gaussian Isserlis covariance plugged into the sample, is between 0.915 and 0.958 across the reported coefficients and the six correlations. The genotypes are ± 1, so Isserlis is an approximation. *(our assumption inside the standard error)* The coverage near 95% says the approximation is tolerable at $n = 150$ for these coefficients. Variance-component means sit a little low (1.54 against 1.6 at $\rho = 0$) because the likelihood uses divisor n; the probability limit is still the population value.

7 Calibration to three phenotypic correlations

Campbell et al., Crop Science 43 (2003), Table 2, print phenotypic correlations for 95 recombinant inbred chromosome lines pooled across seven environments. There are no standard deviations and no environment or block adjustment. *(taken from a cited source, with the limits just stated)* The text layer of the copy we could read dropped minus signs. The paragraph above the table says that SPSM and kernels per square meter were negatively correlated with thousand-kernel weight. We therefore use $\mathrm{Corr}(\mathrm{TKWT}, \mathrm{SPSM}) = -0.44$. The cells $\mathrm{Corr}(\mathrm{TKWT}, \mathrm{KPS}) = 0.08$ and $\mathrm{Corr}(\mathrm{KPS}, \mathrm{SPSM}) = 0.59$ are the signs in that text layer; the prose does not restate them. *(our assumption for those two signs)* GYLD is the fourth trait in the application and is not in equation (7). The three-trait order is $Y_1 = \mathrm{TKWT}$, $Y_2 = \mathrm{KPS}$, $Y_3 = \mathrm{SPSM}$, which is the chain used in the indirect-effect arithmetic on PDF page 17 of Mi et al. *(our assumption)*

Treating the correlation matrix as a covariance, the diagonal recursive factorization is

$$B = \begin{pmatrix} 0 & 0.521 & -0.747 \\ 0 & 0 & 0.590 \\ 0 & 0 & 0 \end{pmatrix}, \qquad \Psi = \mathrm{diag}(0.629, 0.652, 1).$$

(our own calculation) The eigenvalues of the correlation are 0.224, 1.077, and 1.699, so this sign choice is positive definite. Naive Gaussian standard errors that pretend the matrix is a sample correlation of 95 i.i.d. lines are about 0.10 for the two coefficients in the first equation and 0.084 for 0.590. Campbell pooled seven environments, so those standard errors are not his sampling variances. The point fit does not depend on N.

The scan of Section 5 was rerun with this B and Ψ and with G copied from equation (7). Seed `20101029`, 100 replicates. Those QTL coefficients were not estimated from the correlations, and on this residual scale they are large: multi-trait power is 100%, 96%, and 100%. That is a consequence of the scale mixture, not evidence that the 2003 correlations produce Mi's powers. *(our own simulation, under an assumption we flag)* The bow check at $\rho = 0.25$ (seed `20101030`) again gives a diagonal probability limit 0.198 for the missing edge and a rejection rate 0.74, with bow coverage 0.935 for the residual covariance.

8 Limitations

The wheat fit is still blocked. It needs the 98 line identifiers, the ANOVA residuals of grain yield, thousand-kernel weight, kernels per spike, and spikes per square meter, the ± 1 QTL and cofactor codes, the divisor-n sample covariance, and the edge set as a matrix. Figure 5 draws arrows and does not print that matrix. Campbell's Table 2 does not substitute for any of those columns.

The Drton–Foygel–Sullivant criterion is not a theorem of this repository. The delta method is not a Lean central limit theorem. The replication does not reproduce Table 1's 97% or Table 2's indirect-effect signs, because those cells disagree with equations (7)–(8) under the printed minus.

Twelve generations of selfing, Haldane's map, the marker placement, and the detection windows are our assumptions. The 2021 Hidalgo-Contreras wheat SEM is a different experiment.

9 Reproducibility

From `formalization/correlated-residuals/simulation`,

`python3 studies.py --reps 100 --bow-reps 400 --studies 1,2,3`

writes `study1.json`, `study2.json`, and `study3.json`. The CSV tables are built from those files.

- Simulation commit: `5cd3c9f`.
- Full hash: `5cd3c9f6d2f6e38f25e32146d74d55f3778436e1`.
- Branch: `cursor/l-f102-correlated-residuals-4c01`.
- Lean 4.22.0 and Mathlib v4.22.0, built with `lake build`.
- Theorem links are pinned to `05d69ea141adf3d5ee936d3b48061bde241250dc`.
- This PDF is `pdflatex paper.tex`. The ledger is `SOURCES.md`.

A Central Lean proofs

The diagonal model reads the structural coefficients from the Schur complement $\Omega = \Sigma_{yy} - \Sigma_{yx}\Sigma_{xx}^{-1}\Sigma_{yx}^{\mathsf{T}}$, then recovers $\Gamma = (I - B)\Sigma_{yx}\Sigma_{xx}^{-1}$ and $\Phi = \Sigma_{xx}$. The bow-free model drops the edge into the first trait from the third and reads the residual covariance c from the same complement. The collision proofs exhibit, for an arbitrary exogenous coefficient, a second parameter point with the same joint blocks. Consistency is continuous mapping of those left inverses, composed with the strong law for the sample Gram matrix. The listings below are the theorem bodies. Their lemmas live in the linked files.

`recover_sigma`, `Diagonal.lean`, lines 246–279.

```
theorem recover_sigma (θ : Params) (h : OnDomain θ) : recover (sigma θ) = θ := by
  obtain ⟨hv1, hv2, hv3, hφ⟩ := h
  have _hpsi : (psi θ).PosDef := by
    rw [psi, posDef_diagonal_iff]
    intro i
    fin_cases i
    · simpa using hv1
    · simpa using hv2
    · simpa using hv3
  have hread : readDiag (omega θ) = (θ.b12, θ.b13, θ.b23, θ.v1, θ.v2, θ.v3) :=
    readDiag_omega θ hv2 hv3
  have hschur := schur_omega θ hφ
  simp only [recover, hschur, hread]
  have hB : (1 -
      ![0, θ.b12, θ.b13;
        0, 0, θ.b23;
        0, 0, 0]) = subB θ := by
    simpa [Bmat] using (subB_eq_one_sub θ).symm
  have hdet : IsUnit θ.phi.det := (isUnit_iff_isUnit_det θ.phi).1 hφ.isUnit
  apply Params.ext
```

```
  · rfl
  · rfl
  · rfl
  · rfl
  · rfl
  · rfl
  · simp only [sigma]
    rw [hB]
    calc
      subB θ * (Lmat θ * θ.gamma * θ.phi) * θ.phi⁻¹
          = (subB θ * Lmat θ) * θ.gamma * (θ.phi * θ.phi⁻¹) := by simp only [mul_assoc]
      _ = (1 : M3) * θ.gamma * (1 : M3) := by rw [subB_mul_L, mul_nonsing_inv _ hdet]
      _ = θ.gamma := by simp
  · rfl
```

recover_bow, BowFree.lean, lines 187–213.

```
theorem recover_bow (θ : Bow) (h : OnBow θ) : recoverBow θ.sigma = θ := by
  obtain ⟨hN, hφ⟩ := h
  have hv2 := Bow.v2_pos θ hN
  have hv3 := Bow.v3_pos θ hN
  have hread := readBow_omega θ hv2 hv3
  have hschur : θ.sigma.yy - θ.sigma.yx * θ.sigma.xx⁻¹ * θ.sigma.yxᵀ = θ.omega := by
    simpa [Bow.sigma, Bow.omega] using
      schur_noise (L := θ.L) (G := θ.gamma) (Φ := θ.phi) (N := θ.noise) hφ
  simp only [recoverBow, hschur, hread]
  have hB : (1 - !![0, θ.b12, 0; 0, 0, θ.b23; 0, 0, 0]) = θ.subB := by
    simpa [Bow.B] using (Bow.subB_eq θ).symm
  have hdet : IsUnit θ.phi.det := (isUnit_iff_isUnit_det θ.phi).1 hφ.isUnit
  apply Bow.ext
  · rfl
  · rfl
  · rfl
  · rfl
  · rfl
  · rfl
  · simp only [Bow.sigma]
    rw [hB]
    calc
      θ.subB * (θ.L * θ.gamma * θ.phi) * θ.phi⁻¹
          = (θ.subB * θ.L) * θ.gamma * (θ.phi * θ.phi⁻¹) := by simp only [mul_assoc]
      _ = (1 : M3) * θ.gamma * (1 : M3) := by rw [Bow.subB_mul_L, mul_nonsing_inv _ hdet]
      _ = θ.gamma := by simp
  · rfl
```

bow12_same_law in Collision.lean, lines 114–128. The siblings bow13_same_law and bow23_same_law repeat the argument on the other two pairs, lines 130–160.

```
theorem bow12_same_law (c : ℝ) (G Φ : M3) :
    bow12L c * (bow12G c * G) * Φ = G * Φ ∧
      bow12L c * ((bow12G c * G) * Φ * (bow12G c * G)ᵀ + bow12G c * (bow12G c)ᵀ) * (bow12L c)ᵀ =
        G * Φ * Gᵀ + 1 := by
  have hG : bow12L c * (bow12G c * G) = (1 : M3) * G := by
    rw [← mul_assoc, bow12_resolve c, one_mul]
  have hN : bow12L c * (bow12G c * (bow12G c)ᵀ) * (bow12L c)ᵀ = (1 : M3) :=
    resolved_noise (bow12_resolve c)
  have hbaseN : (1 : M3) * (1 : M3) * (1 : M3)ᵀ = (1 : M3) := by simp
  obtain ⟨hyx, hyy⟩ := exogenous_lift (L := bow12L c) (G := bow12G c * G)
```

```
    (L' := (1 : M3)) (G' := G) (N := bow12G c * (bow12G c)ᵀ) (N' := (1 : M3)) (Φ := Φ)
    hG (hN.trans hbaseN.symm)
  constructor
  · simpa using hyx
  · simpa using hyy
```

unitUpper_factor_unique, Triangular.lean, lines 68–135.

```
theorem unitUpper_factor_unique {n : ℕ}
    {L M : Matrix (Fin n) (Fin n) ℝ} {d e : Fin n → ℝ}
    (hL : UnitUpper L) (hM : UnitUpper M)
    (hd : ∀ i, 0 < d i) (he : ∀ i, 0 < e i)
    (hΩ : L * diagonal d * Lᵀ = M * diagonal e * Mᵀ) :
    L = M ∧ d = e := by
  induction n with
  | zero =>
    constructor
    · ext i j
      exact i.elim0
    · ext i
      exact i.elim0
  | succ n ih =>
    have hde_last : d (Fin.last n) = e (Fin.last n) := by
      have hLlast := gram_last (d := d) hL (Fin.last n)
      have hMlast := gram_last (d := e) hM (Fin.last n)
      rw [hL.2] at hLlast
      rw [hM.2] at hMlast
      have hentry := congr_fun (congr_fun hΩ (Fin.last n)) (Fin.last n)
      rw [hLlast, hMlast] at hentry
      simpa using hentry
    have hcol : ∀ i, L i (Fin.last n) = M i (Fin.last n) := by
      intro i
      have hLi := gram_last (d := d) hL i
      have hMi := gram_last (d := e) hM i
      have hentry := congr_fun (congr_fun hΩ i) (Fin.last n)
      rw [hLi, hMi, hde_last] at hentry
      exact mul_right_cancel₀ (he (Fin.last n)).ne' hentry
    have hlead :
        lead L * diagonal (fun k : Fin n => d k.castSucc) * (lead L)ᵀ =
          lead M * diagonal (fun k : Fin n => e k.castSucc) * (lead M)ᵀ := by
      ext i j
      have hLij := lead_gram (L := L) (d := d) i j
      have hMij := lead_gram (L := M) (d := e) i j
      have hentry := congr_fun (congr_fun hΩ i.castSucc) j.castSucc
      rw [hLij, hMij] at hentry
      have hr :
          L i.castSucc (Fin.last n) * d (Fin.last n) * L j.castSucc (Fin.last n) =
            M i.castSucc (Fin.last n) * e (Fin.last n) * M j.castSucc (Fin.last n) := by
        rw [hcol, hcol, hde_last]
      rw [hr] at hentry
      exact add_right_cancel hentry
    have hrec :=
      ih (lead_unitUpper hL) (lead_unitUpper hM) (fun i => hd i.castSucc)
        (fun i => he i.castSucc) hlead
    refine ⟨?_, ?_⟩
    · ext i j
      cases i using Fin.lastCases with
      | last =>
        cases j using Fin.lastCases with
```

```
        | last =>
          rw [hL.2, hM.2]
        | cast j =>
          rw [hL.1 _ _ j.castSucc_lt_last, hM.1 _ _ j.castSucc_lt_last]
      | cast i =>
        cases j using Fin.lastCases with
        | last =>
          exact hcol i.castSucc
        | cast j =>
          have hlead_ij := congr_fun (congr_fun hrec.1 i) j
          simpa [lead, submatrix_apply] using hlead_ij
    · ext i
      cases i using Fin.lastCases with
      | last => exact hde_last
      | cast i =>
        have := congr_fun hrec.2 i
        simpa using this
```

sigmaRec_injective, Triangular.lean, lines 298–317.

```
theorem sigmaRec_injective {n p : ℕ} {θ θ' : RecModel n p}
    (hθ : OnRec θ) (hθ' : OnRec θ') (h : sigmaRec θ = sigmaRec θ') : θ = θ' := by
  have hθ0 := hθ
  have hθ'0 := hθ'
  obtain ⟨hB, hd, hΦ⟩ := hθ
  obtain ⟨hC, he, hΦ'⟩ := hθ'
  have hgram :
      (1 - θ.B)⁻¹ * diagonal θ.d * ((1 - θ.B)⁻¹)ᵀ =
        (1 - θ'.B)⁻¹ * diagonal θ'.d * ((1 - θ'.B)⁻¹)ᵀ := by
    have hθs := schur_sigmaRec θ hΦ
    have hθs' := schur_sigmaRec θ' hΦ'
    rw [h] at hθs
    exact hθs.symm.trans hθs'
  obtain ⟨hBeq, hdeq⟩ := recursive_residual_unique hB hC hd he hgram
  have hγ : θ.gamma = θ'.gamma := by
    have hg := (recover_exogenous θ hθ0).2
    have hg' := (recover_exogenous θ' hθ'0).2
    rw [hg, hg', h, hBeq]
  have hφ : θ.phi = θ'.phi := by simpa [sigmaRec] using congrArg Joint.xx h
  exact RecModel.ext hBeq hdeq hγ hφ
```

plug_in_consistent_ae, Law.lean, lines 174–199.

```
theorem plug_in_consistent_ae {Ω : Type*} [MeasurableSpace Ω] {μ : Measure Ω}
    {Z : ℕ → Ω → Idx → ℝ} {θ : Params} (hθ : OnDomain θ)
    (hindep : Pairwise fun i j => IndepFun (Z i) (Z j) μ)
    (hident : ∀ i, IdentDistrib (Z i) (Z 0) μ μ)
    (hint : ∀ a b : Idx, Integrable (fun ω => Z 0 ω a * Z 0 ω b) μ)
    (hyy : ∀ i j : Fin 3,
      ∫ ω, Z 0 ω (.inl i) * Z 0 ω (.inl j) ∂μ = (sigma θ).yy i j)
    (hyx : ∀ i j : Fin 3,
      ∫ ω, Z 0 ω (.inl i) * Z 0 ω (.inr j) ∂μ = (sigma θ).yx i j)
    (hxx : ∀ i j : Fin 3,
      ∫ ω, Z 0 ω (.inr i) * Z 0 ω (.inr j) ∂μ = (sigma θ).xx i j) :
    ∀ᵐ ω ∂μ, Tendsto (fun n =>
        ((recover (sampleCov Z n ω)).b12, (recover (sampleCov Z n ω)).b13,
          (recover (sampleCov Z n ω)).b23, (recover (sampleCov Z n ω)).v1,
          (recover (sampleCov Z n ω)).v2, (recover (sampleCov Z n ω)).v3,
```

```
        (recover (sampleCov Z n ω)).gamma, (recover (sampleCov Z n ω)).phi))
    atTop
    (nhds (θ.b12, θ.b13, θ.b23, θ.v1, θ.v2, θ.v3, θ.gamma, θ.phi)) := by
  filter_upwards [strong_gram Z hindep hident hint] with ω hω
  have hyy' : Tendsto (fun n => (sampleCov Z n ω).yy) atTop (nhds (sigma θ).yy) := by
    simpa [sampleCov, blockYY_pop hyy] using tendsto_blockYY hω
  have hyx' : Tendsto (fun n => (sampleCov Z n ω).yx) atTop (nhds (sigma θ).yx) := by
    simpa [sampleCov, blockYX_pop hyx] using tendsto_blockYX hω
  have hxx' : Tendsto (fun n => (sampleCov Z n ω).xx) atTop (nhds (sigma θ).xx) := by
    simpa [sampleCov, blockXX_pop hxx] using tendsto_blockXX hω
  exact plug_in_consistent hθ hyy' hyx' hxx'
```

plug_in_bow_consistent_ae, BowConsistency.lean, lines 270–304.

```
theorem plug_in_bow_consistent_ae {Ω : Type*} [MeasurableSpace Ω] {μ : Measure Ω}
    {Z : ℕ → Ω → Idx → ℝ} {θ : Bow} (hθ : OnBow θ)
    (hindep : Pairwise fun i j => IndepFun (Z i) (Z j) μ)
    (hident : ∀ i, IdentDistrib (Z i) (Z 0) μ μ)
    (hint : ∀ a b : Idx, Integrable (fun ω => Z 0 ω a * Z 0 ω b) μ)
    (hyy : ∀ i j : Fin 3,
      ∫ ω, Z 0 ω (.inl i) * Z 0 ω (.inl j) ∂μ = θ.sigma.yy i j)
    (hyx : ∀ i j : Fin 3,
      ∫ ω, Z 0 ω (.inl i) * Z 0 ω (.inr j) ∂μ = θ.sigma.yx i j)
    (hxx : ∀ i j : Fin 3,
      ∫ ω, Z 0 ω (.inr i) * Z 0 ω (.inr j) ∂μ = θ.sigma.xx i j) :
    ∀ᵐ ω ∂μ, Tendsto (fun n =>
        ((recoverBow (sampleCov Z n ω)).b12, (recoverBow (sampleCov Z n ω)).b23,
          (recoverBow (sampleCov Z n ω)).v1, (recoverBow (sampleCov Z n ω)).v2,
          (recoverBow (sampleCov Z n ω)).v3, (recoverBow (sampleCov Z n ω)).c,
          (recoverBow (sampleCov Z n ω)).gamma, (recoverBow (sampleCov Z n ω)).phi))
      atTop
      (nhds (θ.b12, θ.b23, θ.v1, θ.v2, θ.v3, θ.c, θ.gamma, θ.phi)) := by
  filter_upwards [strong_gram Z hindep hident hint] with ω hω
  have hyy' : Tendsto (fun n => (sampleCov Z n ω).yy) atTop (nhds θ.sigma.yy) := by
    have hpop : blockYY (popGram μ (Z 0)) = θ.sigma.yy := by
      ext i j
      simpa [blockYY, popGram] using hyy i j
    simpa [sampleCov, hpop] using tendsto_blockYY hω
  have hyx' : Tendsto (fun n => (sampleCov Z n ω).yx) atTop (nhds θ.sigma.yx) := by
    have hpop : blockYX (popGram μ (Z 0)) = θ.sigma.yx := by
      ext i j
      simpa [blockYX, popGram] using hyx i j
    simpa [sampleCov, hpop] using tendsto_blockYX hω
  have hxx' : Tendsto (fun n => (sampleCov Z n ω).xx) atTop (nhds θ.sigma.xx) := by
    have hpop : blockXX (popGram μ (Z 0)) = θ.sigma.xx := by
      ext i j
      simpa [blockXX, popGram] using hxx i j
    simpa [sampleCov, hpop] using tendsto_blockXX hω
  exact plug_in_bow_consistent hθ hyy' hyx' hxx'
```

References

[1] C. Brito and J. Pearl. A new identification condition for recursive models with correlated errors. *Structural Equation Modeling* 9 (2002), 459–474.

[2] B. T. Campbell, P. S. Baenziger, K. S. Gill, K. M. Eskridge, H. Budak, M. Erayman, and Y. Yen. Identification of QTLs and environmental interactions associated with agronomic traits on chromosome 3A of wheat. *Crop Science* 43 (2003), 1493–1505.

[3] M. Drton, R. Foygel, and S. Sullivant. Global identifiability of linear structural equation models. *Annals of Statistics* 39 (2011), 865–886.

[4] R. Foygel, J. Draisma, and M. Drton. Half-trek criterion for generic identifiability of linear structural equation models. *Annals of Statistics* 40 (2012), 1682–1713.

[5] J. V. Hidalgo-Contreras, J. Salinas-Ruiz, K. M. Eskridge, and S. P. Baenziger. Incorporating molecular markers and causal structure among traits using a Smith–Hazel index and structural equation models. *Agronomy* 11 (2021), 1953.

[6] R. Li, S.-W. Tsaih, K. Shockley, I. M. Stylianou, J. Wergedal, B. Paigen, and G. A. Churchill. Structural model analysis of multiple quantitative traits. *PLoS Genetics* 2 (2006), e114.

[7] X. Mi, K. Eskridge, D. Wang, P. S. Baenziger, B. T. Campbell, K. S. Gill, I. Dweikat, and J. Bovaird. Regression-based multi-trait QTL mapping using a structural equation model. *Statistical Applications in Genetics and Molecular Biology* 9 (2010), article 38.

[8] B. D. Valente, G. J. M. Rosa, G. de los Campos, D. Gianola, and M. A. Silva. Searching for Recursive Causal Structures in Multivariate Quantitative Genetics Mixed Models. *Genetics* 185 (2010), 633–644.