SRINIVAS UNIVERSITY INSTITUTE OF ENGINEERING & TECHNOLOGY

(SUBJECT:ARTIFICIAL NEURAL NETWORK) (SUBJECT CODE:24SBT113)

A Individual Task on

"Models Equations of Activation and Synaptic Dynamics in Artificial Neural Networks"

Submitted in the partial fulfillment of the requirements for the fourth semester

BACHELOR OF TECHNOLOGY IN AIML

SubmittedBy,

ALEENA V.K (01SU24AI119)

UNDER THE GUIDANCE OF

Prof.MaheshKumarVB

DEPARTMENT OF ARTIFICIAL INTELLIGENCE & MACHINE LEARNING

SRINIVAS UNIVERSITY INSTITUTE OF ENGINEERING & TECHNOLOGY
MUKKA, MANGALURU-574146

2025-26

SRINIVAS UNIVERSITY

INSTITUTEOFENGINEERING&TECHNOLOGY

DEPARTMENT ARTIFICIAL INTELLIGENCE & MACHINE LEARNING

CERTIFICATE

This is to certify that ALEENA V.K (01SU24AI119) has satisfactorily completed the assessment (Individual-Task – Module 2) in **"ARTIFICIAL NEURAL NETWORK"** prescribed by the Srinivas University for the 4st semester B. Tech course during the year **2025-26**.

MARKS AWARDED

Staff In charge

Name: Prof. Mahesh Kumar V B

Assistant Professor,Department Of AIML

Introduction

Artificial Neural Networks (ANNs) are systems of interconnected processing units whose behaviour is governed by two fundamental mechanisms: **activation dynamics** and **synaptic dynamics**.

- **Activation dynamics** describe how neuron states(activations) evolve with time for a fixed input.
- **Synaptic dynamics** describe how connection weights change during learning.

Mathematically, both are represented using **first-order differential equations**, which model the rate of change of activation or weight with respect to time.

2. PassiveDecayModel

$$\frac{dx_i(t)}{dt} = -A_i x_i(t)$$

Symbols

- $x_i(t)$:activation(state)of neuron indexed by i at time t
- i: index identifying the neuron
- t:time variable
- $\frac{dx_i(t)}{dt}$: derivative of $x_i(t)$ with respect to time (rate of change)
- A_i:positive decay(leakage) constant for neuron i
- $-$:indicates reduction(decay) of activation

3.Modified Passive Decay Model

$$\frac{dx_i(t)}{dt} = -\frac{A_i}{C_i} x_i(t)$$

Symbols

- $x_i(t)$:activation of neuron i at time t
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change of activation

- A_i:decay constant
- C_i:membrane capacitance of neuron i
- $\frac{A_i}{C_i}$: effective decayrate
- $-$:decay effect

4.Non - Zero Resting Potential Model

$$\frac{dx_i(t)}{dt} = -A_i x_i(t) + P_i$$

Symbols

- $x_i(t)$:activation of neuron i at time t
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change of activation
- A_i:decay constant
- P_i:constant input (resting potential or bias)
- $+$: additive effect

5.External Input Activation Model

$$\frac{dx_i(t)}{dt} = -A_i x_i(t) + B_i I_i$$

Symbols

- $x_i(t)$:activation of neuron i at time t
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change of activation
- A_i:decay constant
- B_i: input gain(scaling factor)
- I_i:external input signal applied to neuron i
- $+$: additive contribution

6. Additive Autoassociative Model

$$\frac{dx_i(t)}{dt} = -A_i x_i(t) + B_i I_i + \sum_{j=1}^{N} W_{ij} f_j(x_j(t))$$

Symbols

- $x_i(t)$:activation of neuron i at time t
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change of activation
- A_i:decay constant
- B_i:input gain
- I_i:external input to neuron i
- $\sum_{j=1}^{N}$:summation over index j from 1 to N
- j:index of pre synaptic neuron
- N:total number of neurons
- W_{ij}:synaptic weight from neuron j to neuron i
- $f_j(\cdot)$:output function of neuron j
- $x_j(t)$:activation of neuron j at time t

7. Inhibitory Feedback Model

$$\frac{dx_i(t)}{dt} = -A_i x_i(t) - B_i I_i - \sum_{j=1}^{N} W_{ij} f_j(x_j(t))$$

Symbols

- $x_i(t)$:activation of neuron i at time t
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change
- A_i:decay constant
- B_i:scaling constant

- I_i:external input
- $\sum_{j=1}^{N}$: summation over neurons
- j:neuron index
- N:total neurons
- W_{ij}:weight from neuron j to neuron i
- $f_j(x_j(t))$: output of neuron j

8. Perkel's Model

$$\frac{dx_i(t)}{dt} = \frac{1}{R_i}x_i(t) + \sum_{j=1}^{N} \frac{1}{R_{ij}}\phi_j(x_j(t))$$

Symbols

- $x_i(t)$:activation of neuron i
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change
- R_i: resistance of neuron i
- $\frac{1}{R_i}$:conductance
- $\sum_{j=1}^{N}$: summation over neurons
- j:neuron index
- N:total neurons
- R_{ij}:resistance between neurons i and j
- $\phi_j(\cdot)$:output function
- $x_j(t)$: activation of neuron j

9. Hopfield Model

$$\frac{dx_i(t)}{dt} = -A_i x_i(t) + \sum_{j=1}^{N} W_{ij} f(x_j(t)) + I_i$$

Symbols

- $x_i(t)$:activation of neuron i
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change
- A_i:decay constant
- $\sum_{j=1}^{N}$: summation
- j:neuron index
- N:total neurons
- W_{ij}:symmetric connection weight
- $f(\cdot)$:bounded activation function
- $x_j(t)$: activation of neuron j
- I_i:external input

10. Heteroassociative Network

Layer-1

$$\frac{dx_i(t)}{dt} = -A_i x_i(t) + \sum_{j=1}^{M} f_j(y_j(t)) V_{ji} + I_i$$

Symbols

- $x_i(t)$:activation of neuron $i$in layer-1
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change
- A_i:decay constant
- $\sum_{j=1}^{M}$:summation overlayer-2neurons

- j:index of neuron in layer-2
- M:number of neurons in layer-2
- $f_j(\cdot)$:output function of neuron$j$
- $y_j(t)$:activation of neuron j inlayer-2
- V_{ji}:weight from neuron j to neuron$i$
- I_i:external input

Layer-2

$$\frac{dy_j(t)}{dt} = -B_j y_j(t) + \sum_{i=1}^{N} g_i(x_i(t)) W_{ij} + J_j$$

Symbols

- $y_j(t)$:activation of neuron j in layer-2
- t:time
- $\frac{dy_j(t)}{dt}$:rate of change
- B_j:decay constant
- $\sum_{i=1}^{N}$:summation overlayer-1neurons
- i:neuron index in layer-1
- N:number of neurons in layer-1
- $g_i(\cdot)$:output function of neuron$i$
- $x_i(t)$:activation of neuron i
- W_{ij}:weight from neuron i to neuron$j$
- J_j:external input

11. Bidirectional Associative Memory (BAM)

Same equations asheteroassociative net work with

- $V = W^T$:weight matrix V equals transpose of weight matrix W

12. Basic Shunting Model

$$\frac{dx_i(t)}{dt} = -A_i x_i(t) + (B_i - x_i(t)) I_i$$

Symbols

- $x_i(t)$:activation
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change
- A_i:decay constant
- B_i: upper saturation limit
- I_i:external input
- $(B_i - x_i(t))$:remaining activation capacity

13. On-CenterOff-Surround Model

$$\frac{dx_i(t)}{dt} = -A_i x_i(t) + (B_i - x_i(t)) I_i - x_i(t) \sum_{j \neq i} I_j$$

Symbols

- $x_i(t)$:activation
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change
- A_i:decay constant
- B_i:upper bound
- I_i: excitatory input
- $\sum_{j \neq i}$:summation overall neurons except i
- $j \neq i$: index condition
- I_j:inhibitory inputs

14. Modified Shunting Model

$$\frac{dx_i(t)}{dt} = -A_i x_i(t) + (B_i - x_i(t)) I_i - (E_i + x_i(t)) \sum_{j \neq i} I_j$$

Symbols

- $x_i(t)$:activation
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change
- A_i:decay constant
- B_i:upper bound
- E_i: lower bound parameter
- I_i: excitatory input
- $\sum_{j \neq i}$:summation excluding neuron i
- I_j:inhibitory inputs

15. Shunting Model with Feedback

$$\frac{dx_i(t)}{dt} = -A_i x_i + (B_i - C_i x_i)[I_i + f_i(x_i)] - (E_i + D_i x_i)[J_i + \sum_{j \neq i} f_j(x_j) w_{ji}]$$

Symbols

- x_i:activation of neuron i
- t:time
- $\frac{dx_i(t)}{dt}$:rate of change
- A_i:decay constant
- B_i:upper bound
- C_i:scaling constant for excitatory term
- I_i:external excitatory input
- $f_i(x_i)$:feedback function of neuron i

- E_i:lower bound parameter
- D_i:scaling constant for inhibitory term
- J_i: external inhibitory input
- $\sum_{j \neq i}$:summation over neurons except i
- $f_j(x_j)$:output of neuron j
- w_{ji}:feedback weight from neuron j to neuron i

16. Synaptic Dynamics Model

$$\frac{dw_{ij}(t)}{dt} = -w_{ij}(t) + f_i(x_i(t)) f_j(x_j(t))$$

Symbols

- $w_{ij}(t)$:synaptic weight from neuron j to neuron i at time t
- $\frac{dw_{ij}(t)}{dt}$:rate of change of weight
- t:time
- $f_i(x_i(t))$: output of neuron i
- $x_i(t)$:activation of neuron i
- $f_j(x_j(t))$: output of neuron$j$
- $x_j(t)$: activation of neuron j