

Estimation

How to get information from samples

{Arnaud Legrand, Lucas Schnorr, Jean-Marc Vincent}@univ-grenoble-alpes.fr

University Grenoble-Alpes, UFR IM²AG



October 2025



ESTIMATION

1 THE PROBLEM : Estimate a System Parameter

2 MODELLING : Stochastic Model

3 Synthesis

PROTOCOL VALIDATION

Typical random access protocol to a common channel (CSMA family)

```
Protocol-Send ( $M$ )  
  while Message is not sent  
  | Send( $M$ )  
  | if Collision  
  | |  $W = \text{Random}(I_n)$   
  | |  $n = n + 1$   
  | |  $I_{n+1} = g(n, I_n)$   
  | | Wait ( $W$ )
```

PROTOCOL VALIDATION

Typical random access protocol to a common channel (CSMA family)

```
Protocol-Send ( $M$ )  
  while Message is not sent  
  | Send( $M$ )  
  | if Collision  
  | |  $W = \text{Random}(I_n)$   
  | |  $n = n + 1$   
  | |  $I_{n+1} = g(n, I_n)$   
  | | Wait ( $W$ )
```

What should be the *amount of time* ?

PROTOCOL VALIDATION

Typical random access protocol to a common channel (CSMA family)

```
Protocol-Send ( $M$ )  
  while Message is not sent  
  | Send( $M$ )  
  | if Collision  
  | |  $W = \text{Random}(I_n)$   
  | |  $n = n + 1$   
  | |  $I_{n+1} = g(n, I_n)$   
  | | Wait ( $W$ )
```

What should be the *amount of time* ?

Protocol dimensioning

Waiting time :

- ▶ Random
- ▶ Uniformly distributed on an interval $[0, I_n]$
- ▶ Length of the interval depends on the number of collisions
- ▶ Adaptive scheme $I_{n+1} = 2 \times I_n$,
- ▶ I_0 fixed, characteristic of the protocol

PROTOCOL HISTORY

University of Hawaii 1970

<http://www.hicss.hawaii.edu/>



Norman Abramson et al.

Use of a radio network to provide computer communications without centralization or vacations

PROTOCOL HISTORY

University of Hawaiï 1970

<http://www.hicss.hawaii.edu/>



Norman Abramson et al.

Use of a radio network to provide computer communications without centralization or vacations

Ancestor of CSMA/CD (ethernet), CSMA/CA (WiFi)...

QUANTITATIVE SPECIFICATION VALIDATION

Experiment

Propose an experiment to check the specification of the protocol

QUANTITATIVE SPECIFICATION VALIDATION

Experiment

Propose an experiment to check the specification of the protocol

Estimation

How could I_0 be estimated ?

QUANTITATIVE SPECIFICATION VALIDATION

Experiment

Propose an experiment to check the specification of the protocol

Estimation

How could I_0 be estimated ?

Decision

How could you conclude on the validity of the implementation of the protocol ?

ESTIMATION

1 THE PROBLEM : Estimate a System Parameter

2 **MODELLING : Stochastic Model**

3 Synthesis

SAMPLES

The observations : a sequence of n waiting times.

We suppose that the experiment has been well driven (no outliers, ...) and the observed values denoted by

$$\{x_1, \dots, x_n\}.$$

The stochastic model : the observations are considered to be realizations of independent random variables with the same probability law F

$$\{X_1, \dots, X_n\}$$

Question : What could be said on F from the observations $\{x_1, \dots, x_n\}$?

A priori knowledge :

- ▶ The shape of the law is known and depends on some parameter(s) unknown : **parametric estimation**

ex: X follows a uniform distribution on $[0, \theta[$ with θ unknown

$$F_{\theta}(x) = \begin{cases} 0 & x \leq 0; \\ \frac{1}{\theta}x & x \in [0, \theta[; \\ 1 & \theta \leq x. \end{cases}$$

- ▶ The shape of the law is unknown and some parameters are under study (expectation, variance, moments,...) : **non-parametric estimation**

BASIC CONCEPTS

- ▶ A **statistic** is a function of the observations : $t_n(x_1, \dots, x_n)$, it usually summarize some parameter of the distribution.
- ▶ An **estimator** is a random variable $T_n = t(X_1, \dots, X_n)$ (model of the statistic)

Example

$t(x_1, \dots, x_n) = \max_{1 \leq i \leq n} x_i$ is a statistic on the samples;

and $T_n = \max_{1 \leq i \leq n} X_i$ the corresponding estimator.

Law of T_n under the hypothesis X_i uniformly distributed on $[0, \theta[$:

$$F_n^\theta(x) = \mathbb{P}(\max_{1 \leq i \leq n} X_i \leq x) = \frac{1}{\theta^n} x^n \text{ using independence and uniformity law of } X_i;$$

and density

$$f_n^\theta(x) = \frac{1}{\theta^n} n x^{n-1}.$$

BIAS

An estimator T_n of some parameter θ is **unbiased** iff

$$\mathbb{E}T_n = \theta$$

Example

Bias of estimator T_n

$$\mathbb{E}T_n = \int_0^\theta x f_n^\theta(x) dx = \int_0^\theta x \frac{1}{\theta^n} n x^{n-1} dx = \frac{1}{\theta^n} n \left[\frac{x^{n+1}}{n+1} \right]_0^\theta = \frac{n}{n+1} \theta \neq \theta.$$

T_n is a biased estimator, on average it underestimates θ on average. But, for large samples the bias decreases to 0.

$$\lim_{n \rightarrow +\infty} \mathbb{E}T_n = \theta \quad T_n \text{ is asymptotically unbiased (or consistent)}$$

To compensate the bias,

$$T'_n = \frac{n+1}{n} T_n,$$

which is unbiased.

RISK

The quality T_n of an estimator is evaluated by the **risk** function

$$R(T_n) = \mathbb{E}(T_n - \theta)^2 = \mathbb{E}(T_n - \mathbb{E}T_n)^2 + (\mathbb{E}T_n - \theta)^2 = \mathbb{V}arT_n + (\mathbb{E}T_n - \theta)^2$$

- ▶ $\mathbb{V}arT_n$: the concentration of the distribution
- ▶ $(\mathbb{E}T_n - \theta)^2$: impact of the bias.

For an unbiased estimator

$$R(T_n) = \mathbb{V}arT_n$$

Risk of T'_n

For the example

$$\begin{aligned}\mathbb{V}ar T_n &= \mathbb{E}(T_n - \mathbb{E}T_n)^2 = \mathbb{E}T_n^2 - (\mathbb{E}T_n)^2 = \int_0^\theta x^2 f_n^\theta(x) dx - \left(\frac{n}{n+1}\theta\right)^2 \\ &= \left(\frac{n}{n+2} - \frac{n^2}{(n+1)^2}\right) \theta^2\end{aligned}$$

$$\begin{aligned}\mathbb{V}ar T'_n &= \mathbb{V}ar \frac{n+1}{n} T_n = \frac{(n+1)^2}{n^2} \mathbb{V}ar T_n = \frac{(n+1)^2}{n^2} \left(\frac{n}{n+2} - \frac{n^2}{(n+1)^2}\right) \theta^2 \\ &= \frac{1}{n(n+2)} \theta^2\end{aligned}$$

ANOTHER ESTIMATOR

Consider

$$U_n = \frac{2}{n} \sum_{i=1}^n X_i.$$

$$\mathbb{E}U_n = \frac{2}{n} \sum_{i=1}^n \mathbb{E}X_i = \frac{2}{n} \sum_{i=1}^n \frac{\theta}{2} = \theta.$$

U_n is an unbiased estimator of θ .

Risk of U_n

For the example

$$\begin{aligned} \mathbb{V}ar U_n &= \mathbb{V}ar \left(\frac{2}{n} \sum_{i=1}^n X_i \right); \\ &= \left(\frac{2}{n} \right)^2 \sum_{i=1}^n \mathbb{V}ar X_i \text{ because of the independence of } X_i; \\ &= \frac{4}{n^2} n \frac{\theta^2}{12} = \frac{\theta^2}{3n}. \end{aligned}$$

The risk associated to U_n is much larger than the risk of T'_n so we will prefer T'_n .

ESTIMATION

1 THE PROBLEM : Estimate a System Parameter

2 MODELLING : Stochastic Model

3 **Synthesis**

SYNTHESIS

Observations are realizations of a same experiment in the same conditions

Parametric Estimation

- Assumptions : the results of the experiments are modeled by a sequence of independent identically distributed with a **known probability law**;
- An **estimator** is a random variable function of the modeled random variables, expecting to approximate a parameter of the law;
- The **biais** of an estimator is the expected difference between the estimator and the real value $\mathbb{E}_\theta(T_n - \theta)$
- The **risk** is the expectation of the error around the real value $\mathbb{E}_\theta(T_n - \theta)^2$

Non-Parametric Estimation : estimation of the mean

- Assumptions : the results of the experiments are modeled by a sequence of independent identically distributed with a **unknown probability law**, but with some properties mean m and variance σ^2 ;
- The average $\frac{1}{n} \sum X_i$ is an unbiased estimator of m ;
- The error $\frac{1}{n} \sum X_i - m$ converges almost surely to 0 (strong law of large numbers)
- The law of errors satisfies the central limit theorem (central limit theorem and confidence intervals)

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{X_i - m}{\sigma} \xrightarrow{d} \mathcal{N}(0, 1)$$