

## Numerical Integration Techniques

Error in  $d$  dimensions with  $n$  points:

Sparse Grids:  $\mathcal{O}(n^{-1/d})$

Monte Carlo (MC):  $\mathcal{O}(n^{-1/2})$

Quasi-Monte Carlo (QMC):  $\mathcal{O}(n^{-1})$

**MC & QMC avoid the curse of dimensionality.**  
**QMC is more efficient than MC.**

## Applications

Applied Statistics, Finance, Computer Graphics, AI,  
Computational Sciences, Engineering, Simulation, ...

## The (Quasi-)Monte Carlo Problem

### Integral Transform

$$\mu = \int_{\mathcal{T}} g(\mathbf{t}) \lambda(\mathbf{t}) d\mathbf{t} = \int_{[0,1]^d} f(\mathbf{x}) d\mathbf{x}$$

$g : \mathcal{T} \rightarrow \mathbb{R}$  = original integrand

$\lambda$  = true measure weight

$\mathbf{T} : [0,1]^d \rightarrow \mathcal{T}$  = change of variables


$f : [0,1]^d \rightarrow \mathbb{R}$  = integrand after change of variables

### (Quasi-)Monte Carlo Approximation

$$\hat{\mu}_n = \frac{1}{n} \sum_{i=1}^n f(\mathbf{x}_i) \approx \int_{[0,1]^d} f(\mathbf{x}) d\mathbf{x} = \mu$$

discrete distribution =  $\{\mathbf{x}_1, \mathbf{x}_2, \dots\} \sim \mathcal{U}[0,1]^d$

## Monte Carlo vs Quasi-Monte Carlo



Choose discrete distribution  $\{\mathbf{x}_i\}_{i=1}^n$  to be

IID (independent ...) - Monte Carlo (left)

LD (low discrepancy) - Quasi-Monte Carlo (right)

## Pricing an Asian Call Option

Time Horizon  $\tau = 1$  Interest Rate  $r = 0$  Volatility  $\sigma = 0.5$  Start Price  $S_0 = 100$  Strike Price  $K = 150$

True Measure  $\mathbf{T} = (T_1, \dots, T_d) \sim \mathcal{N}(\mathbf{0}, \Sigma)$ ,  $\Sigma = (\tau/d)(\min(k, j))_{j,k=1}^d$


Asset Path  $S_j(\mathbf{T}) = S_0 \exp((r - \sigma^2/2)\tau j/d + \sigma T_j)$ ,  $j = 1, \dots, d$

Payoff  $\text{Pay}(\mathbf{S}) = \max\left(\frac{1}{2d} \sum_{j=1}^d (S_{j-1} + S_j) - K, 0\right)$ ,  $\mathbf{S} = (S_0, \dots, S_d)$

Discounted Payoff  $g(\mathbf{t}) = \text{Pay}(\mathbf{S}(\mathbf{t})) \exp(-r\tau)$

$$\mu = \mathbb{E}(g(\mathbf{T})) = \int_{\mathbb{R}^d} g(\mathbf{t}) \underbrace{(2\pi)^{-d/2} \det(\Sigma)^{-1/2} \exp(-\mathbf{t}^T \Sigma^{-1} \mathbf{t}/2)}_{\mathcal{N}(\mathbf{0}, \Sigma) \text{ density}} d\mathbf{t}$$

```
>>> import qmcpy as qp
>>> sobol = qp.Sobol(dimension=365) # LD sequence, daily monitoring
>>> params = {"start_price":100, "strike_price":150, "call_put":"call"}
>>> aco = qp.AsianOption(sobol,**params) # defaults other params
>>> algorithm = qp.CubQMCSobolG(aco,abs_tol=1e-2) # tol of a penny
>>> solution,data = algorithm.integrate() # determine n to meet tol
>>> print("Approx discounted payoff $%.2f took %.2f seconds, %d samples."%
...      (solution,data.time_integrate,data.n_total)) # print(data) for details
Approx discounted payoff $1.47 took 0.81 seconds, 16384 samples.
```



## Contributing Projects

- Guaranteed Automatic Integration Library [3]
- Quasi-Random Number Generators [4]
- P. Robbe's Multilevel MC and QMC [5]
- M. Giles' Multilevel MC and QMC [6,7]
- A. Owen's Halton Generator [8]
- LatNet Builder Generating Vectors [9]

## References

- S.-C. T. Choi, F. J. Hickernell, R. Jagadeeswaran, M. J. McCourt, A. G. Sorokin, "Quasi-Monte Carlo Software," Monte Carlo and Quasi-Monte Carlo Methods 2020.
- S.-C. T. Choi, F. J. Hickernell, R. Jagadeeswaran, M. J. McCourt, A. G. Sorokin, "QMCPy: A Quasi-Monte Carlo Python Library," [www.qmcpy.org](http://www.qmcpy.org).
- S.-C. T. Choi, Y. Ding, F. J. Hickernell, L. Jiang, D. Li, R. Jagadeeswaran, L.-A. Jimenez Rugama, X. Tong, K. Zhang, Y. Zhang, and X. Zhou, "GAIL: Guaranteed Automatic Integration Library," Version 2.3.1, MATLAB Software, 2020.
- M. Hofert and C. Lemieux, "qrng: (Randomized) Quasi-Random Number Generators," R package version 0.0-7, (2019).
- P. Robbe. "Multilevel Uncertainty Quantification Methods for Robust Design of Industrial Applications," PhD thesis, KU Leuven, 2019.
- M. B. Giles. "Improved multilevel Monte Carlo convergence using the Milstein scheme," p.343-358 in Monte Carlo and Quasi-Monte Carlo Methods 2006, Springer, 2008.
- M. B. Giles and B. J. Waterhouse. "Multilevel quasi-Monte Carlo path simulation," pp.165-181 in Advanced Financial Modelling, in Radon Series on Computational and Applied Mathematics, de Gruyter, 2009.
- A. B. Owen. "A randomized Halton algorithm in R2017." arXiv:1706.02808 [stat.CO].
- P. L'Ecuyer, P. Marion, M. Godin, and F. Puchhammer, "A Tool for Custom Construction of QMC and RQMC Point Sets," Monte Carlo and Quasi-Monte Carlo Methods 2020.