

# Stats theory (SDA)

Alejandro Caceres

2022-10-31



# Contents

<b>1</b>	<b>About</b>	<b>11</b>
1.1	Recommended reading list . . . . .	11
<b>2</b>	<b>Data description</b>	<b>13</b>
2.1	Objective . . . . .	13
2.2	Statistics . . . . .	13
2.3	Scientific method . . . . .	13
2.4	Outcome . . . . .	13
2.5	Types of outcome . . . . .	14
2.6	Random experiments . . . . .	14
2.7	Absolute frequencies . . . . .	14
2.8	Example . . . . .	15
2.9	Relative frequencies . . . . .	15
2.10	Example . . . . .	16
2.11	Bar plot . . . . .	16
2.12	Pie chart . . . . .	17
2.13	Categorical and ordered variables . . . . .	17
2.14	Example . . . . .	18
2.15	Absolute and relative cumulative frequencies . . . . .	18
2.16	Frequency table . . . . .	18
2.17	Cumulative frequency plot . . . . .	19
2.18	Continuous variables . . . . .	19
2.19	Bins . . . . .	20
2.20	Create a categorical variable from a continuous one . . . . .	20
2.21	Frequency table for a continuous variable . . . . .	21
2.22	Histogram . . . . .	21
2.23	Histogram . . . . .	22
2.24	Cumulative frequency plot: Continuous variables . . . . .	23
2.25	Summary statistics . . . . .	24
2.26	Average . . . . .	25
2.27	Average (categorical ordered) . . . . .	25
2.28	Average (categorical ordered) . . . . .	25
2.29	Average . . . . .	26

2.30	Average . . . . .	26
2.31	Median . . . . .	26
2.32	Median Vs Average . . . . .	27
2.33	Dispersion . . . . .	28
2.34	Dispersion . . . . .	29
2.35	Sample variance . . . . .	29
2.36	Sample variance . . . . .	29
2.37	Standard deviation . . . . .	30
2.38	IQR . . . . .	30
2.39	IQR . . . . .	31
2.40	Box plot . . . . .	31
<b>3</b>	<b>Probability</b>	<b>33</b>
3.1	Objective . . . . .	33
3.2	Random experiments . . . . .	33
3.3	Probability . . . . .	34
3.4	Example . . . . .	34
3.5	Example . . . . .	34
3.6	Relative frequency . . . . .	34
3.7	At infinity . . . . .	35
3.8	Frequentist probability . . . . .	36
3.9	Classical Probability . . . . .	36
3.10	Classical and frequentist probabilities . . . . .	37
3.11	Probability . . . . .	37
3.12	Sample space . . . . .	37
3.13	Examples of sample spaces . . . . .	38
3.14	Discrete and continuous sample spaces . . . . .	38
3.15	Event . . . . .	38
3.16	Event operations . . . . .	39
3.17	Event operations example . . . . .	39
3.18	Outcomes . . . . .	39
3.19	Probability definition . . . . .	40
3.20	Probability properties . . . . .	40
3.21	Addition Rule . . . . .	41
3.22	Example Addition Rule . . . . .	41
3.23	Venn diagram . . . . .	41
3.24	Probability table . . . . .	42
3.25	Example probability table . . . . .	42
3.26	Contingency table . . . . .	42
3.27	Example contingency table . . . . .	43
3.28	Misophonia study . . . . .	43
3.29	Contingency table for frequencies . . . . .	46
3.30	Heat map . . . . .	47
3.31	Continuous variables . . . . .	47
3.32	Heat map for continuous variables . . . . .	50
3.33	Scatter plot . . . . .	51

<b>4</b>	<b>Conditional Probability</b>	<b>53</b>
4.1	Objective . . . . .	53
4.2	Joint Probability . . . . .	53
4.3	Diagnostics . . . . .	54
4.4	Diagnostics Test . . . . .	54
4.5	Observations . . . . .	54
4.6	Contingency tables . . . . .	55
4.7	Conditional probability . . . . .	55
4.8	Conditional probability . . . . .	56
4.9	Conditional contingency table . . . . .	56
4.10	Example conditional contingency table . . . . .	57
4.11	Multiplication rule . . . . .	57
4.12	Diagnostic performance . . . . .	57
4.13	Multiplication rule . . . . .	58
4.14	Contingency table in terms of conditional probabilities . . . . .	58
4.15	Conditional tree . . . . .	59
4.16	Contingency table in terms of conditional probabilities . . . . .	59
4.17	Total probability rule . . . . .	59
4.18	Conditional tree . . . . .	60
4.19	Finding reverse probabilities . . . . .	60
4.20	Recover joint probabilities . . . . .	60
4.21	Reverse conditionals . . . . .	60
4.22	Baye's theorem . . . . .	61
4.23	Example: Bayes' theorem . . . . .	62
4.24	Example: Bayes' theorem . . . . .	62
4.25	Statistical independence . . . . .	63
4.26	Statistical independence . . . . .	63
4.27	Statistical independence . . . . .	63
4.28	Statistical independence . . . . .	64
4.29	Products of marginals products . . . . .	64
4.30	Example . . . . .	65
<b>5</b>	<b>Discrete Random Variables</b>	<b>67</b>
5.1	Objective . . . . .	67
5.2	How do we assign probability values to outcomes? . . . . .	67
5.3	Random variable . . . . .	67
5.4	Random variable . . . . .	68
5.5	Events of observing a random variable . . . . .	68
5.6	Probability of random variables . . . . .	69
5.7	Probability functions . . . . .	69
5.8	Probability functions . . . . .	70
5.9	Probability functions . . . . .	70
5.10	Probability functions . . . . .	71
5.11	Example: Probability mass function . . . . .	71
5.12	Probability table for equally likely outcomes . . . . .	72
5.13	Probability table for $X$ . . . . .	72

5.14	Example . . . . .	72
5.15	Example . . . . .	73
5.16	Probabilities and frequencies . . . . .	73
5.17	Probabilities and relative frequencies . . . . .	74
5.18	Mean and Variance . . . . .	74
5.19	Mean and Variance . . . . .	75
5.20	Mean . . . . .	75
5.21	Example: Mean . . . . .	75
5.22	Mean and Average . . . . .	76
5.23	Variance . . . . .	77
5.24	Example: Variance . . . . .	77
5.25	Functions of $X$ . . . . .	78
5.26	Example: Variance about the origin . . . . .	78
5.27	Probability distribution . . . . .	79
5.28	Example: Probability distribution . . . . .	79
5.29	Probability distribution . . . . .	80
5.30	Probability function and Probability distribution . . . . .	80
5.31	Probability function and Probability distribution . . . . .	81
5.32	Quantiles . . . . .	81
5.33	Summary . . . . .	82
<b>6</b>	<b>Continous Random Variables</b>	<b>83</b>
6.1	Objective . . . . .	83
6.2	Continuous random variable . . . . .	83
6.3	Continuous random variable . . . . .	84
6.4	Continuous random variable . . . . .	84
6.5	Continuous random variable . . . . .	85
6.6	Continuous random variable . . . . .	85
6.7	Total area under the curve . . . . .	86
6.8	Area under the curve . . . . .	86
6.9	Area under the curve . . . . .	87
6.10	Probability distribution . . . . .	88
6.11	Probability distribution . . . . .	90
6.12	Probability distribution . . . . .	90
6.13	Probability distribution . . . . .	91
6.14	Probability graphics . . . . .	91
6.15	Probability graphics . . . . .	92
6.16	Mean . . . . .	93
6.17	Mean . . . . .	93
6.18	Variance . . . . .	94
6.19	Functions of $X$ . . . . .	94
6.20	Example . . . . .	95
<b>7</b>	<b>Discrete Probability Models</b>	<b>97</b>
7.1	Objective . . . . .	97
7.2	Probability mass function . . . . .	97

7.3	Probability model . . . . .	98
7.4	Parametric models . . . . .	98
7.5	Uniform distribution (one parameter) . . . . .	99
7.6	Uniform distribution . . . . .	100
7.7	Uniform distribution (two parameters) . . . . .	100
7.8	Uniform distribution (two parameters) . . . . .	101
7.9	Uniform distribution . . . . .	101
7.10	Uniform distribution (two-parameter) . . . . .	102
7.11	Parameters and Models . . . . .	103
7.12	Parameters and Models . . . . .	103
7.13	Bernoulli trial . . . . .	104
7.14	Bernoulli trial . . . . .	105
7.15	Bernoulli trial . . . . .	105
7.16	Bernoulli trial . . . . .	106
7.17	Binomial distribution . . . . .	106
7.18	Examples: Binomial distribution . . . . .	107
7.19	Binomial distribution . . . . .	107
7.20	Binomial distribution . . . . .	107
7.21	Binomial distribution: Definition . . . . .	108
7.22	Binomial distribution: Mean and Variance . . . . .	109
7.23	Example 1 . . . . .	109
7.24	Example 1 . . . . .	110
7.25	Example 2 . . . . .	110
7.26	Binomial distribution . . . . .	111
7.27	Negative binomial distribution . . . . .	111
7.28	Negative binomial distribution . . . . .	112
7.29	Negative binomial distribution . . . . .	112
7.30	Mean and Variance . . . . .	113
7.31	Geometric distribution . . . . .	113
7.32	Example . . . . .	113
7.33	Example . . . . .	114
7.34	Example . . . . .	114
7.35	Examples . . . . .	114
7.36	Negative binomial distribution . . . . .	115
<b>8</b>	<b>Poisson and Exponential Models</b>	<b>117</b>
8.1	Objective . . . . .	117
8.2	Discrete probability models . . . . .	117
8.3	Counting events . . . . .	118
8.4	Counting events . . . . .	118
8.5	Poisson distribution . . . . .	118
8.6	Poisson distribution . . . . .	118
8.7	Poisson distribution: Derivation details . . . . .	119
8.8	Poisson distribution . . . . .	119
8.9	Poisson distribution . . . . .	120
8.10	Poisson distribution . . . . .	120

8.11 Poisson distribution . . . . .	121
8.12 Continuous probability models . . . . .	121
8.13 Exponential density . . . . .	122
8.14 Exponential density . . . . .	122
8.15 Exponential density . . . . .	122
8.16 Exponential density . . . . .	123
8.17 Exponential density . . . . .	124
8.18 Exponential Distribution . . . . .	124
8.19 Exponential Distribution . . . . .	125
8.20 Exponential Distribution . . . . .	126
<b>9 Normal Distribution</b>	<b>127</b>
9.1 Objective . . . . .	127
9.2 Continuous probability models . . . . .	127
9.3 Normal density . . . . .	128
9.4 Normal density . . . . .	128
9.5 Normal density . . . . .	128
9.6 Normal density . . . . .	128
9.7 Normal density . . . . .	129
9.8 Definition . . . . .	129
9.9 Normal probability density (Gaussian) . . . . .	130
9.10 Normal distribution . . . . .	130
9.11 Normal distribution . . . . .	131
9.12 Normal distribution . . . . .	131
9.13 Normal distribution . . . . .	132
9.14 Normal distribution . . . . .	132
9.15 Standard normal density . . . . .	133
9.16 Standard normal density . . . . .	133
9.17 Standard normal density . . . . .	134
9.18 Normal distribution . . . . .	134
9.19 Standard distribution . . . . .	134
9.20 Standard normal density . . . . .	135
9.21 Standard normal density . . . . .	135
9.22 Normal and standard distributions . . . . .	135
9.23 Normal distribution . . . . .	136
9.24 Summary of probability models . . . . .	136
9.25 R functions of probability models . . . . .	137
<b>10 Sampling Distributions</b>	<b>139</b>
10.1 Objective . . . . .	139
10.2 Normal distribution . . . . .	139
10.3 Example: When we do not know the parameters . . . . .	140
10.4 Example . . . . .	140
10.5 Random sample . . . . .	141
10.6 Example . . . . .	142
10.7 Average or sample mean . . . . .	142



10.8 The average as an estimator . . . . .	142
10.9 Sample variance . . . . .	143
10.10 Sample variance . . . . .	143
10.11 Fitting a model . . . . .	144
10.12 Prediction . . . . .	145
10.13 Inference . . . . .	146
10.14 Example: When we do know the parameters . . . . .	146
10.15 Density for $X$ and $\bar{X}$ . . . . .	146
10.16 Sample mean distribution . . . . .	148
10.17 Inference on the average . . . . .	148
10.18 Density for $\bar{X}$ . . . . .	149
10.19 Inference in the sample variance . . . . .	151
10.20 Probabilities of the sample variance . . . . .	151
10.21 $\chi^2$ -statistic . . . . .	152
10.22 $\chi^2$ -statistic . . . . .	153
<b>11 Exercises</b>	<b>155</b>
11.1 Data description . . . . .	155
11.2 Probability . . . . .	156
11.3 Conditional Probability . . . . .	157
11.4 Random variables . . . . .	160
11.5 Probability Models . . . . .	163
11.6 Sampling and Central Limit Theorem . . . . .	164
11.7 Point Estimators . . . . .	165
11.8 Maximum likelihood . . . . .	166
11.9 Method of moments . . . . .	167
11.10 Confidence intervals . . . . .	168
11.11 Hypothesis testing . . . . .	169



# Chapter 1

## About

- This is the introduction to statistics course from EEBE (UPC).
- Exam dates and additional study material can be found in ATENEA

### 1.1 Recommended reading list

- Douglas C. Montgomery and George C. Runger. “Applied Statistics and Probability for Engineers” 4th Edition. Wiley 2007.



## Chapter 2

# Data description

### 2.1 Objective

- Data: discrete, continuous
- Summarizing data in tables and figures

### 2.2 Statistics

- Solve problems in a systematic way (science, engineering and technology)
- Modern humans use a general **method** historically developed for thousands of years! ... and still under development.
- It has three main components: observation, logic, and generation of new knowledge

---

---

### 2.3 Scientific method

---

---

### 2.4 Outcome

Observation or *Realization*

- an **observation** is the acquisition of a number or a characteristic from an experiment

... 1 0 0 1 0 **1** 0 1 1 ... (the number in bold is an observation in a repetition of the experiment)

### Outcome

- An **outcome** is a possible observation that is the result of an experiment.

1 is an outcome, 0 is the other outcome

---



---

## 2.5 Types of outcome

- **Categorical:** If the result of an experiment can only take discrete values (number of car pieces produced per hour, number of leukocytes in blood)
- **Continuous:** If the result of an experiment can only take continuous values (battery state of charge, engine temperature).

---



---

## 2.6 Random experiments

### Definition:

A **random experiment** is an experiment that gives different outcomes when repeated in the same manner.

### Examples:

- on the same object (person): temperature, sugar levels.
- on different objects but the same measurement: the weight of an animal.
- on events: a number of emails received in an hour.

---



---

## 2.7 Absolute frequencies

When we repeat a random experiment, we record a list of outcomes.

We summarize the **categorical** observations by counting how many times we saw a particular outcome.

**Absolute frequency:**

$$n_i$$

is the number of times we observed the outcome  $i$

---



---

## 2.8 Example

**Random experiment:** Extract a leukocyte from **one** donor and write down its type. Repeat experiment  $N = 119$  times.

(T cell, Tcell, Neutrophil, ..., B cell)

```
##      outcome ni
## 1      T Cell 34
## 2      B cell 50
## 3    basophil 20
## 4    Monocyte  5
## 5 Neutrophil 10
```

- For instance:  $n_1 = 34$  is total number of T cells
- $N = \sum_i n_i = 119$

---



---

## 2.9 Relative frequencies

We can also summarize the observations by computing the **proportion** of how many times we saw a particular outcome.

$$f_i = n_i/N$$

where  $N$  is the total number of observations

In our example there are recorded  $n_1 = 34$  T cells, so we ask for the proportion of T cells from the total of 119.

---



---

## 2.10 Example

```
##      outcome ni      fi
## 1    T Cell 34 0.28571429
## 2    B cell 50 0.42016807
## 3  basophil 20 0.16806723
## 4 Monocyte  5 0.04201681
## 5 Neutrophil 10 0.08403361
```

We have


$$\sum_{i=1..M} n_i = N$$

$$\sum_{i=1..M} f_i = 1$$

where  $M$  is the number of outcomes.

## 2.11 Bar plot

We can plot  $n_i$  Vs the outcomes, giving us a bar plot






## 2.12 Pie chart

We can visualize the relative frequencies with a pie chart

- Where the area of the circle represents 100% of observations (proportion = 1) and the sections the relative frequencies of all the outcomes.




---



---

## 2.13 Categorical and ordered variables

Cell types are not meaningfully ordered concerning the outcomes. However, sometimes **categorical** variables can be **ordered**.

Misophonia study:

- 123 patients were examined for misophonia: anxiety/anger produced by certain sounds
- They were categorized into 4 different groups according to severity.

---



---

## 2.14 Example

The results of the study are:

```
## [1] 4 2 0 3 0 0 2 3 0 3 0 2 2 0 2 0 0 3 3 0 3 3 2 0 0 0 4 2 2 0 2 0 0 0 3 0 2
## [38] 3 2 2 0 2 3 0 0 2 2 3 3 0 0 4 3 3 2 0 2 0 0 0 2 2 0 0 2 3 0 1 3 2 4 3 2 3
## [75] 0 2 3 2 4 1 2 0 2 0 2 0 2 2 4 3 0 3 0 0 0 2 2 1 3 0 0 3 2 1 3 0 4 4 2 3 3
## [112] 3 0 3 2 1 2 3 3 4 2 3 2
```

And its frequency table

```
## outcome ni          fi
## 1          0 41 0.33333333
## 2          1  5 0.04065041
## 3          2 37 0.30081301
## 4          3 31 0.25203252
## 5          4  9 0.07317073
```

---



---

## 2.15 Absolute and relative cumulative frequencies

Misophonia severity is **categorical** and **ordered**.

When outcomes can be ordered then it is useful to ask how many observations were obtained up to a given outcome we call this number the absolute cumulative frequency up to the outcome  $i$ :

$$N_i = \sum_{k=1..i} n_k$$

It is also useful to compute the **proportion** of the observations that was obtained up to a given outcome

$$F_i = \sum_{k=1..i} f_k$$


---




---

## 2.16 Frequency table

```
## outcome ni          fi  Ni          Fi
## 0          0 41 0.33333333 41 0.33333333
## 1          1  5 0.04065041 46 0.3739837
```



##	2	2	37	0.30081301	83	0.6747967
##	3	3	31	0.25203252	114	0.9268293
##	4	4	9	0.07317073	123	1.0000000

- **67%** of patients had misophonia up to severity **2**
- **37%** of patients have severity less or equal than **1**



## 2.17 Cumulative frequency plot

We can also plot the cumulative frequency Vs the outcomes



## 2.18 Continuous variables

The result of a random experiment can also give continuous outcomes.

In the misophonia study, the researchers asked whether the convexity of the jaw would affect the misophonia severity (the scientific hypothesis is that the

convexity angle of the jaw can influence the ear and its sensitivity). These are the results for the convexity of the jaw (degrees)

```
## [1] 7.97 18.23 12.27 7.81 9.81 13.50 19.30 7.70 12.30 7.90 12.60 19.00
## [13] 7.27 14.00 5.40 8.00 11.20 7.75 7.94 16.69 7.62 7.02 7.00 19.20
## [25] 7.96 14.70 7.24 7.80 7.90 4.70 4.40 14.00 14.40 16.00 1.40 9.76
## [37] 7.90 7.90 7.40 6.30 7.76 7.30 7.00 11.23 16.00 7.90 7.29 6.91
## [49] 7.10 13.40 11.60 -1.00 6.00 7.82 4.80 11.00 9.00 11.50 16.00 15.00
## [61] 1.40 16.80 7.70 16.14 7.12 -1.00 17.00 9.26 18.70 3.40 21.30 7.50
## [73] 6.03 7.50 19.00 19.01 8.10 7.80 6.10 15.26 7.95 18.00 4.60 15.00
## [85] 7.50 8.00 16.80 8.54 7.00 18.30 7.80 16.00 14.00 12.30 11.40 8.50
## [97] 7.00 7.96 17.60 10.00 3.50 6.70 17.00 20.26 6.64 1.80 7.02 2.46
## [109] 19.00 17.86 6.10 6.64 12.00 6.60 8.70 14.05 7.20 19.70 7.70 6.02
## [121] 2.50 19.00 6.80
```

---



---

## 2.19 Bins

Continuous outcomes cannot be counted!

We transform them into ordered categorical variables

- We cover the range of the observations into regular intervals of the same size (bins)

```
## [1] "[-1.02,3.46]" "(3.46,7.92]" "(7.92,12.4]" "(12.4,16.8]" "(16.8,21.3]"
```

---



---

## 2.20 Create a categorical variable from a continuous one

- We map each observation to its interval: creating an **ordered categorical** variable; in this case with 5 possible outcomes

```
## [1] "(7.92,12.4]" "(16.8,21.3]" "(7.92,12.4]" "(3.46,7.92]" "(7.92,12.4]"
## [6] "(12.4,16.8]" "(16.8,21.3]" "(3.46,7.92]" "(7.92,12.4]" "(3.46,7.92]"
## [11] "(12.4,16.8]" "(16.8,21.3]" "(3.46,7.92]" "(12.4,16.8]" "(3.46,7.92]"
## [16] "(7.92,12.4]" "(7.92,12.4]" "(3.46,7.92]" "(7.92,12.4]" "(12.4,16.8]"
## [21] "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]" "(16.8,21.3]" "(7.92,12.4]"
## [26] "(12.4,16.8]" "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]"
## [31] "(3.46,7.92]" "(12.4,16.8]" "(12.4,16.8]" "(12.4,16.8]" "[-1.02,3.46]"
## [36] "(7.92,12.4]" "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]"
```

```
## [41] "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]" "(7.92,12.4]" "(12.4,16.8]"
## [46] "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]" "(12.4,16.8]"
## [51] "(7.92,12.4]" "[-1.02,3.46]" "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]"
## [56] "(7.92,12.4]" "(7.92,12.4]" "(7.92,12.4]" "(12.4,16.8]" "(12.4,16.8]"
## [61] "[-1.02,3.46]" "(12.4,16.8]" "(3.46,7.92]" "(12.4,16.8]" "(3.46,7.92]"
## [66] "[-1.02,3.46]" "(16.8,21.3]" "(7.92,12.4]" "(16.8,21.3]" "[-1.02,3.46]"
## [71] "(16.8,21.3]" "(3.46,7.92]" "(3.46,7.92]" "(3.46,7.92]" "(16.8,21.3]"
## [76] "(16.8,21.3]" "(7.92,12.4]" "(3.46,7.92]" "(3.46,7.92]" "(12.4,16.8]"
## [81] "(7.92,12.4]" "(16.8,21.3]" "(3.46,7.92]" "(12.4,16.8]" "(3.46,7.92]"
## [86] "(7.92,12.4]" "(12.4,16.8]" "(7.92,12.4]" "(3.46,7.92]" "(16.8,21.3]"
## [91] "(3.46,7.92]" "(12.4,16.8]" "(12.4,16.8]" "(7.92,12.4]" "(7.92,12.4]"
## [96] "(7.92,12.4]" "(3.46,7.92]" "(7.92,12.4]" "(16.8,21.3]" "(7.92,12.4]"
## [101] "(3.46,7.92]" "(3.46,7.92]" "(16.8,21.3]" "(16.8,21.3]" "(3.46,7.92]"
## [106] "[-1.02,3.46]" "(3.46,7.92]" "[-1.02,3.46]" "(16.8,21.3]" "(16.8,21.3]"
## [111] "(3.46,7.92]" "(3.46,7.92]" "(7.92,12.4]" "(3.46,7.92]" "(7.92,12.4]"
## [116] "(12.4,16.8]" "(3.46,7.92]" "(16.8,21.3]" "(3.46,7.92]" "(3.46,7.92]"
## [121] "[-1.02,3.46]" "(16.8,21.3]" "(3.46,7.92]"
```

---



---

## 2.21 Frequency table for a continuous variable

```
##      outcome ni      fi  Ni      Fi
## 1 [-1.02,3.46] 8 0.06504065 8 0.06504065
## 2 (3.46,7.92] 51 0.41463415 59 0.47967480
## 3 (7.92,12.4] 26 0.21138211 85 0.69105691
## 4 (12.4,16.8] 20 0.16260163 105 0.85365854
## 5 (16.8,21.3] 18 0.14634146 123 1.00000000
```


---



---



## 2.22 Histogram

The histogram is the plot of  $n_i$  or  $f_i$  Vs the outcomes (bins). The histogram depends on the size of the bins




## 2.23 Histogram

The histogram is the plot of  $n_i$  or  $f_i$  Vs the outcomes (bins). The histogram depends on the size of the bins



## 2.24 Cumulative frequency plot: Continous variables

We can also plot the cumulative frequency Vs the outcomes



## 2.25 Summary statistics

The summary statistics are numbers computed from the data that tell us important features of numerical variables (categorical or continuous).

Limiting values:

- minimum: the minimum outcome observed
- maximum: the maximum outcome observed

Central value for the outcomes

- The average is defined as

$$\bar{x} = \frac{1}{N} \sum_{j=1..N} x_j$$

where  $x_j$  is the **observation**  $j$  (convexity) from a total of  $N$ .



## 2.26 Average

The average convexity can be computed directly from the **observations**

$$\begin{aligned}\bar{x} &= \frac{1}{N} \sum_j x_j \\ &= \frac{1}{N} (7.97 + 18.23 + 12.27 \dots + 6.80) = 10.19894\end{aligned}$$


---



---

## 2.27 Average (categorical ordered)

For **categorical ordered** variables we can use the frequency table to compute the average

```
## outcome ni      fi
## 1      0 41 0.33333333
## 2      1  5 0.04065041
## 3      2 37 0.30081301
## 4      3 31 0.25203252
## 5      4  9 0.07317073
```

The average **severity** of misophonia in the study can **also** be computed from the relative frequencies of the **outcomes**

$$\begin{aligned}\bar{x} &= \frac{1}{N} \sum_{i=1 \dots N} x_j = \frac{1}{N} \sum_{i=1 \dots M} x_i * n_i = \sum_{i=1 \dots M} x_i * f_i \\ &= 0 * f_0 + 1 * f_1 + 2 * f_2 + 3 * f_3 + 4 * f_4 = 1.691057\end{aligned}$$

(note the change from  $N$  to  $M$  in the second summation)

---



---

## 2.28 Average (categorical ordered)

In terms of the **outcomes** of categorical ordered variables, the **average** can be written as

$$\bar{x} = \sum_{i=1 \dots M} x_i f_i$$

from a total of  $M$  possible outcomes (number of severity levels).

$\bar{x}$  is the **central value** or center of gravity of the outcomes. As if each outcome had a mass density given by  $f_i$ .


---

## 2.29 Average

- The average is not the result of one observation (random experiment).
- It is the result of a series of observations (sample).
- It describes the number where the observed values balance.

That is why we hear, for instance, that a patient with an infection can infect an average of 2.5 people.

## 2.30 Average



## 2.31 Median

Another measure of centrality is the median. The median  $q_{0.5}$  is the value  $x_p$

$$\text{median}(x) = q_{0.5} = x_p$$

below which we find half of the observations

$$\sum_{x \leq x_p} 1 = \frac{N}{2}$$

or in terms of the frequencies, is the value  $x_p$  that makes the cumulative frequency  $F_p$  equal to 0.5


$$q_{0.5} = \sum_{x \leq x_p} f_x = F_p = 0.5$$

---

---

2.32 Median Vs Average

- Average: Center of mass (compensates distant values)
- Median: Half of the mass




---

---

## 2.33 Dispersion


An important measure of the outcomes is their **dispersion**. Many experiments can share their mean but differ on how dispersed the values are.



---

---

## 2.34 Dispersion




---



---

## 2.35 Sample variance

Dispersion about the mean is measured with the

- The sample variance:

$$s^2 = \frac{1}{N-1} \sum_{j=1..N} (x_j - \bar{x})^2$$

It measures the average square distance of the **observations** to the average. The reason for  $N - 1$  will be explained when we talk about inference.

---



---

## 2.36 Sample variance

- In terms of the frequencies of **categorical and ordered** variables

$$s^2 = \frac{N}{N-1} \sum_x (x - \bar{x})^2 f_x$$

$s^2$  can be thought of as the moment of inertia of the observations.

---



---

### 2.37 Standard deviation

The squared root of the sample variance is called the **standard deviation**  $s$ .

The standard deviation of the convexity angle is

$$s = [\frac{1}{123-1} ((7.97 - 10.19894)^2 + (18.23 - 10.19894)^2 + (12.27 - 10.19894)^2 + \dots)]^{1/2} = 5.086707$$

The jaw convexity deviates from its mean by 5.086707.

---



---

### 2.38 IQR

- Dispersion of data can also be measured with respect to the median by the **interquartile range**
- We define the **first** quartile as the value  $x_p$  that makes the cumulative frequency  $F_p$  equal to 0.25

$$q_{0.25} = \sum_{x \leq x_p} f_x = F_p = 0.25$$

- We also define the **third** quartile as the value  $x_p$  that makes the cumulative frequency  $F_p$  equal to 0.75

$$q_{0.75} = \sum_{x \leq x_p} f_x = F_p = 0.75$$


---



---


## 2.39 IQR

The distance between the third quartile and the first quartile is called the **interquartile range** (IQR) and captures the central 50% of the observations



## 2.40 Box plot

The interquartile range, the median, and the 5% and 95% of the data can be visualized in a **boxplot**, here the values of the outcomes are on the y-axis. The IQR is the box, the median is the line in the middle and the whiskers mark the 5% and 95% of the data.





## Chapter 3

# Probability

### 3.1 Objective

- Definition of probability
  - Probability algebra
  - Joint probability
- 
- 

### 3.2 Random experiments

#### Observation

- An **observation** is the acquisition of a number or a characteristic from an experiment

#### Outcome

- An **outcome** is a possible observation that is the result of an experiment.

#### Random experiment

- An experiment that gives **different** outcomes when repeated in the same manner.
- 
-

### 3.3 Probability

The **probability** of an outcome is a measure of how sure we are to observe that outcome when performing a random experiment.

- 0: We are sure that the observation will **not** happen.
- 1: We are sure that the observation will happen.

---



---

### 3.4 Example

- Consider the following observations of a random experiment:

1 5 1 2 2 1 2 2

- How sure we are to obtain 2 in the following observation?

---



---

### 3.5 Example

The frequency table is

##	outcome	ni	fi
## 1	1	3	0.375
## 2	2	4	0.500
## 3	5	1	0.125

The **relative frequency**  $f_i$

- is a number between 0 and 1.
- measures the proportion of total observations that we observed a particular outcome.
- seems a reasonable probability measure.

As  $f_2 = 0.5$  then we would be half certain to obtain a 2 in the next repetition of the experiment.

---



---

### 3.6 Relative frequency

As a measure of certainty is  $f_i$  enough?

Say we repeated the experiment 12 times more:

1 5 1 2 2 1 2 2 **3 1 1 3 3 1 6 3 5 6 4 4**

The frequency table is now

##	outcome	ni	fi
## 1	1	6	0.3
## 2	2	4	0.2
## 3	3	4	0.2
## 4	4	2	0.1
## 5	5	2	0.1
## 6	6	2	0.1

New outcomes appeared and  $f_2$  is now 0.2, we are now a fifth certain of obtaining 2 in the next experiment... probability should not depend on  $N$




### 3.7 At infinity

Say we repeated the experiment 1000 times:

##	outcome	ni	fi
## 1	1	185	0.185
## 2	2	145	0.145
## 3	3	166	0.166
## 4	4	166	0.166
## 5	5	159	0.159
## 6	6	179	0.179

We find that  $f_i$  is converging to a constant value

$$\lim_{N \rightarrow \infty} f_i = P_i$$



### 3.8 Frequentist probability

We call **Probability**  $P_i$  to the limit when  $N \rightarrow \infty$  of the **relative frequency** of observing the outcome  $i$  in a random experiment.

Championed by Venn (1876)

The frequentist interpretation of probabilities is derived from data/experience (empirical).

- We do not observe  $P_i$ , we observe  $f_i$
- When we **estimate**  $P_i$  with  $f_i$  (typically when  $N$  is large), we write:

$$\hat{P}_i = f_i$$

### 3.9 Classical Probability

Whenever a random experiment has  $M$  possible outcomes that are all **equally likely**, the probability of each outcome is  $\frac{1}{M}$ .

Championed by Laplace (1814).

Since each outcome is **equally probable** we declare complete ignorance and the best we can do is to fairly distribute the same probability to each outcome.

What if I told you that our experiment was the throw of the dice? then

$$P_2 = 1/6 = 0.166666.$$

$$P_i = \lim_{N \rightarrow \infty} \frac{n_i}{N} = \frac{1}{M}$$


---



---

### 3.10 Classical and frequentist probabilities

---



---

### 3.11 Probability

Probability is a number between 0 and 1 that is assigned to each member  $E$  of a collection of **events** of a **sample space** ( $S$ ) from a random experiment.

$$P(E) \in (0, 1)$$

where  $E \in S$

---



---

### 3.12 Sample space

We start by reasoning what are all the possible values (outcomes) that a random experiment could give.

Note that we do not have to observe them in a particular experiment: We are using **reason/logic** and not observation.

**Definition:**

- The set of all possible outcomes of a random experiment is called the **sample space** of the experiment.
- The sample space is denoted as  $S$ .

### 3.13 Examples of sample spaces

- temperature 35 and 42 degrees Celcius
- sugar levels: 70-80mg/dL
- the size of one screw from a production line: 70mm-72mm
- number of emails received in an hour: 0-100
- a dice throw: 1, 2, 3, 4, 5, 6

### 3.14 Discrete and continuous sample spaces

- A sample space is discrete if it consists of a finite or countable infinite set of outcomes.
- A sample space is continuous if it contains an interval (either finite or infinite in length) of real numbers.

### 3.15 Event

#### Definition:

An **event** is a **subset** of the sample space of a random experiment. It is a **collection** of outcomes.

Examples of events:

- The event of a healthy temperature: temperature 37-38 degrees Celsius
- The event of producing a screw with a size: of 71.5mm
- The event of receiving more than 4 emails in an hour.
- The event of obtaining a number less than 3 in the throw of a dice

One event refers to a possible set of **outcomes**.

### 3.16 Event operations

For two events  $A$  and  $B$ , we can construct the following derived events:

- Complement  $A'$ : the event of **not**  $A$
- Union  $A \cup B$ : the event of  $A$  **or**  $B$
- Intersection  $A \cap B$ : the event of  $A$  **and**  $B$




### 3.17 Event operations example

Take

- Event  $A : \{1, 2, 3\}$  a number less or equal to three in the throw of a dice
- Event  $B : \{2, 4, 6\}$  an even number in the throw of a dice

New events:

- Not less than three:  $A' : \{4, 5, 6\}$
- Less or equal to three **or** even:  $A \cup B : \{1, 2, 3, 4, 6\}$
- Less or equal to three **and** even  $A \cap B : \{2\}$



### 3.18 Outcomes

Outcomes are events that are **mutually exclusive**

**Definition:**

Two events denoted as  $E_1$  and  $E_2$ , such that

$$E_1 \cap E_2 = \emptyset$$

They cannot occur at the same time.

Example:

- The outcome of obtaining 1 **and** the outcome of obtaining 5 in the throw of one dice are mutually exclusive:
- The event of obtaining 1 and 5 is empty:

$$\{1\} \cap \{5\} = \emptyset$$

### 3.19 Probability definition

A probability is a number that is assigned to each possible event ( $E$ ) of a sample space ( $S$ ) of a random experiment that satisfies the following properties:

- $P(S) = 1$
- $0 \leq P(E) \leq 1$
- when  $E_1 \cap E_2 = \emptyset$

$$P(E_1 \cup E_2) = P(E_1) + P(E_2)$$

Proposed by Kolmogorov's (1933)

### 3.20 Probability properties

Kolmogorov says that we can build a probability table (likewise the relative frequency table)

outcome	Probability
1	1/6
2	1/6
3	1/6
4	1/6
5	1/6
6	1/6
$P(1 \cup 2 \cup \dots \cup 6)$	1

As  $\{1, 2, 3, 4, 5, 6\}$  are mutually exclusive then

$$P(S) = P(1 \cup 2 \cup \dots \cup 6) = P(1) + P(2) + \dots + P(n) = 1$$



### 3.21 Addition Rule

When  $A$  and  $B$  are not mutually exclusive then:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

Where  $P(A)$  and  $P(B)$  are called the **marginal probabilities**

---



---

### 3.22 Example Addition Rule

Take

- Event  $A : \{1, 2, 3\}$  a number less or equal to three in the throw of a dice
- Event  $B : \{2, 4, 6\}$  an even number in the throw of a dice

then:

- $P(A) : P(1) + P(2) + P(3) = 3/6$
- $P(B) : P(2) + P(4) + P(6) = 3/6$
- $P(A \cap B) : P(2) = 1/6$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 3/6 + 3/6 - 1/6 = 5/6$$

Note:  $P(2)$  appears in  $P(A)$  and  $P(B)$  that's why we subtract it with the intersection

---



---

### 3.23 Venn diagram

Note that can always break down the sample space in **mutually exclusive** sets involving the intersections:

$$S = \{A \cap B, A \cap B', A' \cap B, A' \cap B'\}$$

Marginals:

- $P(A) = P(A \cap B') + P(A \cap B) = 2/6 + 1/6 = 3/6$
- $P(B) = P(A' \cap B) + P(A \cap B) = 2/6 + 1/6 = 3/6$

---



---

### 3.24 Probability table

Let's look at the probability table

outcome	Probability
$A \cap B$	$P(A \cap B)$
$A \cap B'$	$P(A \cap B')$
$A' \cap B$	$P(A' \cap B)$
$A' \cap B'$	$P(A' \cap B')$
sum	1

### 3.25 Example probability table

We also write  $A \cap B$  as  $(A, B)$  and call it the **joint probability** of  $A$  and  $B$

In our example:

outcome	Probability
$(A, B)$	$P(A, B) = 1/6$
$(A, B')$	$P(A, B') = 2/6$
$(A', B)$	$P(A', B) = 2/6$
$(A', B')$	$P(A', B') = 1/6$
sum	1

Note: each outcome has *two* values (one for the characteristic of type  $A$  and another for type  $B$ )

### 3.26 Contingency table

We can organize the probability of **joint outcomes** in a **contingency table**

	$B$	$B'$	sum
$A$	$P(A, B)$	$P(A, B')$	$P(A)$
$A'$	$P(A', B)$	$P(A', B')$	$P(A')$
sum	$P(B)$	$P(B')$	1

Marginals:

- $P(A) = P(A, B') + P(A, B)$
- $P(B) = P(A', B) + P(A, B)$

---



---

### 3.27 Example contingency table

- Event  $A$  :  $\{1, 2, 3\}$  a number less or equal to three in the throw of a dice
- Event  $B$  :  $\{2, 4, 6\}$  an even number in the throw of a dice

	$B$	$B'$	sum
$A$	1/6	2/6	3/6
$A'$	2/6	1/6	3/6
sum	3/6	3/6	1

Three forms of the **addition rule**:

$$\begin{aligned}
 P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\
 &= P(A \cap B) + P(A \cap B') + P(A' \cap B) \\
 &= 1 - P(A' \cap B')
 \end{aligned}$$

---



---

### 3.28 Misophonia study

In the misophonia study, the patients were assessed for their misophonia severity **and** if they were depressed.

The outcome of one random experiment is to measure the misophonia severity **and** depression status of one patient. The repetition of the random experiment was to perform the same two measurements on another patient.

```
##      Misofonia.dic  depression.dic
## 1                4                1
## 2                2                0
## 3                0                0
## 4                3                0
## 5                0                0
```

## 6	0	0
## 7	2	0
## 8	3	0
## 9	0	1
## 10	3	0
## 11	0	0
## 12	2	0
## 13	2	1
## 14	0	0
## 15	2	0
## 16	0	0
## 17	0	0
## 18	3	0
## 19	3	0
## 20	0	0
## 21	3	0
## 22	3	0
## 23	2	0
## 24	0	0
## 25	0	0
## 26	0	0
## 27	4	1
## 28	2	0
## 29	2	0
## 30	0	0
## 31	2	0
## 32	0	0
## 33	0	0
## 34	0	0
## 35	3	0
## 36	0	0
## 37	2	0
## 38	3	1
## 39	2	0
## 40	2	0
## 41	0	0
## 42	2	0
## 43	3	0
## 44	0	0
## 45	0	0
## 46	2	0
## 47	2	0
## 48	3	0
## 49	3	0
## 50	0	0
## 51	0	0

## 52	4	1
## 53	3	0
## 54	3	1
## 55	2	1
## 56	0	1
## 57	2	0
## 58	0	0
## 59	0	0
## 60	0	0
## 61	2	0
## 62	2	0
## 63	0	0
## 64	0	0
## 65	2	0
## 66	3	1
## 67	0	0
## 68	1	0
## 69	3	0
## 70	2	0
## 71	4	1
## 72	3	0
## 73	2	1
## 74	3	0
## 75	0	1
## 76	2	0
## 77	3	0
## 78	2	0
## 79	4	1
## 80	1	0
## 81	2	0
## 82	0	0
## 83	2	0
## 84	0	0
## 85	2	0
## 86	0	1
## 87	2	0
## 88	2	0
## 89	4	1
## 90	3	0
## 91	0	1
## 92	3	0
## 93	0	0
## 94	0	0
## 95	0	0
## 96	2	0
## 97	2	0

## 98	1	0
## 99	3	0
## 100	0	0
## 101	0	0
## 102	3	1
## 103	2	0
## 104	1	0
## 105	3	0
## 106	0	0
## 107	4	1
## 108	4	1
## 109	2	0
## 110	3	0
## 111	3	0
## 112	3	1
## 113	0	0
## 114	3	0
## 115	2	0
## 116	1	0
## 117	2	0
## 118	3	1
## 119	3	0
## 120	4	1
## 121	2	0
## 122	3	0
## 123	2	0

---



---

### 3.29 Contingency table for frequencies

- For the number of observations  $n_{i,j}$  of each outcome  $(x_i, y_i)$ , misophonia:  $x \in \{0, 1, 2, 3, 4\}$  and depression  $y \in \{0, 1\}$  (no:0, yes:1)

##		
##	Depression:0	Depression:1
## Misophonia:4	0	9
## Misophonia:3	25	6
## Misophonia:2	34	3
## Misophonia:1	5	0
## Misophonia:0	36	5

- For the relative frequencies  $f_{i,j}$

##


```
##          Depression:0 Depression:1
## Misophonia:4  0.00000000  0.07317073
## Misophonia:3  0.20325203  0.04878049
## Misophonia:2  0.27642276  0.02439024
## Misophonia:1  0.04065041  0.00000000
## Misophonia:0  0.29268293  0.04065041
```

---

---

3.30 Heat map

The contingency table can be plotted as a **heat map**



3.31 Continous variables

In the misophonia study, the jaw protrusion was also measured as a possible cephalometric factor for de disease.

```
##      Angulo_convexidad protusion.mandibular
## 1          7.97          13.00
```

## 2	18.23	-5.00
## 3	12.27	11.50
## 4	7.81	16.80
## 5	9.81	33.00
## 6	13.50	2.00
## 7	19.30	-3.90
## 8	7.70	16.80
## 9	12.30	8.00
## 10	7.90	28.80
## 11	12.60	3.00
## 12	19.00	-7.90
## 13	7.27	28.30
## 14	14.00	4.00
## 15	5.40	22.20
## 16	8.00	0.00
## 17	11.20	15.00
## 18	7.75	17.00
## 19	7.94	49.00
## 20	16.69	5.00
## 21	7.62	42.00
## 22	7.02	28.00
## 23	7.00	9.40
## 24	19.20	-13.20
## 25	7.96	23.00
## 26	14.70	2.30
## 27	7.24	25.00
## 28	7.80	4.90
## 29	7.90	92.00
## 30	4.70	6.00
## 31	4.40	17.00
## 32	14.00	3.30
## 33	14.40	10.30
## 34	16.00	6.30
## 35	1.40	19.50
## 36	9.76	22.00
## 37	7.90	5.00
## 38	7.90	78.00
## 39	7.40	9.30
## 40	6.30	50.60
## 41	7.76	18.00
## 42	7.30	18.00
## 43	7.00	10.00
## 44	11.23	4.00
## 45	16.00	13.30
## 46	7.90	48.00
## 47	7.29	23.50




## 48	6.91	37.60
## 49	7.10	15.00
## 50	13.40	5.10
## 51	11.60	-2.20
## 52	-1.00	32.00
## 53	6.00	25.00
## 54	7.82	24.00
## 55	4.80	33.60
## 56	11.00	3.30
## 57	9.00	31.50
## 58	11.50	12.80
## 59	16.00	3.00
## 60	15.00	6.00
## 61	1.40	21.40
## 62	16.80	-10.00
## 63	7.70	19.00
## 64	16.14	32.00
## 65	7.12	15.00
## 66	-1.00	10.00
## 67	17.00	-16.90
## 68	9.26	2.00
## 69	18.70	-10.10
## 70	3.40	12.20
## 71	21.30	-11.00
## 72	7.50	5.20
## 73	6.03	16.00
## 74	7.50	5.80
## 75	19.00	5.20
## 76	19.01	13.00
## 77	8.10	13.60
## 78	7.80	16.10
## 79	6.10	33.20
## 80	15.26	4.00
## 81	7.95	12.00
## 82	18.00	-1.50
## 83	4.60	18.30
## 84	15.00	3.00
## 85	7.50	15.80
## 86	8.00	27.10
## 87	16.80	-10.00
## 88	8.54	25.00
## 89	7.00	27.10
## 90	18.30	-8.00
## 91	7.80	12.00
## 92	16.00	-8.00
## 93	14.00	23.00

## 94	12.30	5.00
## 95	11.40	1.00
## 96	8.50	18.90
## 97	7.00	15.00
## 98	7.96	22.00
## 99	17.60	-3.50
## 100	10.00	20.00
## 101	3.50	12.20
## 102	6.70	14.70
## 103	17.00	-5.00
## 104	20.26	-4.15
## 105	6.64	11.00
## 106	1.80	-4.00
## 107	7.02	25.00
## 108	2.46	35.00
## 109	19.00	-5.00
## 110	17.86	-30.00
## 111	6.10	12.20
## 112	6.64	19.00
## 113	12.00	1.60
## 114	6.60	20.00
## 115	8.70	17.10
## 116	14.05	24.00
## 117	7.20	7.10
## 118	19.70	-11.00
## 119	7.70	21.30
## 120	6.02	5.00
## 121	2.50	12.90
## 122	19.00	5.90
## 123	6.80	5.80

### 3.32 Heat map for continuous variables


- Two dimensional **histogram**.
- It illustrates the “continuous contingency” table for continuous variables



### 3.33 Scatter plot

- The **histogram** depends on the size of the bin (pixel).
- If the pixel is small enough to contain a single observation then the heat map results in a **scatter plot**

The scatter plot is the illustration of a “contingency table” for continuous variables when the bin (pixel) is small enough to contain one single observation (consisting of a pair of values).



## Chapter 4

# Conditional Probability

### 4.1 Objective

- Conditional probability
  - Independence
  - Bayes' theorem
- 
- 

### 4.2 Joint Probability

The joint probability of two events  $A$  and  $B$  is

$$P(A, B) = P(A \cap B)$$

Let's imagine a random experiment that measures two different types of outcomes.

- height and weight of an individual:  $(h, w)$
- time and place of an electric charge:  $(p, t)$
- a throw of two dice:  $(n_1, n_2)$
- cross two traffic lights in green:  $(\bar{R}_1, \bar{R}_2)$

In many cases, we are interested in finding out whether the values of one outcome **condition** the values of the other.

---

---

### 4.3 Diagnostics

Let's consider a **diagnostic tool**

We want to find the state of a system (s):

- inadequate (yes)
- adequate (no)

with a test (t):

- positive
- negative

We test a battery to find how long it can live. We stress a cable to find if it resists carrying a certain load. We perform a PCR to see if someone is infected.

---



---

### 4.4 Diagnostics Test

Let's consider diagnosing infection with a new test.

Infection status:

- yes (infected)
- no (not infected)

Test:

- positive
- negative

---



---

### 4.5 Observations

Each individual is a random experiment with two measurements: (Infection, Test)

Subject	Infection	Test
$s_1$	yes	positive
$s_2$	no	negative
$s_3$	yes	positive
...	...	...
$s_i$	no	positive*

Subject	Infection	Test
...	...	...
...	...	...
$s_n$	yes	negative*

## 4.6 Contingency tables

- For the number of observations of each outcome

	Infection: yes	Infection: no	sum
Test: positive	18	12	30
Test: negative	30	300	330
sum	48	312	360

- For the relative frequencies, if  $N \gg 0$  we will take  $f_{i,j} = \hat{P}(x_i, y_j)$

	Infection: yes	Infection: no	sum
Test: positive	0.05	0.0333	0.0833
Test: negative	0.0833	0.833	0.9166
sum	0.133	0.866	1

## 4.7 Conditional probability

Let's think first in terms of those who are **infected**

Within those who are infected (**yes**), what is the probability of those who tested positive?

- Sensitivity (true positive rate)

$$\begin{aligned}\hat{P}(\text{positive}|\text{yes}) &= \frac{n_{\text{positive, yes}}}{n_{\text{yes}}} \\ &= \frac{\frac{n_{\text{positive, yes}}}{N}}{\frac{n_{\text{yes}}}{N}} = \frac{f_{\text{positive, yes}}}{f_{\text{yes}}}\end{aligned}$$

Therefore, in the limit, we expect to have a probability of the type

$$P(\text{positive}|\text{yes}) = \frac{P(\text{positive}, \text{yes})}{P(\text{yes})} = \frac{P(\text{positive} \cap \text{yes})}{P(\text{yes})}$$


---



---

## 4.8 Conditional probability

**Definition:** The conditional probability of an event B given an event A, denoted as  $P(A|B)$ , is

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

- you can prove that the conditional probability satisfies the axioms of probability.
  - it is the probability with the sampling space given by  $B$ :  $S_B$ .
- 
- 

## 4.9 Conditional contingency table

	Infection: Yes	Infection: No
Test: positive	$P(\text{positive}   \text{yes})$	$P(\text{positive}   \text{no})$
Test: negative	$P(\text{negative}   \text{yes})$	$P(\text{negative}   \text{no})$
sum	1	1

- True positive rate (Sensitivity): The probability of testing positive **if** having the disease  $P(\text{positive}|\text{yes})$
  - True negative rate (Specificity): The probability of testing negative **if** not having the disease  $P(\text{negative}|\text{no})$
  - False-positive rate: The probability of testing positive **if** not having the disease  $P(\text{positive}|\text{no})$
  - False-negative rate: The probability of testing negative **if** having the disease  $P(\text{negative}|\text{yes})$
- 
-



## 4.10 Example conditional contingency table

Taking the frequencies as estimates of the probabilities then

	Infection: Yes	Infection: No
Test: positive	$18/48 = 0.375$	$12/312 = 0.038$
Test: negative	$30/48 = 0.625$	$300/312 = 0.962$
sum	1	1

Our diagnostic tool has low sensitivity (0.375) but high specificity (0.962).

---



---

## 4.11 Multiplication rule

Now let's imagine the real situation where we want to compute **joint** probabilities from conditional **probabilities**

- PCRs for coronavirus were (performed)[<https://www.nejm.org/doi/full/10.1056/NEJMp2015897>] in people in the hospital who we are sure to be infected. They have a sensitivity of 70%. They have also been tested in the lab in conditions of no infection with 96% specificity
- A prevalence study in Spain showed that  $P(yes) = 0.05$ ,  $P(no) = 0.95$  before summer.

With this data, what was the probability that a randomly selected person in the population tested positive **and** was infected:  $P(yes \cap positive) = P(yes, positive)$ ?

---



---

## 4.12 Diagnostic performance

To study the performance of a new diagnostic test:

- you select specimens that are inadequate (disease: **yes**) and apply the test, trying to find its sensitivity:  $P(positive|yes)$  (0.70 for PCRs)
- you select specimens that are adequate (disease: **no**) and apply the test, trying to find its specificity:  $P(negative|no)$  (0.96 for PCRs)

---



---

	Infection: Yes	Infection: No
Test: positive	$P(\text{positive} \text{yes})=0.7$	$P(\text{positive} \text{no})=0.06$
Test: negative	$P(\text{negative} \text{yes})=0.3$	$P(\text{negative} \text{no})=0.94$
sum	1	1

From this matrix, can we obtain  $P(\text{yes}, \text{positive})$ ?

---



---

### 4.13 Multiplication rule

How do you recover the joint probability from the conditional probability?

For two events  $A$  and  $B$  we have the multiplication rule

$$P(A, B) = P(A|B)P(B)$$

that follows from the definition of the conditional probability.

---



---

### 4.14 Contingency table in terms of conditional probabilities

	Infection: Yes	Infection: No	sum
Test: positive	$P(\text{positive}   \text{yes})P(\text{yes})$	$P(\text{positive}   \text{no})P(\text{no})$	$P(\text{positive})$
Test: negative	$P(\text{negative}   \text{yes})P(\text{yes})$	$P(\text{negative}   \text{no})P(\text{no})$	$P(\text{negative})$
sum	$P(\text{yes})$	$P(\text{no})$	1

For instance the probability of testing *positive* and being infected *yes*:

- $P(\text{positive}, \text{yes}) = P(\text{positive} \cap \text{yes}) = P(\text{positive}|\text{yes})P(\text{yes})$

---



---

## 4.15 Conditional tree

---



---

## 4.16 Contingency table in terms of conditional probabilities

	Infection: yes	Infection: no	sum
Test: positive	0.035	0.057	0.092
Test: negative	0.015	0.893	0.908
sum	0.05	0.95	1

- $P(\text{positive}, \text{yes}) = 0.035$

But we also found the marginal of being positive:

- $P(\text{positive}) = 0.092$

---



---

## 4.17 Total probability rule

	Infection: Yes	Infection: No	sum
Test: positive	$P(\text{positive}   \text{yes})P(\text{yes})$	$P(\text{positive}   \text{no})P(\text{no})$	$P(\text{positive})$
Test: negative	$P(\text{negative}   \text{yes})P(\text{yes})$	$P(\text{negative}   \text{no})P(\text{no})$	$P(\text{negative})$
sum	$P(\text{yes})$	$P(\text{no})$	1

When we write the unknown marginals in terms of their conditional probabilities we call it the **total probability rule**

- $P(\text{positive}) = P(\text{positive} | \text{yes})P(\text{yes}) + P(\text{positive} | \text{no})P(\text{no})$
- $P(\text{negative}) = P(\text{negative} | \text{yes})P(\text{yes}) + P(\text{negative} | \text{no})P(\text{no})$

---



---

### 4.18 Conditional tree

**Total probability rule** for the marginal of  $B$ : In how many ways I can obtain the outcome  $B$ ?

$$P(B) = P(B|A)P(A) + P(B|A')P(A')$$

---



---

### 4.19 Finding reverse probabilities

From the conditional contingency table

	Infection: Yes	Infection: No
Test: positive	$P(\text{positive}   \text{yes})$	$P(\text{positive}   \text{no})$
Test: negative	$P(\text{negative}   \text{yes})$	$P(\text{negative}   \text{no})$
sum	1	1

How can we calculate the probability of being infected if tested positive:  $P(\text{yes}|\text{positive})$ ?

---



---

### 4.20 Recover joint probabilities

1. We recover the contingency table for joint probabilities

	Infection: Yes	Infection: No	sum
Test: positive	$P(\text{positive}   \text{yes})P(\text{yes})$	$P(\text{positive}   \text{no})P(\text{no})$	$P(\text{positive})$
Test: negative	$P(\text{negative}   \text{yes})P(\text{yes})$	$P(\text{negative}   \text{no})P(\text{no})$	$P(\text{negative})$
sum	$P(\text{yes})$	$P(\text{no})$	1

---



---

### 4.21 Reverse conditionals

2. We compute the conditional probabilities for the test:

$$P(\text{infection}|\text{test}) = \frac{P(\text{test}|\text{infection})P(\text{infection})}{P(\text{test})}$$

	Infection: Yes	Infection: No	sum
Test: positive	P(yes positive)	P(no positive)	1
Test: negative	P(yes negative)	P(no negative)	1

For instance:

$$P(\text{yes}|\text{positive}) = \frac{P(\text{positive}|\text{yes})P(\text{yes})}{P(\text{positive})}$$

since we usually don't have  $P(\text{positive})$  we use the **total probability** rule in the denominator

$$P(\text{yes}|\text{positive}) = \frac{P(\text{positive}|\text{yes})P(\text{yes})}{P(\text{positive}|\text{yes})P(\text{yes}) + P(\text{positive}|\text{no})P(\text{no})}$$

## 4.22 Baye's theorem

The expression:

$$P(\text{yes}|\text{positive}) = \frac{P(\text{positive}|\text{yes})P(\text{yes})}{P(\text{positive}|\text{yes})P(\text{yes}) + P(\text{positive}|\text{no})P(\text{no})}$$

is called the **Bayes theorem**

### Theorem

If  $E_1, E_2, \dots, E_k$  are  $k$  mutually exclusive and exhaustive events and  $B$  is any event,

$$P(E_i|B) = \frac{P(B|E_i)P(E_i)}{P(B|E_1)P(E_1) + \dots + P(B|E_k)P(E_k)}$$

It allows to reverse the conditionals:

$$P(B|A) \rightarrow P(A|B)$$

Or **design** a test  $B$  in controlled condition  $A$  and then use it to **infer** the probability of the condition when the test is positive.

---



---

### 4.23 Example: Bayes' theorem

Baye's theorem:

$$P(yes|positive) = \frac{P(positive|yes)P(yes)}{P(positive|yes)P(yes) + P(positive|no)P(no)}$$

we know:

- $P(positive|yes) = 0.70$
- $P(positive|no) = 1 - P(negative|no) = 0.06$
- the probability of infection and not infection in the population:  $P(yes) = 0.05$  and  $P(no) = 1 - P(yes) = 0.95$ .

Therefore:

$$P(yes|positive) = 0.47$$

Tests are not so good to **confirm** infections.

---



---

### 4.24 Example: Bayes' theorem

Let's now apply it to the probability of not being infected if the test is negative

$$P(no|negative) = \frac{P(negative|no)P(no)}{P(negative|no)P(no) + P(negative|yes)P(yes)}$$

Substitution of all the values gives

$$P(no|negative) = 0.98$$

Tests are good to **rule out** infections.

---



---

## 4.25 Statistical independence

In many applications, we want to know if the knowledge of one event conditions the outcome of another event.

- there are cases where we want to know if the events are not conditioned

---



---

## 4.26 Statistical independence

Consider conductors for which we measure their surface flaws and if their conduction capacity is defective

The estimated **joint probabilities** are

	flaws (F)	no flaws (F')	sum
defective (D)	0.005	0.045	0.05
no defective (D')	0.095	0.855	0.95
sum	0.1	0.9	1

where, for instance, the joint probability of  $F$  and  $D$  is

- $P(D, F) = 0.005$

The marginal probabilities are

- $P(D) = P(D, F) + P(D, F') = 0.05$
- $P(F) = P(D, F) + P(D', F) = 0.1$ .

---



---

## 4.27 Statistical independence

What is the **conditional probability** of observing a defective conductor if they have a flaw?

	F	F'
D	$P(D F) = 0.05$	$P(D F')=0.05$
D'	$P(D' F)=0.95$	$P(D' F')=0.95$
sum	1	1

The marginals and the conditional probabilities are the same!

- $P(D|F) = P(D|F') = P(D)$
- $P(D'|F) = P(D'|F') = P(D')$

The probability of observing a defective conductor **does not** depend on having observed or not a flaw.

$$P(D) = P(D|F)$$


---



---

## 4.28 Statistical independence

Two events  $A$  and  $B$  are statistically independent if

- $P(A|B) = P(A)$ ;  $A$  is independent of  $B$
- $P(B|A) = P(B)$ ;  $B$  is independent of  $A$

and by the multiplication rule, their joint probability is

- $P(A \cap B) = P(A|B)P(B) = P(A)P(B)$

the multiplication of their marginal probabilities.

---



---

## 4.29 Products of marginals products

	F	F'	sum
D	0.005	0.045	0.05
D'	0.095	0.855	0.95
sum	0.1	0.9	1

Confirm that all the entries of the matrix are the product of the marginals.

For example:

- $P(F)P(D) = P(D \cap F)$
  - $P(D')P(F') = P(D' \cap F')$
- 
-



4.30 Example

Outcomes of throwing two coins:  $S = (H, H), (H, T), (T, H), (T, T)$

	H	T	sum
H	1/4	1/4	1/2
T	1/4	1/4	1/2
sum	1/2	1/2	1

- Obtaining a head in the first coin does not condition obtaining a tail in the result of the second coin  $P(T|H) = P(T) = 1/2$
- the probability of obtaining a head and then a tail is the product of each independent outcome  $P(H, T) = P(H) * P(T) = 1/4$

---

---



## Chapter 5

# Discrete Random Variables

### 5.1 Objective

- Random variables
- Probability mass function
- Mean and variance
- Probability distribution

---

---

### 5.2 How do we assign probability values to outcomes?

---

---

### 5.3 Random variable

**Definition:**

A **random variable** is a function that assigns a real **number** to each **outcome** in the sample space of a random experiment.

- Most commonly a random variable is the value of the **measurement** of interest that is made in a random experiment.

A random variable can be:

- Discrete (nominal, ordinal)

- Continuous (interval, ratio)

---



---

## 5.4 Random variable

A **value** (or **outcome**) of a random variable is one of the possible numbers that the variable can take in a random experiment.

We write the random variable in **capitals**.

Example:

If  $X \in \{0, 1\}$ , we then say  $X$  is a random variable that can take the values 0 or 1.

**Observation** of a random variable

- An observation is the **acquisition** of the value of a random variable in a random experiment

Example:

1 0 0 1 0 **1** 0 1 1

The number in bold is an observation of  $X$

---



---

## 5.5 Events of observing a random variable

- $X = 1$  is the **event** of observing the random variable  $X$  with value 1
- $X = 2$  is the **event** of observing the random variable  $X$  with value 2

...

**In general:**

- $X = x$  is the **event** of observing the random variable  $X$  with value  $x$  (little  $x$ )
- Any two values of a random variable define two **mutually exclusive** events.

---



---

## 5.6 Probability of random variables


We are interested in assigning probabilities to the values of a random variable.

We have already done this for the dice:  $X \in \{1, 2, 3, 4, 5, 6\}$  (classical interpretation of probability)

$X$	Probability
1	$P(X = 1) = 1/6$
2	$P(X = 2) = 1/6$
3	$P(X = 3) = 1/6$
4	$P(X = 4) = 1/6$
5	$P(X = 5) = 1/6$
6	$P(X = 6) = 1/6$

## 5.7 Probability functions

- We can write the probability table
- plot it



- or write as the function

$$f(x) = P(X = x) = 1/6$$



---



---

## 5.8 Probability functions

We can **create** any type of probability function if we respect the probability rules:



## 5.9 Probability functions

For a discrete random variable  $X \in \{x_1, x_2, \dots, x_M\}$ , a **probability mass function**

is always positive

- $f(x_i) \geq 0$

is used to compute probabilities

- $f(x_i) = P(X = x_i)$

and its sum over all the values of the variable is 1:

- $\sum_{i=1}^M f(x_i) = 1$

---



---

## 5.10 Probability functions

- Note that the definition of  $X$  and its probability mass function is general **without reference** to any experiment. The functions live in the model (abstract) space.
- $X$  and  $f(x)$  are abstract objects that may or may not map to an experiment
- We have the freedom to construct them as we want as long as we respect their definition.
- They have some **properties** that are derived exclusively from their definition.

---



---

## 5.11 Example: Probability mass function

Consider the following random variable  $X$  over the outcomes

outcome	$X$
$a$	0
$b$	0
$c$	1.5
$d$	1.5
$e$	2
$f$	3

If each outcome is equally probable then what is the probability mass function of  $x$ ?

---



---

### 5.12 Probability table for equally likely outcomes

outcome	Probability(outcome)
$a$	$1/6$
$b$	$1/6$
$c$	$1/6$
$d$	$1/6$
$e$	$1/6$
$f$	$1/6$

### 5.13 Probability table for $X$

$X$	$f(x) = P(X = x)$
0	$P(X = 0) = 2/6$
1.5	$P(X = 1.5) = 2/6$
2	$P(X = 2) = 1/6$
3	$P(X = 3) = 1/6$

We can compute, for instance, the following probabilities for events on the values of  $X$

- $P(X > 3)$
- $P(X = 0 \cup X = 2)$
- $P(X \leq 2)$

### 5.14 Example

**Probability model:**

Consider the following experiment: In one urn put 8 balls and:

- mark 1 ball with  $-2$
- mark 2 balls with  $-1$
- mark 2 balls with  $0$



- mark 2 balls with 1
- mark 1 ball with 2

**experiment:** Take one ball and read the number.

$X$	$P(X = x)$
-2	$1/8 = 0.125$
-1	$2/8 = 0.25$
0	$2/8 = 0.25$
1	$2/8 = 0.25$
2	$1/8 = 0.125$

---



---

## 5.15 Example

Consider another experiment where we do not know what is in the previous urn. We draw a ball 30 times, write its number and put it back in the urn.

- we do not know what the primary events with equal probabilities are.
- we then **estimate** the probability mass function from the relative frequencies observed for a random variable

$X$	$f_i$
-2	0.132
-1	0.262
0	0.240
1	0.248
2	0.118

---



---

## 5.16 Probabilities and frequencies

For computing the relative frequencies  $f_i$  you have to

- **repeat** the experiment  $N$  times (you have to put the ball back in the urn each time) and at the end compute

$$f_i = n_i/N$$

We are assuming that:


$$\lim_{N \rightarrow \infty} f_i = f(x_i) = P(X = x_i)$$


---



---

### 5.17 Probabilities and relative frequencies



- In this example we **know** the probability **model**  $f(x) = P(X = x)$  by design.
- We never observe  $f(x)$
- We can use relative frequencies to estimate the probabilities

$$f_i = \hat{f}(x_i) = \hat{P}(X = x_i)$$

( $f_i$  depends on  $N$ )

---



---

### 5.18 Mean and Variance

The probability mass functions  $f(x)$  have two main properties

- its center
- its spread

We can ask,

- around which values of  $X$  the probability concentrated?
- How dispersed are the values of  $X$  in relation to their probabilities?

---



---

## 5.19 Mean and Variance

---



---

## 5.20 Mean

Remember that the **average** in terms of the relative frequencies of the values of  $x_i$  (categorical ordered outcomes) can be written as

$$\bar{x} = \sum_{i=1}^M x_i \frac{n_i}{N} = \sum_{i=1}^M x_i f_i$$

### Definition

The **mean** ( $\mu$ ) or expected value of a discrete random variable  $X$ ,  $E(X)$ , with mass function  $f(x)$  is given by

$$\mu = E(X) = \sum_{i=1}^M x_i f(x_i)$$

It is the center of gravity of the **probabilities**: The point where probability loadings on a road are balanced

---




---

## 5.21 Example: Mean

What is the mean of  $X$  if its probability mass function  $f(x)$  is given by

$$P(X = 0) = 1/16 \quad P(X = 1) = 4/16 \quad P(X = 2) = 6/16 \quad P(X = 3) = 4/16 \\ P(X = 4) = 1/16$$



$$\mu = E(X) = \sum_{i=1}^m x_i f(x_i)$$

$$E(X) = 0 * 1/16 + 1 * 4/16 + 2 * 6/16 + 3 * 4/16 + 4 * 1/16 = 2$$


---



---

## 5.22 Mean and Average

- The mean  $\mu$  is the centre of gravity of the probability mass function **it does not change**

For instance for

$X$	$P(X = x)$
-2	$1/8 = 0.125$
-1	$2/8 = 0.25$
0	$2/8 = 0.25$
1	$2/8 = 0.25$
2	$1/8 = 0.125$

- The average  $\bar{x}$  is the centre of gravity of the observations (relative frequencies) it **changes** with different data

For instance for

$X$	$f_i$
-2	0.132
-1	0.262
0	0.240
1	0.248
2	0.118

## 5.23 Variance

In similar terms we define the mean squared distance from the mean:

### Definition

The variance, written as  $\sigma^2$  or  $V(X)$ , of a discrete random variable  $X$  with mass function  $f(x)$  is given by

$$\sigma^2 = V(X) = \sum_{i=1}^M (x_i - \mu)^2 f(x_i)$$

- $\sigma = \sqrt{V(X)}$  is called the **standard deviation** of the random variable
- Think of it as the moment of inertia of probabilities about the mean.

## 5.24 Example: Variance

What is the variance of  $X$  if its probability mass function  $f(x)$  is given by

$$P(X = 0) = 1/16 \quad P(X = 1) = 4/16 \quad P(X = 2) = 6/16 \quad P(X = 3) = 4/16 \\ P(X = 4) = 1/16$$

$$\sigma^2 = V(X) = \sum_{i=1}^m (x_i - \mu)^2 f(x_i)$$

$$V(X) = (0-2)^2 * 1/16 + (1-2)^2 * 4/16 + (2-2)^2 * 6/16 + (3-2)^2 * 4/16 + (4-2)^2 * 1/16 = 1$$

$$V(X) = \sigma^2 = 1$$

$$\sigma = 1$$


---



---

## 5.25 Functions of $X$

### Definition

For any function  $h$  of a random variable  $X$ , with mass function  $f(x)$ , its expected value is given by

$$E[h(X)] = \sum_{i=1}^M h(x_i) f(x_i)$$

This is an important definition that allows us to prove three important properties of the median and variance:

- The mean of a linear function is the linear function of the mean:

$$E(a \times X + b) = a \times E(X) + b$$

for  $a$  and  $b$  scalars (numbers).

- The variance of a linear function of  $X$  is:

$$V(a \times X + b) = a^2 \times V(X)$$

- The variance **about the origin** is the variance **about the mean** plus the mean squared:

$$E(X^2) = V(X) + E(X)^2$$


---



---

## 5.26 Example: Variance about the origin

What is the variance  $X$  about the origin,  $E(X^2)$ , if its probability mass function  $f(x)$  is given by

$$P(X = 0) = 1/16 \quad P(X = 1) = 4/16 \quad P(X = 2) = 6/16 \quad P(X = 3) = 4/16 \\ P(X = 4) = 1/16$$

$$E(X^2) = \sum_{i=1}^m x_i^2 f(x_i)$$

$$E(X^2) = (0)^2 * 1/16 + (1)^2 * 4/16 + (2)^2 * 6/16 + (3)^2 * 4/16 + (4)^2 * 1/16 = 5$$

We can also verify:

$$E(X^2) = V(X) + E(X)^2$$

$$5 = 1 + 2^2$$

---



---

## 5.27 Probability distribution

**Definition:**

The **probability distribution** function is defined as

$$F(x) = P(X \leq x) = \sum_{x_i \leq x} f(x_i)$$

That is the accumulated probability up to a given value  $x$

$F(x)$  satisfies:

- $0 \leq F(x) \leq 1$
- If  $x \leq y$ , then  $F(x) \leq F(y)$

---



---

## 5.28 Example: Probability distribution

For the probability mass function:

$$f(0) = P(X = 0) = 1/16 \quad f(1) = P(X = 1) = 4/16 \quad f(2) = P(X = 2) = 6/16 \\ f(3) = P(X = 3) = 4/16 \quad f(4) = P(X = 4) = 1/16$$

The probability distribution is:

$$F(x) = \begin{cases} 1/16, & \text{if } x < 1 \\ 5/16, & 1 \leq x < 2 \\ 11/16, & 2 \leq x < 3 \\ 15/16, & 3 \leq x < 4 \\ 16/16, & x \leq 5 \end{cases}$$


For  $X \in \mathbb{Z}$

---



---

### 5.29 Probability distribution




---



---

### 5.30 Probability function and Probability distribution

Compute the mass probability function of the following probability distribution:

$$F(0) = 1/16, F(1) = 5/16, F(2) = 11/16, F(3) = 15/16, F(4) = 16/16,$$



Let's work backward.

$$\begin{aligned} f(0) &= F(0) = 1/16 & f(1) &= F(1) - f(0) = 5/32 - 1/32 = 4/16 & f(2) &= F(2) - \\ & & f(1) - f(0) &= F(2) - F(1) = 6/16 & f(3) &= F(3) - f(2) - f(1) - f(0) = F(3) - \\ & & & & F(2) &= 4/16 & f(4) &= F(4) - F(3) = 1/16 \end{aligned}$$



## 5.31 Probability function and Probability distribution

The Probability distribution is another way to specify the probability of a random variable

$$f(x_i) = F(x_i) - F(x_{i-1})$$

with

$$f(x_1) = F(x_1)$$

for  $X$  taking values in  $x_1 \leq x_2 \leq \dots \leq x_n$



## 5.32 Quantiles

We define the **q-quantile** as the value  $x_p$  **under** which we have accumulated  $q \cdot 100\%$  of the probability

$$q = \sum_{i=1}^p f(x_i) = F(x_p)$$

- The **median** is value  $x_m$  such that  $q = 0.5$

$$F(x_m) = 0.5$$

- The 0.05-quantile is the value  $x_r$  such that  $q = 0.05$

$$F(x_r) = 0.05$$

- The 0.25-quantile is **first quartile** the value  $x_s$  such that  $q = 0.25$

$$F(x_s) = 0.25$$


---



---

### 5.33 Summary

quantity names	model (unobserved)	data (observed)
probability mass function // relative frequency	$f(x_i) = P(X = x_i)$	$f_i = \frac{n_i}{N}$
probability distribution // cumulative relative frequency	$F(x_i) = P(X \leq x_i)$	$F_i = \sum_{k \leq i} f_k$
mean // average	$\mu = E(X) = \sum_{i=1}^M x_i f(x_i)$	$\bar{x} = \sum_{j=1}^N x_j / N$
variance // sample variance	$\sigma^2 = V(X) = \sum_{i=1}^M (x_i - \mu)^2 f(x_i)$	$s^2 = \sum_{j=1}^N (x_j - \bar{x})^2 / (N - 1)$
standard deviation // sample sd	$\sigma = \sqrt{V(X)}$	$s$
variance about the origin // 2nd sample moment	$E(X^2) = \sum_{i=1}^M x_i^2 f(x_i)$	$m_2 = \sum_{j=1}^N x_j^2 / n$

Note that:

- $i = 1 \dots M$  is an **outcome** of the random variable  $X$ .
- $j = 1 \dots N$  is an **observation** of the random variable  $X$ .

Properties:

- $\sum_{i=1}^N f(x_i) = 1$
- $f(x_i) = F(x_i) - F(x_{i-1})$
- $E(a \times X + b) = a \times E(X) + b$ ; for  $a$  and  $b$  scalars.
- $V(a \times X + b) = a^2 \times V(X)$
- $E(X^2) = V(X) + E(X)^2$

## Chapter 6

# Continuous Random Variables

### 6.1 Objective

- Probability density function
  - Mean and variance
  - Probability distribution
- 
- 

### 6.2 Continuous random variable

What happens with continuous random variables?

Let's reconsider the convexity angle of misophonia patients (Section 2.21).

- We redefined the outcomes as little regular intervals (bins) and computed the relative frequency for each of them as we did in the discrete case.

```
##          outcome ni          fi
## 1 [-1.02,3.46]  8 0.06504065
## 2  (3.46,7.92] 51 0.41463415
## 3  (7.92,12.4] 26 0.21138211
## 4  (12.4,16.8] 20 0.16260163
## 5  (16.8,21.3] 18 0.14634146
```

---

---

### 6.3 Continuous random variable

Let's consider again that their relative frequencies are the probabilities when  $N \rightarrow \infty$

$$f_i = \frac{n_i}{N} \rightarrow f(x_i) = P(X = x_i)$$

The probability depends now on the length of the bins  $\Delta x$ . If we make the bins smaller and smaller then the frequencies get smaller and therefore

$P(X = x_i) \rightarrow 0$  when  $\Delta x \rightarrow 0$ , because  $n_i \rightarrow 0$

##	outcome	ni	fi
## 1	[-1.02,0.115]	2	0.01626016
## 2	(0.115,1.23]	0	0.00000000
## 3	(1.23,2.34]	3	0.02439024
## 4	(2.34,3.46]	3	0.02439024
## 5	(3.46,4.58]	2	0.01626016
## 6	(4.58,5.69]	4	0.03252033
## 7	(5.69,6.8]	11	0.08943089
## 8	(6.8,7.92]	34	0.27642276
## 9	(7.92,9.04]	12	0.09756098
## 10	(9.04,10.2]	4	0.03252033
## 11	(10.2,11.3]	3	0.02439024
## 12	(11.3,12.4]	7	0.05691057
## 13	(12.4,13.5]	2	0.01626016
## 14	(13.5,14.6]	6	0.04878049
## 15	(14.6,15.7]	4	0.03252033
## 16	(15.7,16.8]	8	0.06504065
## 17	(16.8,18]	4	0.03252033
## 18	(18,19.1]	9	0.07317073
## 19	(19.1,20.2]	3	0.02439024
## 20	(20.2,21.3]	2	0.01626016

---



---

### 6.4 Continuous random variable

We define a quantity at a point  $x$  that is the amount of probability per unit distance that we would find in an **infinitesimal** bin  $dx$  at  $x$

$$f(x) = \frac{P(x \leq X \leq x + dx)}{dx}$$

$f(x)$  is called the probability **density** function.

Therefore, the probability of observing  $x$  between  $x$  and  $x + dx$  is given by

$$P(x \leq X \leq x + dx) = f(x)dx$$


---



---

## 6.5 Continuous random variable

### Definition

For a continuous random variable  $X$ , a **probability density** function is such that

The function is positive:

- $f(x) \geq 0$

The probability of observing a value within an interval is the **area under the curve**:

- $P(a \leq X \leq b) = \int_a^b f(x)dx$

The probability of observing **any** value is 1:

- $\int_{-\infty}^{\infty} f(x)dx = 1$
- 
- 

## 6.6 Continuous random variable

- The probability density function is a step forward in the abstraction of probabilities: we add the continuous limit ( $dx \rightarrow 0$ ).
  - All the properties of probabilities are translated in terms of densities ( $\sum \rightarrow \int$ ).
  - Assignment of probabilities to a random variable can be done with equiprobability (classical) arguments.
  - Densities are mathematical quantities some will map to experiments some will not. *Which density will map best to my experiment?*
- 
-


## 6.7 Total area under the curve

Example: take the **probability density** that may describe the random variable that measures where a raindrop falls in a rain gutter of length 100cm.

$$f(x) = \begin{cases} \frac{1}{100}, & \text{if } x \in (0, 100) \\ 0, & \text{otherwise} \end{cases}$$

Then the probability of **any** observation is the total **area under the curve**


$$P(-\infty \leq X \leq \infty) = \int_{-\infty}^{\infty} f(x)dx = 100 * 0.01 = 1$$



## 6.8 Area under the curve

The probability of observing  $x$  in an interval is the **area under the curve** within the interval

- $P(20 \leq X \leq 60) = \int_{20}^{60} f(x)dx = (60 - 20) * 0.01 = 0.4$




---




---

## 6.9 Area under the curve

In general  $f(x)$  should satisfy:

- $0 \leq P(a \leq X \leq b) = \int_a^b f(x)dx \leq 1$




## 6.10 Probability distribution

The probability accumulated up to  $b$  is defined by the probability distribution  $F$


- $F(b) = P(X \leq b) = \int_{-\infty}^b f(x)dx$





The probability accumulated up to  $a$  is


- $F(a) = P(X \leq a)$



## 6.11 Probability distribution

The probability between  $a$  and  $b$  is defined by the probability distribution  $F$

- $P(a \leq X \leq b) = \int_a^b f(x)dx = F(b) - F(a)$



## 6.12 Probability distribution

The probability distribution of a continuous random variable is defined as

$$F(a) = P(X \leq a) = \int_{-\infty}^a f(x)dx$$

with the properties that:

It is between 0 and 1:

- $F(-\infty) = 0$  and  $F(\infty) = 1$

It always increases:

- if  $a \leq b$  then  $F(a) \leq F(b)$

It can be used to compute probabilities:

- $P(a \leq X \leq b) = F(b) - F(a)$

It recovers the probability density:

- $f(x) = \frac{dF(x)}{dx}$

We use **probability distributions** to **compute probabilities** of a random variable with intervals




## 6.13 Probability distribution

For the uniform density function:

$$f(x) = \begin{cases} \frac{1}{100}, & \text{if } x \in (0, 100) \\ 0, & \text{otherwise} \end{cases}$$


The probability distribution is

$$F(a) = \begin{cases} 0, & a \leq 0 \\ \frac{a}{100}, & \text{if } a \in (0, 100) \\ 1, & 10 \leq a \end{cases}$$




## 6.14 Probability graphics

The probability  $P(20 < X < 60)$  is the *area* under the **density** curve



### 6.15 Probability graphics

The probability  $P(20 < X < 60)$  is the *difference* in **distribution** values




---



---

## 6.16 Mean

As in the discrete case, the **mean** measures the center of the distribution

### Definition

Suppose  $X$  is a continuous random variable with probability **density** function  $f(x)$ . The mean or expected value of  $X$ , denoted as  $\mu$  or  $E(X)$ , is

$$\mu = E(X) = \int_{-\infty}^{\infty} x f(x) dx$$

It is the continuous version of the center of mass.

---




---

## 6.17 Mean

$$f(x) = \begin{cases} \frac{1}{100}, & \text{if } x \in (0, 100) \\ 0, & \text{otherwise} \end{cases}$$

$$E(X) = 50$$



## 6.18 Variance

As in the discrete case, the variance measures the dispersion about the mean

### Definition

Suppose  $X$  is a continuous random variable with probability density function  $f(x)$ . The variance of  $X$ , denoted as  $\sigma^2$  or  $V(X)$ , is

$$\sigma^2 = V(X) = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$$

## 6.19 Functions of $X$

### Definition

For any function  $h$  of a random variable  $X$ , with mass function  $f(x)$ , its expected value is given by

$$E[h(X)] = \int_{-\infty}^{\infty} h(x)f(x)dx$$

And we have the same properties as in the discrete case

- The mean of a linear function is the linear function of the mean:

$$E(a \times X + b) = a \times E(X) + b$$


for  $a$  and  $b$  scalars.

- The variance of a linear function of  $X$  is:

$$V(a \times X + b) = a^2 \times V(X)$$

- The variance about the origin is the variance about the mean plus the mean squared:

$$E(X^2) = V(X) + E(X)^2$$




## 6.20 Example

- for the probability density

$$f(x) = \begin{cases} \frac{1}{100}, & \text{if } x \in (0, 100) \\ 0, & \text{otherwise} \end{cases}$$

- compute the mean
- compute variance using  $E(X^2) = V(X) + E(X)^2$
- compute  $P(\mu - \sigma \leq X \leq \mu + \sigma)$
- What are the first and third quartiles?





## Chapter 7

# Discrete Probability Models

### 7.1 Objective

Discrete probability models:

- Uniform and Bernoulli probability functions
  - Binomial and negative binomial probability functions
- 
- 

### 7.2 Probability mass function

A probability mass function of a **discrete random variable**  $X$  with possible values  $x_1, x_2, \dots, x_M$  is **any function** such that

Positive:

- $f(x_i) \geq 0$

*Allow us to compute probabilities:*

- $f(x_i) = P(X = x_i)$

The probability of any outcome is 1

- $\sum_{i=1}^M f(x_i) = 1$

**Properties:**

Central tendency:

- $E(X) = \sum_{i=1}^M x_i f(x_i)$

Dispersion:

- $V(X) = \sum_{i=1}^M (x_i - \mu)^2 f(x_i)$

They are abstract objects with general properties that may or may not **describe** a natural or engineered process.

### 7.3 Probability model

A **probability model** is a probability mass function that may represent the probabilities of a random experiment.

**Examples:**

- $f(x) = P(X = x) = 1/6$  represents the probability of the outcomes of **one** throw of a dice.
- The probability mass function

$X$	$f(x)$
-2	1/8
-1	2/8
0	2/8
1	2/8
2	1/8

Represents the probability of drawing **one** ball from an urn where there are two balls per label:  $-1, 0, 1$  and one ball per label:  $-2, 2$ .

### 7.4 Parametric models

When we perform a random experiment and **do not** know the probabilities of the outcomes:

- We can always formulate the model given by the relative frequencies:  
 $\hat{P}(X = x_i) = f_i$  ( where  $i = 1 \dots M$  ).

We need to find  $M$  numbers each depending on  $N$ .

In many cases:

- We can formulate probability functions  $f(x)$  that depend on **very few** numbers only.

**Example:**

A random experiment with  $M$  equally likely outcomes has a probability mass function:

$$f(x) = P(X = x) = 1/M$$

We only need to know  $M$ .

The numbers we **need to know** to fully determine a probability function are called **parameters**.

---



---

## 7.5 Uniform distribution (one parameter)

**Definition** A random variable  $X$  with outcomes  $\{1, \dots, M\}$  has a discrete **uniform distribution** if all its  $M$  outcomes have the same probability

$$f(x) = \frac{1}{M}$$

With mean and variance:

$$E(X) = \frac{M+1}{2}$$

$$V(X) = \frac{M^2-1}{12}$$

Note:  $E(X)$  and  $V(X)$  are also **parameters**. If we know any of them then we can fully determine the distribution.


$$f(x) = \frac{1}{2E(X) - 1}$$

---



---

## 7.6 Uniform distribution



## 7.7 Uniform distribution (two parameters)

Let's introduce a new uniform probability model with **two parameters**: The minimum and maximum outcomes.

If the random variable takes values in  $\{a, a+1, \dots, b\}$ , where  $a$  and  $b$  are integers and all the outcomes are equally probable then

$$f(x) = \frac{1}{b - a + 1}$$

as  $M = b - a + 1$ .

- We then say that  $X$  distributes uniformly between  $a$  and  $b$  and write

$$X \rightarrow Unif(a, b)$$

## 7.8 Uniform distribution (two parameters)

### Example:

What is the probability of observing a child of a particular age in a primary school (if all classes have the same amount of children)?


From the experiment we know:  $a = 6$  and  $b = 11$  then

$$X \rightarrow Unif(a = 6, b = 11)$$

that is

$$f(x) = \frac{1}{6}$$

for  $x \in \{6, 7, 8, 9, 10, 11\}$ , and 0 otherwise



## 7.9 Uniform distribution

The probability model of a random variable  $X$

$$f(x) = \frac{1}{b-a+1}$$

for  $x \in \{a, a+1, \dots, b\}$

has mean and variance:

- $E(X) = \frac{b+a}{2}$
- $V(X) = \frac{(b-a+1)^2-1}{12}$

(Change variables  $X = Y + a - 1$ ,  $y \in \{1, \dots, M\}$ )

We can either specify  $a$  and  $b$  or  $E(X)$  and  $V(X)$ .

In our example:


- $E(X) = (11 + 6)/2 = 8.5$
- $V(X) = (6^2 - 1)/12 = 2.916667$

---



---

## 7.10 Uniform distribution (two-parameter)




---



---

## 7.11 Parameters and Models

- A **model** is a particular function  $f(x)$  that **describes** our experiment
- If the model is a **known** function that depends on a few parameters then changing the value of the parameters we produce a **family of models**
- Knowledge of  $f(x)$  is reduced to the knowledge of the value of the parameters
- Ideally, the model and the parameters are **interpretable**

*Example:*

**Model:** The data of our experiment is produced by a random process in which each age has the **same probability** of being observed.

**Parameters:**  $a$  is the minimum age,  $E(X)$  is the expected age ... they are **physical properties** of the experiment.




## 7.12 Parameters and Models

**Example:**

A **family** of models obtained from two-parameter uniform distributions changing the **variances** and keeping a constant mean ( $E(X) = 8.5$ ). It results on **changing** both **minimum** and **maximum** outcomes.

- Note: Only one model makes sense for our experiment (only one model can represent the ages of children in a school).



- We can think of **families** that change only the **mean**, only the **minimum**, or only the **maximum**

---



---

### 7.13 Bernoulli trial

Let's try to advance from the equal probability case and suppose a model with two outcomes ( $A$  and  $B$ ) that have **unequal** probabilities

**Examples:**

- Writing down the sex of a patient who goes into an emergency room of a hospital ( $A$  : *male* and  $B$  : *female*).
- Recording whether a manufactured machine is defective or not ( $A$  : *defective* and  $B$  : *fine*).
- Hitting a target ( $A$  : *success* and  $B$  : *failure*).
- Transmitting one pixel correctly ( $A$  : *yes* and  $B$  : *no*).

In these examples, the probability of outcome  $A$  is usually **unknown**.

---



---



## 7.14 Bernoulli trial

We will introduce the probability of an outcome ( $A$ ) as the **parameter** of the model:

- outcome A (success): has probability  $p$  (parameter)
- outcome B (failure): has a probability  $1 - p$

Or write, the probability mass function of  $K$  taking values  $\{0, 1\}$  for  $A$  and  $B$

$$f(k) = \begin{cases} 1 - p, & k = 0 \text{ (event } B) \\ p, & k = 1 \text{ (event } A) \end{cases}$$

or more shortly

$$f(k; p) = p^k(1 - p)^{1-k}$$

for  $k = (0, 1)$

We only need to know  $p$ .



## 7.15 Bernoulli trial

A Bernoulli variable  $K$  with outcomes  $\{0, 1\}$  has a probability mass function


$$f(k; p) = p^k(1 - p)^{1-k}$$

With mean and variance:


- $E(K) = p$
- $V(K) = (1 - p)p$

Note:

- The probability of the outcome  $A$  is the parameter  $p$  which is the same as  $f(1) = P(X = 1)$ .
- As  $p$  is usually **unknown** we typically estimated it by the relative frequency (more on this in the inference sections):  $\hat{p} = f_A = \frac{n_A}{N}$



## 7.16 Bernoulli trial




---



---

## 7.17 Binomial distribution

When we are interested in learning about a particular Bernoulli trial

- We repeat the Bernoulli trial  $N$  times and count how many times we obtained  $A$  ( $n_A$ ).
- We define a random variable  $X = n_A$  taking values  $x \in 0, 1, \dots, N$

We now ask for the probability of observing  $x$  events of type  $A$  in the repetition of  $n$  independent Bernoulli trials, when the probability of observing  $A$  is  $p$ .

$$P(X = x) = f(x) = ?$$

---



---

## 7.18 Examples: Binomial distribution

- Writing down the sex of  $n = 10$  patients who go into an emergency room of a hospital. What is the probability that  $x = 6$  patients are men when  $p = 0.9$ ?
- Trying  $n = 5$  times to hit a target ( $A : \text{success}$  and  $B : \text{failure}$ ). What is the probability that I hit the target  $x = 5$  times when I usually hit it 25% of the times ( $p = 0.25$ )?
- Transmitting  $n = 100$  pixels correctly ( $A : \text{yes}$  and  $B : \text{no}$ ). What is the probability that  $x = 2$  pixels are errors, when the probability of error is  $p = 0.1$ ?

---



---

## 7.19 Binomial distribution

What is the probability of observing  $X = 4$  errors when transmitting 4 pixels, if the probability of an error is  $p$ ?

Consider 4 random variables:  $K_1, K_2, K_3$  and  $K_4$  that record whether an error has been made in the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> pixel.

Then

- $k_i$  takes values  $\{\text{correct} : 0; \text{error} : 1\}$
- $X = \sum_{i=1}^4 K_i$  takes values  $\{0, 1, 2, 3, 4\}$

Then the probability of observing 4 errors is:

- $P(X = 4) = P(1, 1, 1, 1) = p * p * p * p = p^4$  because  $K_i$  are independent.

The probability of 0 errors is:

- $P(X = 0) = P(0, 0, 0, 0) = (1 - p)(1 - p)(1 - p)(1 - p) = (1 - p)^4$

The probability of 3 errors is:

$$P(X = 3) = P(0, 1, 1, 1) + P(1, 0, 1, 1) + P(1, 1, 0, 1) + P(1, 1, 1, 0) = 4p^3(1 - p)^1$$

---



---

## 7.20 Binomial distribution

Therefore the probability of  $x$  errors is

$$f(x) = \begin{cases} 1 * p^0(1-p)^4, & x = 0 \\ 4 * p^1(1-p)^3, & x = 1 \\ 6 * p^2(1-p)^2, & x = 2 \\ 4 * p^3(1-p)^1, & x = 3 \\ 1 * p^4(1-p)^0, & x = 4 \end{cases}$$

or more shortly

$$f(x) = \binom{4}{x} p^x (1-p)^{4-x}$$

for  $x = 0, 1, 2, 3, 4$

where  $\binom{4}{x}$  is the number of possible outcomes (transmissions of 4 pixels) with  $x$  errors.

---



---

## 7.21 Binomial distribution: Definition

The binomial probability function is the probability mass function of observing  $x$  outcomes of type  $A$  in  $n$  independent Bernoulli trials, where  $A$  has the same probability  $p$  in each trial.

The function is given by

$$f(x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \dots, n$$

$\binom{n}{x} = \frac{n!}{x!(n-x)!}$  is called **the binomial coefficient** and gives the number of ways one can obtain  $x$  events of type  $A$  in a set of  $n$ .

When a variable  $X$  has a binomial probability function we say it distributes binomially and write

$$X \rightarrow \text{Bin}(n, p)$$

where  $n$  and  $p$  are parameters.

---



---

## 7.22 Binomial distribution: Mean and Variance

The mean and variance of  $X \hookrightarrow \text{Bin}(n, p)$  are

- $E(X) = np$
- $V(X) = np(1 - p)$
- Since  $X$  is the sum of  $n$  independent Bernoulli variables

$$E(X) = E(\sum_{i=1}^n K_i) = np$$

and

$$V(X) = V(\sum_{i=1}^n K_i) = n(1 - p)p$$

Example:

- The expected value for the number of errors in the transmission of 4 pixels is  $np = 4 * 0.1 = 0.4$  when the probability of an error is 0.1.
- The variance is  $n(1 - p)p = 0.36$

**Remember:** We can specify either the parameters  $n$  and  $p$ , or the parameters  $E(X)$  and  $V(X)$

---



---

## 7.23 Example 1

Now let's answer:

- What is the probability of observing 4 errors when transmitting 4 pixels, if the probability of an error is 0.1?

Since we are repeating a Bernoulli trial  $n = 4$  times and counting the number of events of type  $A$  (errors), when  $P(A) = p = 0.1$  then

$$X \rightarrow \text{Bin}(n = 4, p = 0.1)$$

That is

$$f(x) = \binom{4}{x} 0.1^x (1 - 0.1)^{4-x}$$

---



---

### 7.24 Example 1

- We want to compute:

$$P(X = 4) = f(4) = \binom{4}{4} 0.1^4 0.9^0 = 10^{-4}$$

In R `dbinom(4,4,0.1)`

- We can also compute:

$$P(X = 2) = \binom{4}{2} 0.1^2 0.9^2 = 0.0486$$

In R `dbinom(2,4,0.1)`

---



---

### 7.25 Example 2

- What is the probability of observing at most 8 voters of the ruling party in an election poll of size 10, if the probability of a positive vote is 0.9

For this case

$$X \rightarrow \text{Bin}(n = 10, p = 0.9)$$

That is

$$f(x) = \binom{10}{x} 0.9^x (0.1)^{4-x}$$

We want to compute:  $P(X \leq 8) = F(8) = \sum_{i=1..8} f(x_i) = 0.2639011$


in R `pbinom(8,10, 0.9)`

---



---

## 7.26 Binomial distribution



## 7.27 Negative binomial distribution

Now let us imagine that we are interested in counting the well-transmitted pixels before a **given number** of errors occur. Say we can **tolerate**  $r$  errors in transmission.

- Experiment: Suppose performing Bernoulli trials until we observe the outcome  $A$  appears  $r$  times.
- Random variable: We count the number of events  $B$
- Example: What is the probability of observing  $y$  well-transmitted ( $B$ ) pixels before  $r$  errors ( $A$ )?

## 7.28 Negative binomial distribution

Let's first find the probability of one particular transmission with  $y$  number of correct pixels ( $B$ ) and  $r$  number of errors ( $A$ ).

$(0, 0, 1, \dots, 0, 1, \dots, 0, 1)$  (there are  $y$  zeros, and  $r$  ones)

We observe  $y$  correct pixels in a total of  $y + r$  trials.

Then

- $P(0, 0, 1, \dots, 0, 1, \dots, 0, 1) = p^r(1 - p)^y$  (Remember:  $p$  is the probability of error)

How many transmissions can have  $y$  correct pixels before  $r$  errors?

Note:

- The last bit is fixed (marks the end of transmission)
- The total number of transmissions with  $y$  number of correct pixels ( $B$ ) that we can obtain in  $y + r - 1$  trials is:  $\binom{y+r-1}{y}$

---



---

## 7.29 Negative binomial distribution

Therefore, the probability of observing  $y$  events of type  $B$  before  $r$  events of type  $A$  (with probability  $p$ ) is

$$P(Y = y) = f(y) = \binom{y+r-1}{y} p^r (1-p)^y$$

for  $y = 0, 1, \dots$

We then say that  $Y$  follows a negative binomial distribution and we write

$$Y \rightarrow NB(r, p)$$

where  $r$  and  $p$  are parameters representing the tolerance and the probability of a single error.

---



---



### 7.30 Mean and Variance

A random variable with  $Y \rightarrow NB(r, p)$  has

- mean:  $E(Y) = r \frac{1-p}{p}$
- variance:  $V(Y) = r \frac{1-p}{p^2}$

---



---

### 7.31 Geometric distribution

We call **geometric distribution** to the negative binomial distribution with  $r = 1$

The probability of observing  $B$  events before observing the **first** event of type  $A$  is

$$P(Y = y) = f(y) = p(1 - p)^y$$

$$Y \rightarrow Geom(p)$$

with mean

- mean:  $E(Y) = \frac{1-p}{p}$
- variance:  $V(Y) = \frac{1-p}{p^2}$

---



---

### 7.32 Example

- A website has three servers.
- One server operates at a time and only when a request fails another server is used.
- If the probability of failure for a request is known to be  $p = 0.0005$  then
- what is the expected number of successful requests before the three computers fail?

---



---

### 7.33 Example

Since we are repeating a Bernoulli trial until  $r = 3$  events of type  $A$  (failure) are observed (each with  $P(A) = p = 0.0005$ ) and are counting the number of events of type  $B$  (successful requests) then

$$Y \rightarrow NB(r = 3, p = 0.0005)$$

Therefore, the expected number of requests before the system fails is:

$$E(Y) = r \frac{1-p}{p} = 3 \frac{1-0.0005}{0.0005} = 5997$$

- Note that there are actually 6000 trials

---



---

### 7.34 Example

What is the probability of dealing with at most 5 successful requests before the system fails?

Recall the cumulative function distribution  $F(y) = P(Y \leq 5)$

$$\begin{aligned} F(5) &= P(Y \leq 5) = \sum_{y=0}^5 f(y) \\ &= \sum_{y=0}^5 \binom{y+2}{y} 0.0005^3 0.9995^y \\ &= \binom{2}{0} 0.0005^3 0.9995^0 + \binom{3}{1} 0.0005^3 0.9995^1 \\ &\quad + \binom{4}{2} 0.0005^3 0.9995^2 + \binom{5}{3} 0.0005^3 0.9995^3 \\ &\quad + \binom{6}{4} 0.0005^3 0.9995^4 + \binom{7}{5} 0.0005^3 0.9995^5 \\ &= 6.9 \times 10^{-9} \end{aligned}$$

In R `pnbinom(5,3,0.0005)`

---



---

### 7.35 Examples

With the negative binomial probability function:

$$f(y) = \binom{y+r-1}{y} p^r (1-p)^y$$

We can now answer questions like:

- What is the probability of observing 10 correct pixels before 2 errors, if the probability of an error is 0.1?

$$f(10; r = 2, p = 0.1) = 0.03835463$$

in R `dnbinom(10, 2, 0.1)`

- What is the probability that 2 girls enter the class before 4 boys if the probability that a girl enters is 0.5?

$$f(2; r = 4, p = 0.5) = 0.15625$$





in R `dnbinom(2, 4, 0.5)`

---



---

## 7.36 Negative binomial distribution





## Chapter 8

# Poisson and Exponential Models

### 8.1 Objective

Discrete probability model:

- Poisson

Continuous probability model:

- Exponential
- 
- 

### 8.2 Discrete probability models

We are building up more complex models from simple ones:

**Uniform:** Classical interpretation of probability  $\downarrow$  **Bernoulli:** Introduction of a **parameter**  $p$  (family of models)  $\downarrow$  **Binomial:** **Repetition** of a random experiment ( $n$ -times Bernoulli trials)  $\downarrow$  **Poisson:** Repetition of random experiment within a continuous interval, having **no control** on when/where the Bernoulli trial occurs.

---

---

### 8.3 Counting events

Imagine that we are observing events that **depend** on time or distance **intervals**.

- cars arriving at a traffic light
- getting messages on your mobile phone
- impurities occurring at random in a copper wire

Suppose that the events are outcomes of **independent** Bernoulli trials each appearing randomly on a continuous interval, and we want to **count** them.

---



---

### 8.4 Counting events

What is the probability of observing  $X$  events in an interval's unit (time or distance)?

Imagine that some impurities in a copper wire deposit randomly along a wire

- at each centimeter, you would count an average of  $\lambda = 10/cm$ .
- divide the centimeter into micrometers ( $0.0001cm$ )

---



---

### 8.5 Poisson distribution

micrometers are small enough so

- either there is or there is not an impurity in each micrometer
- each micrometer can be considered a **Bernoulli trial**

---



---

### 8.6 Poisson distribution

The probability of observing  $X$  impurities in  $n = 10,000\mu$  (1cm) approximately follows a binomial distribution

$$P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x}$$

where  $p$  is the probability of finding an impurity in a micrometer.

Remember that  $E(X) = np$  so for  $\lambda = np$  (average number of impurities per 1cm), we can write


$$P(X = x) = \binom{n}{x} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x}$$

- There **could** still be two impurities in a micrometer so we need to increase the partition of the wire and  $n \rightarrow \infty$ .

Then in the limit:

$$P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$$

Where  $\lambda$  is constant because it is the density of impurities per centimeter, a **physical property** of the system.




## 8.7 Poisson distribution: Derivation details

For  $P(X = x) = \binom{n}{x} \left(\frac{\lambda}{n}\right)^x \left(1 - \frac{\lambda}{n}\right)^{n-x}$

in the limit ( $n \rightarrow \infty$ )

- $\frac{1}{n^x} \binom{n}{x} = \frac{1}{n^x} \frac{n!}{x!(n-x)!} = \frac{(n-x)!(n-x+1)\dots(n-1)n}{n^x x!(n-x)!} = \frac{n(n-1)\dots(n-x+1)}{n^x x!} \rightarrow \frac{1}{x!}$
- $\left(1 - \frac{\lambda}{n}\right)^n \rightarrow e^{-\lambda}$  (definition of exponential)
- $\left(1 - \frac{\lambda}{n}\right)^{-x} \rightarrow 1$

Therefore  $P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$



## 8.8 Poisson distribution

### Definition

Given

- an interval in the real numbers
- counts occur at random in the interval
- the average number of counts on the interval is known ( $\lambda$ )
- if one can find a small regular partition of the interval such that each of them can be considered Bernoulli trials

Then...

---



---

## 8.9 Poisson distribution

### Definition

The random variable  $X$  that counts events across the interval is a **Poisson** variable with probability mass function

$$f(x) = \frac{e^{-\lambda} \lambda^x}{x!}, \lambda > 0$$

### Properties:

- mean  $E(X) = \lambda$
- variance  $V(X) = \lambda$

---



---

## 8.10 Poisson distribution

With the Poisson probability function:

$$f(x) = \frac{e^{-\lambda} \lambda^x}{x!}$$

for  $x \in \{0, 1, \dots\}$

We can now answer questions like:

- What is the probability of receiving 4 emails in an hour, when the average number of emails in **one** hour is 1?

$$f(4; \lambda = 1) = 0.18$$

in R `dpois(2,1)`

- What is the probability of counting at least 10 cars arriving at a road toll in a minute, when the average number of cars that arrive at the toll in a minute is 5;  $P(X \leq 10) = F(10; \lambda = 5) = 0.98$ ?

in R `ppois(10,5)`


---



---



## 8.11 Poisson distribution



## 8.12 Continuous probability models

Continuous probability models are probability density functions  $f(x)$  of a continuous random variables that we **believe** describe real random experiments.

Definition:

Positive:

- $f(x) \geq 0$

*Allows us to compute probabilities using the area under the curve:*

- $P(a \leq X \leq b) = \int_a^b f(x)dx$

The probability of any value is 1:

- $\int_{-\infty}^{\infty} f(x)dx = 1$

### 8.13 Exponential density

Let's go back to the Poisson probability for the number of events ( $k$ ) in an interval

$$f(k) = \frac{e^{-\lambda} \lambda^k}{k!}, \lambda > 0$$

- Let's now consider only the first event
- the distance/time we have to wait until the first event is a **continuous** random variable.

We can ask for the probability that the first event is at distance  $X$ .

---



---

### 8.14 Exponential density

The probability of observing 0 events **if** an interval has unit  $x$  is

$$f(0|x) = \frac{e^{-x\lambda} x \lambda^0}{0!}$$

or

$$f(0|x) = e^{-x\lambda}$$

We can treat this as the conditional probability of 0 events in a distance  $x$ :  $f(K=0|X=x)$  and apply the Bayes theorem to reverse it:

$$f(x|0) = C f(0|x) = C e^{-x\lambda}$$

So we can calculate the **probability of observing a distance  $x$  with 0 events** (this is the distance until the first event or the distance between any two events).

---



---

### 8.15 Exponential density

In a Poisson process with parameter  $\lambda$  the probability of waiting a distance/time  $X$  until the first event has a **probability density**

$$f(x) = Ce^{-x\lambda}$$

- $C$  is a constant that ensures:  $\int_{-\infty}^{\infty} f(x)dx = 1$
- by integration  $C = \lambda$

Therefore

$$f(x) = \lambda e^{-\lambda x}$$



## 8.16 Exponential density

An exponential random variable  $X$  has a probability density

$$f(x) = \lambda e^{-\lambda x}, x \geq 0$$

### Properties:


- Mean:  $E(X) = \frac{1}{\lambda}$
- Variance:  $V(Y) = \frac{1}{\lambda^2}$

Where  $\lambda$  is its single parameter, known as a **decay rate**.

**Note:** The exponential model is a general model. It can describe the time/length until the first count in a Poisson process of the size of a whole made by a drill.



## 8.17 Exponential density



## 8.18 Exponential Distribution

In a Poisson process: ¿What is the probability of observing distance **smaller** than  $a$  until the first event?

Remember that this probability  $F(a) = P(X \leq a)$  is the probability density

$$F(a) = \lambda \int_{-\infty}^a e^{-x\lambda} dx = 1 - e^{-a\lambda}$$

- ¿What is the probability of observing a distance **larger** than  $a$  until the first event?

$$P(X > a) = 1 - P(X \leq a) = 1 - F(a) = e^{-a\lambda}$$

## 8.19 Exponential Distribution

With the exponential density function:

$$f(x) = \lambda e^{-\lambda x}$$

We can answer questions like:

- What is the probability that we have to wait for a bus for more than 1 hour when on average there are two buses per hour?

$$P(X > 1) = 1 - P(X \leq 1) = 1 - F(1, \lambda = 2) = 0.1353$$

in R `1-pexp(1,2)`

- What is the probability of having to wait less than 2 seconds to detect one particle when the radioactive decay rate is 2 particles each second;  
 $F(2, \lambda = 2)$

$$P(X \leq 2) = F(2, \lambda = 2) = 0.981$$


in R `pexp(2,2)`

---



---

## 8.20 Exponential Distribution



The median  $x_m$  is such that  $F(x_m) = 0.5$ . That is  $x_m = \frac{\log(2)}{\lambda}$

## Chapter 9

# Normal Distribution

### 9.1 Objective

Continuous probability model:

- Normal distribution

---

---

### 9.2 Continuous probability models

Continuous probability models are probability density functions  $f(x)$  of a continuous random variables that we **believe** describe real random experiments.

Definition:

Positive:

- $f(x) \geq 0$

*Allows us to compute probabilities using the area under the curve:*

- $P(a \leq X \leq b) = \int_a^b f(x)dx$

The probability of any value is 1:

- $\int_{-\infty}^{\infty} f(x)dx = 1$

---

---

### 9.3 Normal density

In 1801 Gauss analyzed the orbit of Ceres (large asteroid between Mars and Jupiter).

- People suspected it was a new planet.
- The measurements had errors.
- He was interested in finding how the observations were distributed so he could find the most probable orbit.
- He wanted to predict where astronomers should point their telescopes to find it a few months after it had passed behind the Sun.

---



---

### 9.4 Normal density

Errors due to measurement.

---



---

### 9.5 Normal density

He assumed that

- small errors were more likely than large errors
- error at a distance  $-\epsilon$  or  $\epsilon$  from the most likely measurement were equally likely
- the most **likely** altitude of Ceres at a given time in the sky was the **average** of multiple altitude measurements at that latitude.

---



---

### 9.6 Normal density

That was enough to show that the random deviations  $y$  **from the orbit** distributed like

$$f(y) = \frac{h}{\sqrt{\pi}} e^{-h^2 y^2}$$

\*The evolution of the Normal distribution, Saul Stahl, Mathematics Magazine, 2006.



---



---

## 9.7 Normal density

Let's write the distribution of errors

$$f(y) = \frac{h}{\sqrt{\pi}} e^{-h^2 y^2}$$

for the errors of measurements from the horizon  $X$  then  $y = x - x_0$

$$f(x) = \frac{h}{\sqrt{\pi}} e^{-h^2 (x-x_0)^2}$$

- The **mean** of this probability density is:

$E(X) = \mu = x_0$ , that represents the **true** position of Ceres from the horizon (property of the physical system).

- The **variance** is:

$V(X) = \sigma^2 = \frac{1}{2h^2}$ , that represents the dispersion of the error in the observations (property of the measurement system).

---



---

## 9.8 Definition

A random variable  $X$  defined in the real numbers has a **Normal** density if it takes the form

$$f(x; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, x \in \mathbb{R}$$

with mean and variance:


- $E(X) = \mu$
- $V(X) = \sigma^2$

$\mu$  and  $\sigma$  are the **two parameters** that fully describe the normal density function and their **interpretation** depends on the random experiment.

When  $X$  follows a Normal density, i.e. distributes normally, we write

$$X \rightarrow N(\mu, \sigma^2)$$

## 9.9 Normal probability density (Gaussian)



## 9.10 Normal distribution

The probability distribution of the Normal density:

$$F_{normal}(a) = P(Z \leq a)$$

is the **error** function defined by the area under the curve from  $-\infty$  to  $a$

$$F_{normal}(a) = \int_{-\infty}^a \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx$$

The function is found in most computer programs.

## 9.11 Normal distribution

When

$$X \rightarrow N(\mu, \sigma^2)$$

We can ask questions like:

- What is the probability that a woman in the population is at most 150cm tall if women have a mean height of 165cm with standard deviation of 8cm?

$$P(X \leq 150) = F(150, \mu = 165, \sigma = 8) = 0.03039636$$

in R `pnorm(150, 165, 8)`


- What is the probability that a woman's height in the population is between 165cm and 170cm?

$$P(165 \leq X \leq 170) = F(170, \mu = 165, \sigma = 8) - F(165, \mu = 165, \sigma = 8) = 0.2340145$$

in R `pnorm(170, 165, 8)-pnorm(165, 165, 8)`



## 9.12 Normal distribution




### 9.13 Normal distribution

- the mean  $\mu$  is also the median as it splits the measurements in two
- $x$  values that fall farther than  $2\sigma$  are considered **rare** 5%
- $x$  values that fall farther than  $3\sigma$  are considered **extremely rare** 0.2%

### 9.14 Normal distribution

We can define the limits of **common observations** for the distribution of women's height in the population.

- $P(165 - 8 \leq X \leq 165 + 8) = P(157 \leq X \leq 173) = 0.68$
- $P(165 - 2 \times 8 \leq X \leq 165 + 2 \times 8) = P(149 \leq X \leq 181) = 0.95$
- $P(165 - 3 \times 8 \leq X \leq 165 + 3 \times 8) = P(141 \leq X \leq 189) = 0.997$



## 9.15 Standard normal density

Let's change variables to a **standardized variable**

$$Z = \frac{X - \mu}{\sigma}$$

in the density

$$f(x; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, x \in \mathbb{R}$$

replacing  $x = \sigma z + \mu$  and  $dx = \sigma dz$  in the probability expression we have

$$P(x \leq X \leq x + dx) = P(z \leq Z \leq z + dz)$$

$$= \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx$$

$$= \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} dz$$

we obtain the **standardized** form of the normal density.

---



---

## 9.16 Standard normal density

### Definition

A random variable  $Z$  defined in the real numbers has a **standard** density if it takes the form

$$f(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} dz, z \in \mathbb{R}$$

with mean and variance

- $E(X) = 0$
- $V(X) = 1.$

---



---

## 9.17 Standard normal density

The standard density:

$$f(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} dz, z \in \mathbb{R}$$

- is the normal density  $N(\mu = 0, \sigma^2 = 1)$
- any normally distributed variable  $X$  can be transformed to a variable  $Z$

$$Z = \frac{x - \mu}{\sigma}$$

that follows a standard distribution:

$$Z \rightarrow N(0, 1)$$

---



---

## 9.18 Normal distribution

All normal densities can be obtained from the standard density with the values of  $\mu$  and  $\sigma$

---



---

## 9.19 Standard distribution

The probability distribution of the standard density:

$$\phi(a) = F_{standard}(a) = P(Z \leq a)$$

is the **error** function defined by

$$\phi(a) = \int_{-\infty}^a \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} dz$$

You can find it in most computer programs

---



---

## 9.20 Standard normal density

---




---

## 9.21 Standard normal density

We define the limits of the **most common observations** for the standard variable

- $P(-0.67 \leq X \leq 0.67) = 0.50$
- $P(-1.96 \leq X \leq 1.96) = 0.95$
- $P(-2.58 \leq X \leq 2.58) = 0.99$



## 9.22 Normal and standard distributions

---



---

For any normally distributed variable  $X$ , such that

$$X \rightarrow N(\mu, \sigma^2)$$

its distribution  $F(a) = P(X \leq a)$  can be computed from

$$F(a) = \Phi\left(\frac{a-\mu}{\sigma}\right)$$


---



---

## 9.23 Normal distribution

For computing  $P(a \leq X \leq b)$ , we use the property of the probability distributions

$$F(b) - F(a) = P(X \leq b) - P(X \leq a)$$

Let's standardize

$$\begin{aligned} &= P\left(\frac{X-\mu}{\sigma} \leq \frac{a-\mu}{\sigma}\right) - P\left(\frac{X-\mu}{\sigma} \leq \frac{b-\mu}{\sigma}\right) \\ &= P\left(Z \leq \frac{b-\mu}{\sigma}\right) - P\left(Z \leq \frac{a-\mu}{\sigma}\right) \\ &= \Phi\left(\frac{b-\mu}{\sigma}\right) - \Phi\left(\frac{a-\mu}{\sigma}\right) \end{aligned}$$

Then

$$F(b) - F(a) = \Phi\left(\frac{b-\mu}{\sigma}\right) - \Phi\left(\frac{a-\mu}{\sigma}\right)$$

The probabilities of **any normal variable** can be obtained from the **standard distribution**, after standardization (subtract the mean and divide by the standard deviation).

---



---

## 9.24 Summary of probability models

Model	X	range of x	f(x)	E(X)	V(X)
Uniform	integer or real number	$[a, b]$	$\frac{1}{n}$	$\frac{b+a}{2}$	$\frac{(b-a+1)^2-1}{12}$
Bernoulli	event A	0,1	$(1-p)^{1-x}p^x$	$p$	$p(1-p)$



Model	X	range of x	f(x)	E(X)	V(X)
Binomial	# of A events in $n$ repetitions of Bernoulli trials	0,1,...	$\binom{n}{x}(1-p)^{n-x}p^x$	$np$	$np(1-p)$
Negative Binomial for events	# of B events in Bernoulli repetitions before $r$ As are observed	0,1,..	$\binom{x+r-1}{x}(1-p)^x p^r$	$\frac{r(1-p)}{p}$	$\frac{r(1-p)}{p^2}$
Hypergeometric	# A events in a sample $n$ from population $N$ with $K$ As	$\max(0, n + K - N), \dots \min(K, n)$	$\frac{1}{\binom{N}{n}} \binom{K}{x} \binom{N-K}{n-x}$	$n * \frac{N}{K}$	$n \frac{N}{K} (1 - \frac{N}{K}) \frac{N-n}{N-1}$
Poisson	# of events A in an interval	0,1, ..	$\frac{e^{-\lambda} \lambda^x}{x!}$	$\lambda$	$\lambda$
Exponential	Interval between two events A	$[0, \infty)$	$\lambda e^{-\lambda x}$	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}$
Normal	measurement with symmetric errors whose most likely value is the average	$(-\infty, \infty)$	$\frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$	$\mu$	$\sigma^2$

## 9.25 R functions of probability models

Model	R
Uniform (continuous)	dunif(x, a, b)

Model	R
Binomial	dbinom(x,n,p)
Negative Binomial for events	dnbinom(x,r,p)
Hypergeometric	dhyper(x, K, N-K, n)
Poisson	dpois(x, lambda)
Exponential	dexp(x, lambda)
Normal	dnorm(x, mu, sigma)

## Chapter 10

# Sampling Distributions

### 10.1 Objective

Distributions for

- Sample mean (average)
- sample sum
- Sample variance

---

---

### 10.2 Normal distribution

When we have a normal random variable

$$X \rightarrow N(x; \mu, \sigma^2)$$

How do we estimate  $\mu$  and  $\sigma^2$ ?

- we need to take a **random sample**
- we need to **estimate** each parameter

---

---

### 10.3 Example: When we do not know the parameters

Imagine a client asking your metallurgical company to sell them 8 cables that can carry up to 96 Tons; that is 12 Tons each.

- You have in **stock** a set of cables that could do the job.

Can you use the cables in stock or do you need to produce new ones?

---



---

### 10.4 Example

you take a sample of 8 random experiments, each of which consists of loading a cable until it breaks and recording the breaking load.

These are the results: The observation of a **sample** of size 8

```
## [1] 13.34642 13.32620 13.01459 13.10811 12.96999 13.55309 13.75557 12.62747
```


- None of them broke at 12 Tons.
- There was one that broke at 12.62747 Tons.

Do you take the risk and sell a random sample of 8 cables from your stock?

---



---



## 10.5 Random sample

A **random sample** of size  $n$  is the **repetition** of a random experiment  $n$  **independent** times.

- A random sample is a  $n$ -dimensional **random variable**

$$(X_1, X_2, \dots, X_n)$$

where  $X_i$  is the  $i$ -th repetition of the random experiment with common distribution  $f(x; \theta)$  for any  $i$

- **One observation** of a random sample is the set of  $n$  values obtained from the experiments

$$(x_1, x_2, \dots, x_n)$$

Our **observation** of the sample of 8 cables was

```
## [1] 13.34642 13.32620 13.01459 13.10811 12.96999 13.55309 13.75557 12.62747
```

## 10.6 Example

We would like to compute  $P(X \leq 12)$ .

We are going to **assume** that the braking point is **normally distributed**.

$$X \rightarrow N(x; \mu, \sigma^2)$$

- For computing  $P(X \leq 12)$  we need the parameters  $\mu$  and  $\sigma^2$ .
  - How do we estimate the parameters from the observed sample?
- 
- 

## 10.7 Average or sample mean

### Definition

The sample mean (or average) of a **random sample** of size  $n$  is defined as

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

The average is a **random variable** that in our 8-size sample took the value

$$\bar{x}_{stock} = 13.21$$


---



---

## 10.8 The average as an estimator

This number can be used to **estimate** the unknown parameter  $\mu$  because:

- $E(\bar{X}) = E(X) = \mu$
- $V(\bar{X}) = \frac{V(X)}{n} = \frac{\sigma^2}{n}$

(since each random experiment in the sample is independent)

as

- $n \rightarrow \infty, V(\bar{X}) \rightarrow 0$

then

- $\bar{x}$  concentrates closer and closer to  $\mu$  as  $n$  increases.

We can take one value of  $\bar{x}$  as estimation for  $\mu$  or

$$\bar{x} = \hat{\mu}$$


---



---

## 10.9 Sample variance

### Definition

The **sample variance**  $S^2$  of a random sample of size  $n$

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

is the dispersion of the measurements about  $\bar{X}$ . In our 8-size sample  $S^2$  took the value

$$s_{stock}^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 = 0.1275608$$

The expected value of  $S^2$  is

- $E(S^2) = V(X) = \sigma^2$  (unbiased)

and therefore  $S^2$  is

- an estimator of  $V(X)$
- it also concentrates around  $\sigma^2$  because as  $n \rightarrow \infty$ ,  $V(\bar{S}^2) \rightarrow 0$  (consistent)

We can take one value of  $s^2$  as estimation for  $\sigma^2$  or

$$s^2 = \hat{\sigma}^2$$


---



---

## 10.10 Sample variance

$S^2$  aims to estimate the dispersion of the outcomes about  $\mu$  (the variance)


If we use  $\bar{X}$  as an estimator of  $\mu$  we need to correct for its dispersion (i.e. mean squared error of  $\bar{X}$ ).

The correction is achieved by dividing by  $n - 1$  and not  $n$  in the definition of  $S^2$

For:

$$S_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$E(S_n^2) = \sigma^2 - \frac{\sigma^2}{n} \neq \sigma^2 \text{ (we say that } S_n^2 \text{ is **biased** estimator of } \sigma)$$



## 10.11 Fitting a model

We **fit a model** when we

- **estimate** the parameters of the model

We also say we **train** a model (machine learning)


**Assuming** that

$$X \rightarrow N(x; \mu, \sigma^2)$$

Since we do not know the parameters, we **plugin** the estimates  $\bar{x}$  and  $s^2$  as the values of  $\mu$  and  $\sigma^2$



$$X \rightarrow N(x; \mu = 13.21, \sigma^2 = 0.3571565^2)$$



## 10.12 Prediction

We **predict** the value of an **outcome** when we compute its **probability**

What is the probability that the cable breaks at 12 Tons?

If we assumed the random variable

$$X \rightarrow N(x; \mu, \sigma^2)$$

We plug in the estimates  $\bar{x}$  and  $s^2$  into the probability distribution

$$P(X \leq 12) = F_{normal}(12; \mu = 13.21, \sigma^2 = 0.1275608)$$

In R `pnorm(12, 13.21, 0.3571565) = 0.000352188`

Given the **observed** sample, there is an estimated probability of 0.03% that a single cable will break at 12 Tons.

### 10.13 Inference

When we have a normal random variable

$$X \rightarrow N(\mu, \sigma^2)$$

and we know  $\mu$  and  $\sigma^2$ .

We can make inferences about  $\bar{X}$ , that is, calculate probabilities of the random variable  $\bar{X}$ .

When we make **inferences**, we usually ask the question:

How **sure** are we that the value of the estimator is **close** to the **true parameter**?

### 10.14 Example: When we do know the parameters

Let us imagine that our cables are certified to break with an average load of  $\mu = 13$  Tons with variance  $\sigma^2 = 0.35^2$ .

We take a random sample of 8 cables

## [1] 13.34642 13.32620 13.01459 13.10811 12.96999 13.55309 13.75557 12.62747

Can we say that we actually produced stronger cables because we got  $\bar{x} = 13.21$  from this sample of 8 cables?



We need to calculate probabilities of  $\bar{X}$ .

- What is the probability that the distance between  $\bar{X}$  and  $\mu$  is less than  $\bar{x}_{stock} - \mu = 0.21$ ?

$$P(-0.21 \leq \bar{X} - \mu \leq 0.21)$$

### 10.15 Density for $X$ and $\bar{X}$

If we **know** that the **true** parameters are  $\mu = 13$  and  $\sigma = 0.35$  this is what we would see



---

---

## 10.16 Sample mean distribution

**Theorem:** When  $X$  follows a normal distribution  $X \rightarrow N(\mu, \sigma^2)$

$\bar{X}$  is normal:

$$\bar{X} \rightarrow N\left(\mu, \frac{\sigma^2}{n}\right)$$

Then, if we **know**  $\mu$  and  $\sigma$  we can compute the true **probabilities of  $\bar{X}$**  using the normal distribution.

The mean and variance of  $\bar{X}$  are

- $E(\bar{X}) = \mu$
- $V(\bar{X}) = \frac{\sigma^2}{n}$

---



---

## 10.17 Inference on the average

**Example:**

If we **know** that the breaking load of our cables truly distribute as

$$X \rightarrow N(\mu = 13, \sigma^2 = 0.35^2)$$

then

$$\bar{X} \rightarrow N\left(13, \frac{0.35^2}{8}\right)$$

- $E(\bar{X}) = 13$
- $V(\bar{X}) = \frac{0.35^2}{8} = 0.01530169$

Our **observed error** in the estimation of the mean is the difference

$$\bar{x}_{stock} - \mu = 13.21 - 13 = 0.21$$

We ask: Is this a **typical** error?


---



---

## 10.18 Density for $\bar{X}$

If we **knew** that the **true** parameters were  $\mu = 13$  and  $\sigma = 0.35$  this is the error we would see



### 10.18.1 Probabilities of $\bar{X}$

If we **know** that the braking load of our cables **truly** distribute as

$$\bar{X} \rightarrow N\left(\mu = 13, \frac{\sigma^2}{n} = 0.1237^2\right)$$

What is the probability of observing an **error in estimation** of  $\mu$  (distance between  $\bar{X}$  and  $\mu$ ) smaller than 0.21?

We want to compute

$$P(-0.21 \leq \bar{X} - 13 \leq 0.21) = P(12.79 \leq \bar{X} \leq 13.21)$$

$$= F_{normal}(13.21; \mu, se^2) - F_{normal}(12.79; \mu, se^2)$$

In R we can compute it as:

`pnorm(13.21, 13, 0.1237)-pnorm(12.79, 13, 0.1237)=0.9104.`

91.0% of the errors are less than 0.21, therefore the **observed** error does not seem to be too typical (only 9% of the errors are higher).

Maybe we have stronger cables than we thought.

---

### 10.18.2 Sample sum

If we are interested in using all the 8 cables at the same time to carry a total of 96 Tons, then we should consider adding their individual contributions.

The **sample sum** is the **statistic**:

$$Y = n\bar{X} = \sum_{i=1}^n X_i$$

**Theorem:** if  $X \rightarrow N(\mu, \sigma^2)$  then

$$Y \rightarrow N(n\mu, n\sigma^2)$$

With mean and variance:

- $E(Y) = n\mu$
  - $V(Y) = n\sigma^2$
- 

### 10.18.3 Inference on the sample sum

If we **know** that for our cables

$$X \rightarrow N(\mu = 13, \sigma^2 = 0.35^2)$$

then

$$Y \rightarrow N(n\mu = 104, n\sigma^2 = 8 \times 0.35^2)$$

- $E(Y) = 104$
- $V(Y) = 8 \times 0.35^2 = 0.98$

For our 8-sample, we observed

- $y_{stock} = 105.7014$

and, therefore, the **observed error** in the estimation of the mean of the **true** total braking load ( $n\mu$ ) of 8 cables was

- $y_{stock} - n\mu = 1.7014$

Is this a **typical** error?

---



---

#### 10.18.4 Probabilities of the sample sum: Propagation of error

What is the probability of observing a difference  $Y - E(Y)$  smaller than 1.7014?

We want to compute the probability

$$P(-1.7014 \leq \bar{Y} - 104 \leq 1.7014) = P(102.2986 \leq Y \leq 105.7014)$$

$$= F_{normal}(105.7014; n\mu, n\sigma^2) - F_{normal}(102.2986; n\mu, n\sigma^2)$$

In R we can compute it as:

$$\text{pnorm}(105.7014, 104, \text{sqrt}(0.98)) - \text{pnorm}(102.2986, 104, \text{sqrt}(0.98)) = 0.914.$$

91.4% of the times we obtain sample sums that are smaller than 1.7014

This proportion is higher than the proportion for individual cables.

---



---

### 10.19 Inference in the sample variance

Consider a quality control process that requires that the cables are produced close to the specified value  $\mu$ .

If a sample of 8 cables is too dispersed ( $S^2 > 0.3$ ), we stop production: the process is out of control.

What is the probability that the sample variance of a sample of 8 cables is greater than the required 0.3?

---



---

### 10.20 Probabilities of the sample variance

**Theorem:** When  $X$  follows a normal distribution

$$X \rightarrow N(\mu, \sigma^2)$$

The **statistic**:

$$W = \frac{(n-1)S^2}{\sigma^2} \rightarrow \chi^2(n-1)$$

has a  $\chi^2$  (chi-squared) distribution with  $df = n-1$  degrees of freedom given by

$$f(w) = C_n w^{\frac{n-3}{2}} e^{-\frac{w}{2}}$$

where:


- $C_n = \frac{1}{2^{(n-1)/2} \sqrt{\pi(n-1)}}$  ensures  $\int_{-\infty}^{\infty} f(t) dt = 1$
- $\Gamma(x)$  is Euler's factorial for real numbers
- If we **know** the true values of  $\mu$  and  $\sigma$  we can compute probabilities of  $S^2$  using the  $\chi^2$  distribution for  $W$ .

---



---

## 10.21 $\chi^2$ -statistic




---



---



## 10.22 $\chi^2$ -statistic

If we **know** that our cables trully distribute as

$$X \rightarrow N(\mu = 13, \sigma^2 = 0.35^2)$$

then we can compute

$$\begin{aligned} P(S^2 > 0.2) &= P\left(\frac{(n-1)S^2}{\sigma^2} > \frac{(n-1)0.3}{\sigma^2}\right) \\ &= P\left(W > \frac{(n-1)0.3}{\sigma^2}\right) \\ &= 1 - P\left(W \leq \frac{(n-1)0.3}{\sigma^2}\right) = 1 - P\left(W \leq \frac{(8-1)0.3}{0.1225}\right) \\ &= 1 - F_{\chi^2, df=7}(17.14286) = 0.016 \end{aligned}$$

In R 1-pchisq(17.14286, df=7)=0.016

There is only a probability of 1% of obtaining a value greater than  $s^2 = 0.3$ .

- $s^2 > 0.3$  seems to be a good criterion to stop production and revise the process.
- our observed value was  $s_{stock}^2 = 0.1275608$
- the sample is not too dispersed and we believe that the production is under control.



# Chapter 11

## Exercises

### 11.1 Data description

#### 11.1.0.1 Exercise 1

We have performed an experiment 8 times with the following results

```
## [1] 3 3 10 2 6 11 5 4
```

Answer the following questions:

- Compute the relative frequencies of each outcome.
- Compute the cumulative frequencies of each outcome.
- What is the average of the observations?
- What is the median?
- What is the third quartile?
- What is the first quartile?

#### 11.1.0.2 Exercise 2

We have performed an experiment 10 times with the following results

```
## [1] 2.875775 7.883051 4.089769 8.830174 9.404673 0.455565 5.281055 8.924190  
## [9] 5.514350 4.566147
```

Consider 10 bins of size 1:  $[0,1]$ ,  $(1,2]$ ... $(9,10]$ .

Answer the following questions:

- Compute the relative frequencies of each outcome and draw the histogram
- Compute the cumulative frequencies of each outcome and sketch the cumulative plot.
- Sketch a boxplot.

## 11.2 Probability

### 11.2.0.1 Exercise 1

The outcome of one random experiment is to measure the misophonia severity **and** depression status of one patient.

- Misophonia severity:  $x \in \{0, 1, 2, 3, 4\}$
- Depression:  $y \in \{0, 1\}$  (no:0, yes:1)

```
## Misofonia.dic depression.dic
## 1           4           1
## 2           2           0
## 3           0           0
## 4           3           0
## 5           0           0
## 6           0           0
```

A large study on 123 patients showed the frequencies  $n_{x,y}$  given in the contingency table:

```
##
##           Depression:0 Depression:1
## Misophonia:4           0           9
## Misophonia:3          25           6
## Misophonia:2          34           3
## Misophonia:1           5           0
## Misophonia:0          36           5
```

Let's assume that  $N \gg 0$  and that the frequencies **estimate** the probabilities  $f_{x,y} = \hat{P}(X, Y)$

```
##
##           Depression:0 Depression:1
## Misophonia:4  0.00000000  0.07317073
## Misophonia:3  0.20325203  0.04878049
## Misophonia:2  0.27642276  0.02439024
## Misophonia:1  0.04065041  0.00000000
## Misophonia:0  0.29268293  0.04065041
```

- What is the marginal probability of misophonia severity 3? (R/0.3)
- What is the probability of not being misophonic **and** not depressed? (R/0.293)
- What is the probability of being misophonic **or** depressed? (R/0.293)
- What is the probability of being misophonic **and** depressed? (R/0.707)
- Describe in English the outcomes with probability 0.

### 11.2.0.2 Exercise 2

We have performed an experiment 10 times with the following results

```
##      A      B
## 1   male  dead
## 2   male  dead
## 3   male  dead
## 4  female alive
## 5   male  dead
## 6  female alive
## 7  female  dead
## 8  female alive
## 9   male  alive
## 10  male  alive
```

- Create the contingency table for the number ( $n_{i,j}$ ) of observations of each outcome ( $A, B$ )
- Create the contingency table for the relative frequency ( $f_{i,j}$ ) of the outcomes
- What is the marginal frequency of being male? (R/0.6)
- What is the marginal frequency of being alive? (R/0.5)
- What is the frequency of being alive **or** female? (R/0.6)

## 11.3 Conditional Probability

### 11.3.0.1 Exercise 1

A machine is tested for its performance to produce high-quality turning rods. These are the results of the testing

	Rounded: Yes	Rounded: No
smooth surface: yes	200	1
smooth surface: no	4	2

- What is the estimated probability that the machine produces a rod that does not satisfy any quality control? (R: 2/207)
- What is the estimated probability that the machine produces a rod that does not satisfy at least one quality control?(R: 7/207)
- What is the estimated probability that the machine produces rounded and smoothed surfaced rods? (R: 200/207)
- what is the estimated probability that the rod is rounded if the rod is smooth? (R: 201/201)
- what is the estimated probability that the rod is smooth if it is rounded? (R: 201/204)
- what is the estimated probability that the rod is neither smooth nor rounded if it does not satisfy at least one quality control? (R: 2/7)

- Are smoothness and roundness independent events? (no)

### 11.3.0.2 Exercise 2

We develop a test to detect the presence of bacteria in a lake. We find that if the lake contains the bacteria the test is positive 70% of the time. If there are no bacteria then the test is negative 60% of the time. We deploy the test in a region where we know that 20% of the lakes have bacteria.

- What is the probability that one lake that tests positive is contaminated with bacteria? (R: 0.30)

### 11.3.0.3 Exercise 3

Two machines are tested for their performance to produce high-quality turning rods. These are the results of the testing

#### Machine 1

	Rounded: Yes	Rounded: No
smooth surface: yes	200	1
smooth surface: no	4	2

#### Machine 2

	Rounded: Yes	Rounded: No
smooth surface: yes	145	4
smooth surface: no	8	6

- what is the probability that the rod is rounded? (R: 357/370)
- What is the probability that the rod has been produced by machine 1? (R: 207/370)
- what is the probability that the rod is not smooth? (R: 20/370)
- What is the probability that the rod is smooth or rounded or produced by machine 1? (R: 364/370)
- What is the probability that the rod is rounded if it is smoothed and from machine 1? (R: 200/201)
- What is the probability that the rod is not rounded if it is not smoothed and is from machine 2? (R: 6/8)
- what is the probability that the rod has come from machine 1 if it is smoothed and rounded? (R: 200/345)
- what is the probability that the rod has come from machine 2 if it does not pass at least one of the quality controls? (R:0.72)

**11.3.0.4 Exercise 4**

We want to cross an avenue with two traffic lights. The probability of finding the first traffic light in red is 0.6. If we stopped at the first traffic light, the probability of stopping at the second one is 0.15. Whereas the probability of stopping on the second one if we do not stop on the first one is 0.25.

When we try to cross both traffic lights:

- what is the probability of having to stop at each traffic light? (R:0.09)
- What is the probability of having to stop at at least one traffic light? (R:0.7)
- What is the probability of having to stop at only one traffic light? (R:0.61)
- If I stopped at the second traffic light, what is the probability that I had to stop at the first one? (R: 0.62)
- If I had to stop at any traffic light, what is the probability that I had to do it twice? (R: 0.12)
- Is stopping at the first traffic light an independent event from stopping at the second traffic light? (no)

Now, we want to cross an avenue with three traffic lights. The probability of finding a traffic light in red only depends on the previous one. In particular, the probability of finding one traffic light in red given that the previous one was in red is 0.15. Whereas, the probability of finding one traffic right in red given that the previous one was in green is 0.25. Also, the probability of finding the first traffic light in red is 0.6.

- What is the probability of having to stop at each traffic light? (R:0.013)
- What is the probability of having to stop at at least one traffic light? (R:0.775)
- What is the probability of having to stop at only one traffic light? (R:0.5425)

hints:

- If the probability that one traffic light is red depends only on the previous one then  $P(R_3|R_2, R_1) = P(R_3|R_2, \bar{R}_1) = P(R_3|R_2)$  and  $P(R_3|\bar{R}_2, R_1) = P(R_3|\bar{R}_2, \bar{R}_1) = P(R_3|\bar{R}_2)$
- The joint probability of finding three traffic lights in red can be written as:  $P(R_1, R_2, R_3) = P(R_3|R_2)P(R_2|R_1)P(R_1)$

**11.3.0.5 Exercise 5**

A quality test on a random brick is defined by the events:

- Pass quality test:  $E$ , do no pass quality test:  $\bar{E}$
- Defective:  $D$ , non-defective:  $\bar{D}$

If the diagnostic test has sensitivity  $P(E|\bar{D}) = 0.99$  and specificity  $P(\bar{E}|D) = 0.98$ , and the probability of passing a test is  $P(E) = 0.893$  then

- what is the probability that a brick chosen at random is defective  $P(D)$ ? (R:0.1)
- What is the probability that a brick that has passed the test is really defective? (R:0.022)
- The probability that a brick is not defective **and** that it does not pass the test (R:0.009)
- Are  $D$  and  $\bar{E}$  statistical independent? (no)

## 11.4 Random variables

### 11.4.0.1 Exercise 1

Given the probability mass function

$x$	$f(x) = P(X = x)$
10	0.1
12	0.3
14	0.25
15	0.15
17	?
20	0.15

- what is its expected value and standard deviation? (R: 14.2; 2.95)

### 11.4.0.2 Exercise 2

Given the probability distribution for a discrete variable  $X$

$$F(x) = \begin{cases} 0, & x < -1 \\ 0.2, & x \in [-1, 0) \\ 0.35, & x \in [0, 1) \\ 0.45, & x \in [1, 2) \\ 1, & x \geq 2 \end{cases}$$

- find  $f(x)$
- find  $E(X)$  and  $V(X)$  (R:1; 1.5)
- what is the expected value and variance of  $Y = 2X + 3$  (R: 6)
- what is the median and the first and third quartiles of  $X$ ? (R:2,0,2)

### 11.4.0.3 Exercise 3

We are testing a system to transmit digital pictures. We first consider the experiment of sending 3 pixels and having as **possible** outcomes events such



like  $(0, 1, 1)$ . This is the event of receiving the first pixel with no error, the second with error and third with error.

- List in one column the sample space of the random experiment.
- In the a second column assign the random variable that counts the number of errors transmitted for each outcome

Consider that we have a totally noisy channel, that is any outcome of three pixels is equally likely.


- What is the probability of receiving 0, 1, 2, or 3 errors in the transmission of 3 pixels? (R:  $1/8$ ;  $3/8$ ;  $3/8$ ;  $1/8$ )
- Sketch the probability mass function for the number of errors
- What is the expected value for the number of errors? (R:1.5)
- What is its variance? (R: 0.75)
- Sketch the probability distribution
- What is the probability of transmitting at least 1 error? (R:  $7/8$ )

#### 11.4.0.4 Exercise 4

- for the probability density

$$f(x) = \begin{cases} \frac{1}{100}, & \text{if } x \in (0, 100) \\ 0, & \text{otherwise} \end{cases}$$

- compute the mean (R:50)
- compute variance using  $E(X^2) = V(X) + E(X)^2$  (R:  $100^2/12$ )
- compute  $P(\mu - \sigma \leq X \leq \mu + \sigma)$  (R: 0.57)
- What are the first and third quartiles? (R: 25; 75)



#### 11.4.0.5 Exercise 5

Given

$$f(x) = \begin{cases} 0, & x < 0 \\ ax, & x \in [0, 3] \\ b, & x \in (3, 5) \\ \frac{b}{3}(8 - x), & x \in [5, 8] \\ 0, & x > 8 \end{cases}$$

- What are the values of  $a$  and  $b$  such that  $f(x)$  is a continuous probability density function? (R: 1/15; 1/5)
- what is the mean of  $X$ ? (R:4)

#### 11.4.0.6 Exercise 6

For the probability density

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & \text{if } x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

- Confirm that this is a probability density
- Compute the mean (R:  $1/\lambda$ )
- Compute the expected value of  $X^2$  (R:  $2/\lambda^2$ )
- Compute variance (R:  $1/\lambda^2$ )

- Find the probability distribution  $F(a)$  (R:  $1 - \exp(-\lambda a)$ )
- Find the median (R:  $\log 2/\lambda$ )

**11.4.0.7 Exercise 7**

Given the cumulative distribution for a random variable X

$$F(x) = \begin{cases} 0, & x < -1 \\ \frac{1}{80}(17 + 16x - x^2), & x \in [-1, 7) \\ 1, & x \geq 7 \end{cases}$$

compute:

- $P(X > 0)$  (R: 63/80)
- $E(X)$  (R: 1.93)
- $P(X > 0 | X < 2)$  (R: 28/45)

**11.5 Probability Models****11.5.0.1 Exercise 1**

In a population, the probability that a baby boy is born is  $p = 0.51$ . Consider a family of 4 children

- What is the probability that a family has only one boy? (R: 0.240)
- What is the probability that a family has only one girl? (R: 0.259)
- What is the probability that a family has only one boy or only one girl? (R: 0.4999)
- What is the probability that the family has at least two boys? (R: 0.7023)
- What is the number of children that a family should have such that the probability of having at least one girl is more than 0.75? (R:  $n = 3 > \log(0.25)/\log(0.51)$ )

**11.5.0.2 Exercise 2**

A search engine fails to retrieve information with a probability 0.1

- If we system receives 50 search requests, what is the probability that the system fails to answer three of them? (R: 0.1385651)
- What is the probability that the engine successfully completes 15 searches before the first failure? (R: 0.020)
- We consider that a search engine works sufficiently well when it is able to find information for more than 10 requests for every 2 failures. What is the probability that in a reliability trial our search engine is satisfactory? (R: 0.697)

**11.5.0.3 Exercise 3**

The average number of radioactive particles hitting a Geiger counter in a nuclear energy plant under control is 2.3 per minute.

- What is the probability of counting exactly 2 particles in a minute? (R:0.265)
- What is the probability of detecting exactly 10 particles in 5 minute? (R:0.112)
- What is the probability of at least one count in two minutes? (R:0.9899)
- What is the probability of having to wait less than 1 second to detect a radioactive particle, after we switch on the detector? (R:0.037)
- We suspect that a nuclear plant has a radioactive leak if we wait less than 1 second to detect a radioactive particle, after we switch on the detector. What is the probability that when we visit in 5 plants that are under control, we suspect that at least one has a leak? (R:0.1744).

**11.5.0.4 Exercise 4**

- What is the probability that a man's height is at least 165cm if the population mean is 175cm y the standard deviation is 10cm? (R:0.841)
- What is the probability that a man's height is between 165cm and 185cm? (R:0.682)
- What is the height that defines the 5% of the smallest men? (R:158.55)

**11.6 Sampling and Central Limit Theorem****11.6.0.1 Exercise 1**

A battery model charges up to 75% of its capacity within an hour with a standard deviation of 15%.

- If we charge 25 batteries, what is the probability that the difference in charge between the sample average and the mean charge is at most 5%? (R:0.9044)
- If we charge 100, what is that probability? (R:0.9991)
- If, instead we only charge 9 batteries, which value  $c$  is surpassed by the sample mean with probability of 0.015? (R:85.850)

**11.6.0.2 Exercise 2**

An electronic component is needed for the correct functioning of a telescope. It needs to be replaced immediately when it wears out.

The mean life of the component ( $\mu$ ) is 100 hours and its standard deviation  $\sigma$  is 30 hours.

- what is the probability that the average of the mean life of 50 components is within 1 hour from the mean life of a single component?
- How many components do we need such that the telescope is operational 2750 consecutive hours with 0.95 probability?

### 11.6.0.3 Exercise 3

An automated machine fills test tubes with biological samples with mean  $\mu = 130\text{mg}$  and a standard deviation of  $\sigma = 5\text{mg}$ .

- for a random sample of size 50. What is the probability that the sample mean (average) is between 128 and 132gr?
- what should be the size of the sample ( $n$ ) such that the sample mean  $\bar{X}$  is higher than 131gr with a probability less or equal than 0.025?

### 11.6.0.4 Exercise 4

In the Caribbean, there appears to be an average of 6 hurricanes per year. Considering that hurricane formation is a Poisson process, meteorologists plan to estimate the mean time between the formation of two hurricanes. They plan to collect a sample of size 36 for the times between two hurricanes.

- What is the probability that their sample average is between 45 and 60 days?
- Which should be the sample size such that they have a probability of 0.025 that the sample mean is greater than 70 days?

### 11.6.0.5 Exercise 5

The probability that a particular mutation is found in the population is 0.4. If we test 2000 people for the mutation:

- What is the probability that the total number of people with the mutation is between 791 and 809?

hint: Use the CLT with a sample of 2000 Bernoulli trials. This is known as the normal approximation of the binomial distribution.

## 11.7 Point Estimators

### 11.7.0.1 Exercise 1

Consider the probability model

$$f(x) = \begin{cases} 1/2 - a, & \text{if } x = -1 \\ 1/2, & \text{if } x = 0 \\ a, & \text{if } x = 1 \end{cases}$$

where  $a$  is a parameter.

Compute the mean and variance of the statistic:

$$T = \frac{\bar{X}}{2} + \frac{1}{4}$$

where  $\bar{X} = \frac{1}{N} \sum_{i=1}^N X_i$

- is  $T$  a biased estimator of  $a$ ?
- is  $T$  consistent? i.e.  $V(T) \rightarrow 0$  when  $N \rightarrow \infty$

#### 11.7.0.2 Exercise 2

- Is  $\bar{X}^2 = (\frac{1}{N} \sum_{i=1}^N X_i)^2$  an unbiased estimator of  $E(X)^2$ ?

## 11.8 Maximum likelihood

#### 11.8.0.1 Exercise 1

For a random variable with a binomial probability function

$$f(x; p) = \binom{n}{x} p^x (1-p)^{n-x}$$

- What is the maximum-likelihood estimator of  $p$  for a sample of size 1 of this random variable?
- In **one** exam of 100 students we observed  $x_1 = 68$  students that passed the exam. What is the estimate of the  $p$ ?

#### 11.8.0.2 Exercise 2

Take a random variable with the following probability density function

$$f(x) = \begin{cases} (1 + \theta)x^\theta, & \text{if } x \in (0, 1) \\ 0, & \text{if } x \notin (0, 1) \end{cases}$$

- What is the maximum likelihood estimate for  $\theta$ ?
- If we take a 5-sample with observations  $x_1 = 0.92$ ;  $x_2 = 0.79$ ;  $x_3 = 0.90$ ;  $x_4 = 0.65$ ;  $x_5 = 0.86$

What is the estimated value of the parameter  $\theta$ ?

**11.8.0.3 Exercise 3**

Take a random variable with the following probability density function

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & \text{if } 0 \leq \\ 0, & \text{otherwise} \end{cases}$$

- What is the maximum likelihood estimate for  $\lambda$ ?
- If we take a 5-sample with observations  $x_1 = 0.223$   $x_2 = 0.681$ ;  $x_3 = 0.117$ ;  $x_4 = 0.150$ ;  $x_5 = 0.520$

What is the estimated value of the parameter  $\lambda$ ?

**11.9 Method of moments****11.9.0.1 Exercise 1**

What are the estimators of the following parametric models given by the method of moments?

Model	f(x)	E(X)
Bernoulli	$p^x(1-p)^{1-x}$	$p$
Binomial	$\binom{n}{x}p^x(1-p)^{n-x}$	$np$
Shifted geometric	$p(1-p)^{x-1}$	$\frac{1}{p}$
Negative Binomial	$\binom{x+r-1}{x}p^r(1-p)^x$	$r\frac{1-p}{p}$
Poisson	$\frac{e^{-\lambda}\lambda^x}{x!}$	$\lambda$
Exponential	$\lambda e^{-\lambda x}$	$\frac{1}{\lambda}$
Normal	$\frac{1}{\sqrt{2\pi}\sigma}e^{-\frac{(x-\mu)^2}{2\sigma^2}}$	$\mu$

**11.9.0.2 Exercise 2**

Take a random variable with the following probability density function

$$f(x) = \begin{cases} (1+\theta)x^\theta, & \text{if } x \in (0, 1) \\ 0, & x \notin (0, 1) \end{cases}$$

- Compute  $E(X)$  as a function of  $\theta$
- What is the estimate for  $\theta$  using the method of moments?
- If we take a 5-sample with observations  $x_1 = 0.92$ ;  $x_2 = 0.79$ ;  $x_3 = 0.90$ ;  $x_4 = 0.65$ ;  $x_5 = 0.86$

What is the estimated value of the parameter  $\theta$ ?

**11.9.0.3 Exercise 3**

Consider a discrete random variable  $X$  that follows a negative binomial distribution with probability mass function:

$$f(x) = \binom{x+r-1}{x} p^r (1-p)^x$$

Given that

- $E(X) = \frac{r(1-p)}{p}$
- $V(X) = \frac{r(1-p)}{p^2}$

compute:

- An estimate for the parameter  $r$  and an estimate for the parameter  $p$  obtained from a random sample of size  $n$  using the method of moments.
- The values of the estimates of  $r$  y  $p$  for the folowing random sample:

$$x_1 = 27; \quad x_2 = 8; \quad x_3 = 22; \quad x_4 = 29; \quad x_5 = 19; \quad x_5 = 32$$

**11.10 Confidence intervals****11.10.0.1 Exercise 1**

In a scientific paper the authors report a 95% confidence interval of (228, 232) for the natural frequency (Hz) of metallic beam. They used a sample of size 25 and considered that the measurements distributed normally.

- What is the mean and the standard deviation of the measurements?
- Compute the 99% confidence interval.

hints:

- in R  $t_{0.025, 24} = \text{qt}(0.975, 24) \sim 2$
- in R  $t_{0.005, 24} = \text{qt}(0.995, 24) \sim 2.8$

**11.10.0.2 Exercise 2**

compute 95% CI the mean of a normal variable with known variance  $\sigma^2 = 9$  and  $\bar{x} = 22$ , using a sample of size 36.



**11.10.0.3 Exercise 3**

This year, 17 from 1000 of patients with influenza developed complications.

- Compute the 99% confidence interval for the proportion of complications.
- The previous year 2% showed complications. Can we say with 99% confidence that this year there is a significant drop in influenza complications?

**11.11 Hypothesis testing****11.11.0.1 Exercise 1**

Imagine we take a random sample of size  $n = 41$  of a normal random variable  $X$ , and find that the sample average is 10 and the sample variance is 1.5.

- What is then the confidence interval for the mean of  $X$  at 95% confidence level?

Consider that  $t_{0.025,40} = \text{qt}(0.975, 40) \sim 2$ .

- Test the hypothesis that the mean of  $X$  is **different** than 10.5, using a 5% significance threshold.
- Write the code to calculate the P-value to test the hypothesis that the mean of  $\mu$  is **lower** than 10.5, using a 5% significance threshold.

Consider that the code for the T probability distribution with  $n - 1$  degrees of freedom is `pt(tobs, n-1)`.

**11.11.0.2 Exercise 2**

10 gas condensates showed the following concentrations of mercury (in *ng/ml*):

23.3, 22.5, 21.9, 21.5, 19.9, 21.3, 21.7, 23.8, 22.6, 24.7

Assuming that the mercury concentration is distributed normally across gas condensates, test the hypothesis that a condensate does not surpass the toxicity limit established at *24ng/ml*.

**11.11.0.3 Exercise 3**

The manufacturer of gene expression microarrays guarantees that at least 97% of the microarrays they produce have high quality signals. A customer receives a batch of 200 pieces and finds that 8 unperformed.

Should the customer return the lot due to poor quality?