

	INFORMATION AND CONSENT DOCUMENT
--	---

**INFORMED CONSENT DOCUMENT
FOR MINOR SURGERY (LIPOMODELLING)**

In observance of Act 41/2002, regulating patient's autonomy and rights and duties concerning information and clinical documents (BOE 15-11-02), we offer you, in writing and in an understandable manner, information about **LIPOMODELLING**. However, said information has already been concisely and orally provided to you by the undersigned doctor, who also, and at your request, has addressed all your questions, as well as those related to the treatment. This document is just the graphic transcription of said information, to help you understand it and motivate a thoughtful and deliberate authorization.

It is important that you read this information carefully and in its entirety. Please sign or put your initials on every page, thus indicating that you have read and signed the consent document for the procedure recommended by your doctor.

In _____, on _____,
Patient Name _____
Date of Birth _____ ID _____
Address _____
Town _____
Phone Numbers _____

I HEREBY STATE:

That I REQUEST AND AUTHORIZE Doctor _____,
_____, graduated in Medicine and Surgery,
with medical license number _____, to perform upon me a treatment known as
LIPOMODELLING (liposuction, lipolaser), having chosen this instead of the alternatives, namely
_____.

I KNOW and ACCEPT the physician's professional training to perform this treatment.

The substances and machines involved have been authorized for use in aesthetic medicine, and show the CE brand and its corresponding sanitary registration number.

I CONFIRM that the treatment mentioned above has been fully explained to me by the physician in words I can understand, including typical risks involved, undesired effects, the risks specific to me, and also discomfort or, occasionally, pain I may feel with normal post-treatment. I have also been told about other options available in the market, together with pros and cons of each one. Taking this into account, I have chosen the treatment described above.

I have also been informed, in terms of probability, about the results of the procedure according to references from comparative scientific literature, and the professional's previous experience executing these procedures.

I have been informed about, and I accept, the commonly known risks of premedication, anesthesia and/or the treatment to be performed.

I CONSENT, if needed, to the administration of local anesthesia by, or under instructions of, Dr.

I KNOW and ACCEPT the physician's professional training to perform the suggested treatment.

BRIEF EXPLANATION OF THE TREATMENT:

Liposuction is a surgical technique for the removal of unwanted fat deposits from specific areas of the body, including the face and neck, arms, trunk, abdomen, buttocks, hips and thighs, and the knees, calves and ankles. This is not a substitute for weight reduction, but a method to remove localized deposits of fatty tissue that do not respond to diet or exercise.

The best candidates for liposuction are individuals who have excess fat in particular body areas. Having firm, elastic skin will result in a better final contour after liposuction. Skin that has lost its tone due to stretch marks, weight loss or natural aging will not reshape itself to the new contours and may require additional surgical techniques to remove and tighten excess skin. Body-contour irregularities due to structures other than fat cannot be improved by this technique. Liposuction alone will not improve areas of dimpled skin known as "cellulite".

Liposuction is performed with a hollow metal surgical instrument known as a *cannula* that is inserted through small skin incision(s), and passed back and forth through the area of fatty deposit. The cannula is attached to a vacuum source, which provides the suction needed to remove the fatty tissue.

In some situations, a special cannula may be used that emits LASER energy to break down fatty deposits. This technique is known as LASER-assisted liposuction or lipolaser. Depending on your needs, your doctor may recommend liposuction alone, or in combination with LASER-assisted liposuction.

Liposuction can be performed with local or general anesthesia. The tumescent liposuction technique (local anesthesia) involves the infiltration of fluid containing diluted local anesthetic and epinephrine into areas of fatty deposits. This technique can reduce discomfort at the time of surgery, blood loss and post-operative bruising.

Support garments (girdles) and dressings are worn to control swelling and stimulate healing.

This procedure consists of:

- Infiltration with local anesthesia or Klein's solution in the area to be treated.
- Prior aspiration, or without lipolaser (980 or Nd:YAG 1064 diode) of the fat in the areas to be treated.

- Removal of stitches 5-7 days later.
- The residual scar will look red for a few months, subsequently turning white.

This intervention can be performed with Kalinox (50% nitrous oxide) or with intravenous sedation. Said sedation will be administered by the anesthetist, Dr. _____.

RISKS of LIPOSUCTION (lipolaser or ultrasound-assisted)

Every surgical procedure involves a certain amount of risk, and it is important that you understand the risks and possible complications associated with said procedures. Furthermore, all procedures have limitations. An individual's choice to undergo a surgical procedure is based on the comparison of the risk to the potential benefit. Although most patients do not experience these complications, you should discuss each of them with your doctor to make sure you understand every possible consequence of liposuction.

Patient Screening: Individuals with poor skin tone, medical problems, morbid obesity, or unrealistic expectations may not be candidates for liposuction.

Bleeding: With local anesthesia, it is possible, though unusual, to have a bleeding episode during or after surgery. Should post-operative bleeding occur, you may require emergency treatment to drain accumulated blood or a blood transfusion. Hematomas can occur at any time after surgery and may contribute to infections or other problems. Do not take any aspirin or anti-inflammatory medications for seven days before and after the surgery, as these may increase the risk of bleeding. Nonprescription "herbs" and dietary supplements can increase the risk of surgical bleeding.

Infection: Infections are very rare after the surgery. Should an infection occur, additional treatment, including antibiotics, hospitalization or additional surgery may be necessary.

Scarring: All surgeries leave a scar; in the case of liposuction, they are very small. Although the wound is expected to heal well after a surgical procedure, abnormal scars can occur in the skin and deeper tissues. Scars can be unattractive and have a different color than the surrounding skin. The scar appearance can also vary within the scar itself, it can present variations in outline, and it may "frown" due to excess skin. Scars can be asymmetrical (appear different on the right and left sides of the body). There is the possibility of visible marks in the skin caused by sutures.

Changes in skin sensitivity: It is common to experience a decrease in (or loss of) skin sensitivity in the areas where a surgical procedure was performed. This usually resolves with time. The decrease in (or complete loss of) skin sensitivity is rare, and may not totally resolve.

Skin discoloration/swelling: Bruising and swelling normally occur following liposuction. The skin in or near the surgical site may appear either lighter or darker than the surrounding skin. In rare situations, swelling and skin discoloration may persist for long periods of time. Permanent skin discoloration is rare. This process can be improved with manual lymphatic drainage immediately after surgery.

Skin-contour irregularities: Contour and shape irregularities and depressions may occur after liposuction. Visible and palpable wrinkles can also occur in the skin. This process can be improved with manual lymphatic drainage immediately after surgery.

Asymmetry: It may not be possible to achieve symmetrical body appearance from liposuction surgery. Factors such as skin tone, fat deposits, bony prominence, and muscle tone may contribute to normal asymmetry in body features. An additional liposuction to try to improve asymmetry may be necessary.

Seroma: Fluid accumulations in areas where liposuction has been performed are rare. Treatments to drain accumulations of fluid may be necessary. This process can be prevented using a compression girdle, and improved with manual lymphatic drainage immediately after surgery.

Surgical Anesthesia: Both general and local anesthesia involve risks. There is the possibility of complications, injury, and even death from all forms of surgical anesthesia or sedation.

Pain: You will experience pain during the application of local anesthesia and after liposuction, which will vary in intensity and duration, as well as itching, which may persist for a long time. Chronic pain may occur rarely from nerves becoming trapped in scar tissue. This process can be improved with manual lymphatic drainage immediately after surgery.

Slow Healing: Some areas may not heal normally, and they can take a long time to heal.

Allergic Reactions: In rare cases, allergic reactions to dressings or bandages, suture material and adhesives, topical preparations, or injectable agents have been reported. Serious systemic reactions, including shock (anaphylaxis), may occur due to drugs used during the liposuction and prescription medicine. Allergic reactions may require additional treatment.

Fat Necrosis: Deep fatty tissue of the skin may die. Additional surgery may be necessary to remove the areas of fat necrosis. The possibility exists of skin-contour irregularities that may be caused by fat necrosis.

Persistent inflammation (lymphedema): After liposuction, persistent inflammation in the legs may occur. This process can be improved with manual lymphatic drainage immediately after surgery.

Surgical shock: In rare circumstances, liposuction can cause severe trauma, particularly when multiple or extensive areas are suctioned at one time. For patients considering a big-volume liposuction, over 5,000 cc, a post-operative control and care, including an overnight stay at the hospital, is recommended.

Deep vein thrombosis, Cardiac and Pulmonary Complications: If a big-volume liposuction is expected, fat embolism syndrome, or DVT, should be prevented, as well as tumescent fluid overload. Inactivity and other conditions may increase the incidence of blood clots travelling to the lungs, causing a major blood clot that may result in death. Tell your physician about any past history of blood clots, swollen legs, or the use of estrogens or contraceptive pills that may contribute to this condition. If you experience shortness of breath, chest pain or unusual heartbeats, seek medical attention immediately. Should any of these complications occur, you may require hospitalization and additional treatment.

LASER-Assisted Lipectomy: Risks associated with the use of this technique include the risks mentioned above and the following specific risks:

Burns: LASER energy may produce burns and tissue damage either at the location where the cannula is inserted into the skin or in other areas if the cannula touches the undersurface of the skin for prolonged periods of time. In case a burn occurs, additional treatment and surgery may be necessary.

Smoking, Second-Hand Smoke Exposure, Nicotine Products (Patch, Gum, Nasal Spray): Patients who currently smoke or are exposed to second-hand smoke, use tobacco or nicotine products (patch, gum or nasal spray) are at a greater risk for significant surgical complications of skin dying, slow healing and additional scarring. Additionally, smokers may have a significant negative effect on anesthesia and recovery from anesthesia, with coughing and possibly increased bleeding.

Metabolic state of patients who have lost a lot of weight: Your personal metabolic state of blood chemistry and protein levels may be abnormal following massive weight loss and surgical procedures causing the patient to lose weight. Individuals with abnormalities may be at risk of serious medical and surgical complications, including slow wound healing or infection.

Long-term Results: Subsequent alterations in your body's appearance may occur as the result of aging, sun exposure, weight loss, weight gain, pregnancy, menopause and other circumstances not related to liposuction.

Female Patient Information: It is important to inform your doctor if you use birth control pills, replacement therapy with estrogens, or if you believe you may be pregnant. Many medications, including antibiotics, may neutralize the preventive effect of birth control pills, allowing for conception and pregnancy.

Body piercing procedures: Individuals planning to undergo a body piercing procedure have more chances of suffering an infection after liposuction. Do not undergo said procedure until your doctor's approval.

Mental health disorders and elective surgery: It is important that all patients seeking to undergo elective surgery have realistic expectations that focus on improvement rather than perfection. Complications or less-than-satisfactory results are sometimes unavoidable, may require additional surgery and are often stressful. Prior to surgery, openly discuss with your surgeon any history that you may have of significant emotional depression or mental health disorders. Although many individuals may benefit psychologically from the results of elective surgery, effects on mental health cannot be accurately predicted.

Medications: There are many adverse reactions that occur as a result of taking over the counter, herbal, and/or prescription medications. If you have an adverse reaction, discontinue the drugs immediately and call your doctor for further instructions. If the reaction is severe, go immediately to the nearest emergency room. Be sure to take your prescribed medication only as directed.

I have been correctly informed, even in writing (information documents) and/or using images, about this treatment's characteristics: its fundamentals, the manner and details of its execution, its action mechanisms, its immediate effects, the process and evolution to observe the necessary supplementary treatments in the following days, weeks or months, the cares and precautions to adopt in the following hours or days, the variability of the time necessary to fully recover; therefore, I acknowledge that no guarantee can be given to me as to the date when I can resume my regular activities (emotional, social, work and sports).

I ATTEST I have not left out or altered any data when showing my surgical medical history and records, specially those concerning allergies and diseases or personal risks.

The Medical History and results from tests performed on the patient don't advise against using the appropriate medical procedure when there isn't, a priori, any risks that may have determined the failure of the technique objectively considered, and which means to meet your expectations.

I RECOGNIZE that, during the treatment, unforeseen conditions may necessitate different procedures than those described above. Therefore, I hereby expressly authorize said procedures, including my transfer to a hospital center. In case of complications during the intervention, I authorize the Hospital to request the assistance they deem necessary from other specialists, in their best professional judgment.

If, during treatment, the healthcare staff was accidentally exposed to my blood, according to the principles of good medical practice I have been informed of the need to perform the proper analytical determinations on my blood, related to potential pathologies of blood transmission, to adopt the most suitable preventive measures.

I AGREE to do my best to follow the doctor's instructions to the letter before, during and after the treatment mentioned above. I take responsibility for complying with post treatment measures recommended by the Hospital, as well as attending control visits suggested by the doctor. It is important that surgical incisions are not subjected to excessive force, swelling, abrasion, or motion during the healing period. Protective dressings and drains should not be removed unless instructed by your doctor. Successful post-operative function depends on both

the surgery and subsequent care. Any physical activity that increases your pulse or heart rate may cause additional bruising, swelling, fluid accumulation and the need to undergo new treatment. It is wise to refrain from intimate physical activities after surgery until your physician states it is safe. It is important that you participate in follow-up care, that you return for aftercare, and that you promote your recovery after surgery with ultrasounds, HE-Ne laser, and especially with manual lymphatic drainage immediately after surgery. You should also attend regular visits until your final medical discharge (approximately 1 year)

I UNDERSTAND that the goal of the treatment is to improve my appearance, with the possibility that some imperfection remains and that the result may not be what I expected. In this sense, I have been informed that the aesthetic result of the treatment depends on factors like how easily you scar, and the formation or non-formation of keloids. I am aware that the medical field is not an exact science, and that absolute perfection cannot be guaranteed. I understand the result may not be as expected and acknowledge that I haven't been given any guarantee at all.

Medical action might not end in itself, and subsequent treatment action might be necessary to reach the goal you're consenting to.

The cost of the treatment includes several charges for the services provided. The total includes your doctor's fees and the cost of materials, equipment or product. The charges for this procedure don't include possible future costs for additional procedures of your choice, or that you require to review, optimize or complete your result. There may be additional charges due to complications arising from the treatment. Charges for secondary treatments concerning check-ups and charges that haven't been previously arranged will also be your responsibility. By signing the consent for this procedure, you acknowledge you have been informed about its risks and consequences and take responsibility for the medical decisions made, together with the economical costs of any future treatments.

I AUTHORIZE the taking of photographs and videos for an appropriate diagnostic evaluation, and for the control of the assessment and results. I also authorize the taking of photographs of the treated area for scientific, educational or clinical purposes, and I understand that their use is not a violation of privacy or confidentiality, rights to which I am entitled.

I KNOW my data will be treated in an automated manner, which I have authorized after my rights have been explained to me pursuant to the LOPD (Spanish Organic Act on Data Protection) in force: rights of access, rectification, cancellation and objection established by the Act, and after expressing I can address the Clinic in writing, notwithstanding the provisions of Section 8 of the Act about health-related data. With my signature, I give my express consent for my personal data to be handled by the CLINIC for their own purposes.

I am aware that my signature and this consent don't assume any kind of waiver to future claims, both clinical and legal. I am also aware that I can withdraw my signature from this consent at any time prior to the execution of the treatment.

I have clarified any doubts about everything that has been mentioned above, and I fully understand this CONSENT DOCUMENT, standing by each and all of its points; by signing the document, I ratify and consent to the execution of the treatment.

Date and place

Doctor

Patient Signature

Legal Representative

I have also been informed, and I fully acknowledge that this consent can be revoked by the undersigned at any time, and, in this case, said revocation must be done in writing, being fully aware of the consequences of my decision.

Doctor

Patient Signature

Legal Representative