

Preliminary results


Alejandro Rodriguez-Garcia¹, Srikanth Ramaswamy^{1,2}

¹Neural Circuits Laboratory, Biosciences Institute, Faculty of Medical Sciences, Newcastle University NE2 4HH, United Kingdom


²Theoretical Sciences Visiting Program (TSVP), Okinawa Institute of Science and Technology Graduate University, Japan

Model


- **Task:** Reinforce firing of 2 neurons via delayed reward (Izhikevich, 2007)
- **Network architecture:** RNN (1000 neurons with 100 connections/neuron), 4:1 EI ratio, different spiking populations
- **Learning:** DA-STDP learning




Learning




- A reward is delivered with a delay of 1 to 3 seconds whenever pre- and post-synaptic spikes coincide within a 20 ms window.
- The reward triggers dopamine release, enhances the synaptic strength of the connection through DA-STDP learning




Spiking types




Regular spiking (RS)



Fast spiking (FS)




Bursting (BU)




Accommodation (AC)

Impact of inhibitory spiking types



- Considering FS inhibitory neurons enhances learning of the reinforced synapse
- Considering AC inhibitory neurons impairs learning more of the reinforced synapse

Impact of inhibitory spiking heterogeneity




- Including different proportions of spiking heterogeneity has an impact on learning at the synaptic level
- A combination of 70% of FS and 30% of AC inhibitory neurons exhibited the best learning enhancement


Incorporating spiking heterogeneity provides an additional layer for fine-tuning learning in neural systems

Impact of excitatory spiking heterogeneity

- We studied the effects of bursting activity in the network: 80% RS and 20% BU excitatory neurons
 - Bursting activity increases network instability when pre- and post- neurons are RS
 - Bursting activity prompts unstable learning when pre- and post- neurons are BU




pre- and post- RS neurons




pre- and post- BU neurons

Bursting effect at the synaptic level

- Bursting activity increases post-, pre- correlated spikes (within the 20 ms window) → More DA is released → The synapse is enhanced more rapidly




RS neurons



BU neurons



Impact of bursting




- We now consider all excitatory neurons as BU.
- Bursting activity seem to promote learning by enhancing flexibility.
- However, compared with the RS excitatory neurons it is unstable.

Neuromodulated spiking

- Neuromodulators may act as switches on spiking activity, helping to balance fast learning and stability
- DA** drives learning, **NE** enhances fast learning through BU spiking, **ACh** enhances stability through RS spiking



Neuromodulated spiking



- Modulating BU activity to RS activity could help control this instability, preserving the network state after learning and enabling both fast learning and long-term stability

Neuromodulators can help to enhance fast learning and stability by controlling the spiking activity at the synaptic level

Key takeaways

- 1** Incorporating spiking heterogeneity provides an additional layer for fine-tuning learning in neural systems
- 2** Neuromodulators can help to enhance fast learning and stability by controlling the spiking activity at the synaptic level

