





Photo may not be representative of a label-consistent bait placement





Mold Resistant • Moisture Resistant

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for First Aid, additional Precautionary Statements, and Directions for Use.

ACTIVE INGREDIENT:

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, Bushy-tailed woodrat, and Mexican woodrat.



NET WT: 4 lbs (1.81 kg)

Product Code: TD4024

FIRST AID	
If Swallowed:	Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person. Do not induce womiting unless told to do so by the poison control center or doctor. Avoid use of all oils.
If on Skin:	Wash with plenty of soap and water.
If in Eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.
HAVE LABEL WHEN OBTAINING TREATMENT ADVICE	

NOTE TO PHYSICIAN

If serum calcium levels are elevated, treatment with Calcitonin is effective in reducing calcium to normal levels. Continue monitoring serum calcium and treat as necessary for hypercalcemia (Reference, AMA Drug Evaluations, Third Edition (1977), Chapter 16, pp. 248-251).

TREATMENT FOR PET POISONING

If an animal eats this bait, call a veterinarian, or 1-877-854-2494 at once.

NOTE TO VETERINARIAN

Calcium Mobilizer Cholecalciferol: For animals ingesting bait, induce vomiting by using hydrogen peroxide or administration of activated charcoal with a cathartic. If clinical signs develop, treatment consisting of saline diuresis combined with the use of furosemide, corticosteroids, and phosphate binders are recommended. Calcitonin or pamidronate may be needed for animals that remain hypercalcemic despite symptomatic treatment.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin or on clothing.

All handlers, (including applicators) must wear shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof

USER SAFETY REQUIREMENTS: Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten the bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent exposure:



KILLS RATS, HOUSE MICE, AND MEADOW VOLES*

MOLD RESISTANT • MOISTURE RESISTANT

ACTIVE INGREDIENT: Cholecalciferol (CAS #67-97-0): 0.075%

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for First Aid, additional Precautionary Statements, and Directions for Use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Pesticide Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other

Pesticide Disposal: Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. [Plastic:] Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NET WEIGHT: 4 lbs (1.81 kg)

EPA REG. NO. 12455-106 EPA EST, NOs. 12455-WI-1k, 12455-WI-2p, 12455-WI-3W Superscript is first letter of the lot number.

Manufactured by:



DIRECTIONS FOR USE (Continued from other panel)

- 1. Store unused product out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted. units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to
- 3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

Note: Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

USE RESTRICTIONS: This product may only be used to control the following rodent pests in and around man-made structures: House mouse (Mus musculus), Norway rat (Rattus norvegicus), Roof rat (Rattus rattus), Cotton rat* (Sigmodon hispidus), Eastern harvest mouse* (Reithrodontomys humuli), Golden mouse* (Ochrotomys nuttalli), Polynesian rat* (Rattus exulans), Meadow vole* (Microtus pennsylvanicus), White-throated woodrat* (Neotoma albiqula), Southern plains woodrat* (Neotoma micropus), Bushy-tailed woodrat* (Neotoma cinerea) and Mexican woodrat* (Neotoma mexicana). This product must be used in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops. Burrow baiting with TERAD3® Blox is prohibited.

Do not place bait near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait.

Do not sell or distribute this product in individual containers holding less than 4 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats, house mice or meadow voles* will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in corners and concealed places, between floors and walls, or in locations where signs of rodents have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Rats: Place 2 to 8 blocks per placement (usually at intervals of 15- to 30- feet). Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are signs of recent feeding on bait by rats.

House Mice, Eastern Harvest Mice*, Golden Mice* and Meadow Voles*: Use 1 Block per placement. Space placements at intervals of 8- to 12-feet in infested areas. Larger placements, up to 2 Blocks, may be needed at points of extremely high house mouse or meadow vole* activity. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are signs of recent feeding on bait.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Using waterproof gloves, collect and properly dispose of all dead, exposed animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish

*Not permitted for use against the following species in California: Cotton rat. Eastern harvest mouse. Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, Bushy-tailed woodrat, and Mexican woodrat.

043024/04-21