



COLLEGE OF NURSING AND ALLIED HEALTH SCIENCE

NAME OF STUDENT: _____ DATE: _____

INTRAMUSCULAR INJECTION
Performance Checklist

Equipment:

- Medication administration report (MAR)
- Medication as prescribed
- Sterile 3-ml syringe and long bevel, 20 to 22 gauge
- Alcohol swab
- 2-inch needle
- Nonsterile gloves
- Sterile 2X2 gauze pad

PROCEDURE:

STEPS	1	2	3	4	5	Rationale
1. Check with the client and the chart for any known allergies.						
2. Wash hands.						
3. Follow the five rights.						
4. Prepare the medication from an ampule or vial;						
6. Add 0.1 to 0.2 ml of air to the syringe.						
7. Take medication to the client's room & place on a clean surface.						
8. Check the client's identification armband.						
9. Explain the procedure to the client.						
10. Place the client in an appropriate position to						

expose the site. • Deltoid: sitting position. • Ventro-gluteal: • Side-lying: flex the knee; pivot the leg forward from the hip about 20° so it can rest on the bed. • Supine: flex the knee on the injection side. • Prone: point toes inward toward each other to internally rotate the femur.						
11. Wash hands and don non-sterile gloves.						
12. Select and clean the site.						
13. Assess the client's skin for redness, scarring, breaks in the skin, and palpate for lumps or nodules						
14. Select an appropriate site using appropriate anatomic landmarks.						
15. Cleanse the site with an alcohol wipe using a firm circular motion; cleanse from inside to outside; allow alcohol to dry.						
Prepare for injection. 16. Remove the needle cap by pulling it straight off, and expel any air bubbles from the syringe.						
17. Pull the skin down or to one side (Z-track technique) with non-dominant hand.						
Administer the injection. 18. Deltoid: quickly insert the needle with a dart like motion at a 90° angle muscle. Ventro-gluteal: quickly insert the needle using a dart-like motion and steady pressure at a 90° angle to the iliac crest in the middle of the V .						

19. Aspirate by pulling back on the plunger and observe for blood.						
20. If blood appears, remove needle and discard in a sharps container.						
21. Inject medication slowly if there is no blood present. about 10 sec/ml.						
22. Wait 10 seconds after the medication had been injected, then smoothly withdraw the needle at the same angle of insertion.						
23. Apply gentle pressure at the site with a dry, sterile 2 × 2 gauze; do not massage the injection site. Swab using gentle pressure.						
24. Discard the needle and syringe in a sharps container; do not recap the needle.						
25. Position client for comfort; encourage client receiving ventro-gluteal injections to perform leg exercises (flexion and extension).						
26. Remove gloves, wash hands.						
27. Record on the MAR the dosage, route, site, and time.						

Evaluated by: _____