



NAME OF STUDENT: _____ DATE: _____

SUBCUTANEOUS INJECTION
Performance Checklist

Equipment:

- Medication administration report (MAR)
- Medication as prescribed
- Sterile 3-ml syringe and long bevel, 20 to 22 gauge
- Alcohol swab
- 2-inch needle
- Nonsterile gloves
- Sterile 2X2 gauze pad

PROCEDURE:

STEPS	1	2	3	4	5	Rationale
1. Check with the client and the chart for any known allergies.						
2. Wash hands.						
3. Follow the five rights.						
4. Prepare the medication from an ampule or vial;						
6. Add 0.1 to 0.2 ml of air to the syringe.						
7. Take medication to the client's room & place on a clean surface.						
8. Check the client's identification armband.						
9. Explain the procedure to the client.						
10. Place the client in an appropriate position to expose the site.						

● Abdomen: supine position.						
11. Wash hands and don non-sterile gloves.						
12. Select and clean the site (encircling the umbilicus.)						
13. Assess the client’s skin for redness, scarring, breaks in the skin, and palpate for lumps or nodules						
14. Select an appropriate site using appropriate anatomic landmarks.						
15. Cleanse the site with an alcohol wipe using a firm circular motion; cleanse from inside to outside; allow alcohol to dry.						
Prepare for injection. 16. Remove the needle cap by pulling it straight off, and expel any air bubbles from the syringe.						
17. Stretch the skin to side with non-dominant hand.						
18. Quickly insert the needle with a dart like motion at a 45° angle muscle.						
19. Aspirate by pulling back on the plunger, and observe for blood.						
20. If blood appears, remove needle and discard in a sharps container.						
21. Inject medication slowly if there is no blood present. about 10 sec/ml.						
22. Wait 10 seconds after the medication had been injected, then smoothly withdraw the needle at the same angle of insertion with a dry cotton ball applied at the site.						

23. Apply gentle pressure at the site with a dry, sterile 2 × 2 gauze; do not massage the injection site. Swab using gentle pressure.						
24. Discard the needle and syringe in a sharps container; do not recap the needle.						
25. Position client for comfort.						
26. Remove gloves, wash hands.						
27. Record on the MAR the dosage, route, site, and time.						

Evaluated by: _____