



NAME OF STUDENT: _____ DATE: _____

WITHDRAWING FROM AN AMPULE
Performance Checklist

Equipment:

- Medical administration record (MAR)
- Ampule of prescribed medication
- Sterile syringe and needle
- Sterile gauze or alcohol swab
- Extra needle of proper gauge and length in filter needle accord with site

PROCEDURE:

STEPS	1	2	3	4	5	Rationale
1. Do handwashing.						
2. Hold the ampule and quickly and lightly tap the top chamber until all fluid flows into the bottom chamber.						
3. Place a sterile gauze or alcohol wipe around the neck of the ampule.						
4. Firmly grasp the neck of the ampule and quickly snap the top off away from your body. Place the ampule on a flat surface.						
5. Withdraw the medication from the ampule, maintaining sterile technique.						
6. Check connection of needle to syringe by turning barrel to right while holding needle guard.						
7. Remove needle guard, and hold syringe in dominant hand.						
8. With nondominant hand grasp ampule and turn upside down, or stabilize ampule on a flat surface.						
9. Insert the needle into the center of the ampule; do not allow the needle tip or shaft to touch the rim of the ampule.						
10. Keep needle tip below level of meniscus.						
11. Aspirate the medication by gently pulling on the plunger.						

12. If air bubbles are aspirated, remove the needle from the ampule. Hold syringe with needle pointing up and tap sides of the syringe. Draw back slightly on plunger, and gently push the plunger upward to eject air. Reinsert the needle in the middle of the ampule and continue to withdraw the medication.						
13. Remove excess air from the syringe and check the dosage of medication in the syringe. Recap.						
14. Discard any unused portion of the medication and dispose of the ampule top in a suitable container after comparing with MAR.						
15. Change needle and properly discard used needle. Secure needle to syringe by turning the barrel to right while holding the needle guard.						
16. Wash hands.						

Evaluated by:



NAME OF STUDENT: _____ DATE: _____

WITHDRAWING FROM A VIAL
Performance Checklist

Equipment:

- Medication administration record (MAR)
- Vial of medication
- Sterile syringe and needle
- Sterile needle
- Alcohol swab

PROCEDURE:

STEPS	1	2	3	4	5	Rationale
1. Do handwashing.						
Prepare the vial. 2. Open the alcohol wipe.						
3. For a new vial, remove metal cap from vial of medicine and cleanse the rubber top of the vial. For used vial, cleanse the rubber top of the vial.						
Prepare syringe. 4. Choose a syringe of appropriate size to accommodate the volume of medication to be withdrawn.						
5. Grasp needle and turn barrel of syringe to the right.						
6. Remove the needle cap and pull back on plunger to fill syringe with an amount of air equal to amount of solution to be withdrawn from the vial.						
7. Insert the needle into the center of the upright vial and inject air into the vial.						
9. Invert vial; keep the vial at eye level and the						

needle's bevel below the fluid level, and remove the exact amount of medicine while touching only the syringe barrel and plunger tip						
10. Expel air from the syringe while needle remains within the inverted vial by tapping the side of the syringe with your finger.						
11. Check the amount of medicine in the syringe.						
12. Turn vial upright and remove the needle.						
13. Replace the needle cap. Open the sterile package of the new needle. Remove used needle, and dispose in the sharps container.						
14. Attach the new needle to the syringe by turning the barrel to the right.						
15. Compare the medication in the syringe with the prescribed dosage.						
16. Wash hands.						

- Level of Performance:
- 5 - Excellent (E)
 - 4 - Very Satisfactory (VS)
 - 3 - Satisfactory (S)
 - 2 - Poor (P)
 - 1 - Needs Improvement (NI)

Evaluated by:



NAME OF STUDENT: _____ DATE: _____

INTRAMUSCULAR INJECTION
Performance Checklist

Equipment:

- Medication administration report (MAR)
- Medication as prescribed
- Sterile 3-ml syringe and long bevel, 20 to 22 gauge
- Alcohol swab
- 2-inch needle
- Nonsterile gloves
- Sterile 2X2 gauze pad

PROCEDURE:

STEPS	1	2	3	4	5	Rationale
1. Check with the client and the chart for any known allergies.						
2. Wash hands.						
3. Follow the five rights.						
4. Prepare the medication from an ampule or vial;						
6. Add 0.1 to 0.2 ml of air to the syringe.						
7. Take medication to the client's room & place on a clean surface.						
8. Check the client's identification armband.						
9. Explain the procedure to the client.						
10. Place the client in an appropriate position to expose the site. • Deltoid: sitting position. • Ventro-gluteal: • Side-lying: flex the knee; pivot the leg forward from the hip about 20° so it can						

rest on the bed. ● Supine: flex the knee on the injection side. ● Prone: point toes inward toward each other to internally rotate the femur.						
11. Wash hands and don non-sterile gloves.						
12. Select and clean the site.						
13. Assess the client’s skin for redness, scarring, breaks in the skin, and palpate for lumps or nodules						
14. Select an appropriate site using appropriate anatomic landmarks.						
15. Cleanse the site with an alcohol wipe using a firm circular motion; cleanse from inside to outside; allow alcohol to dry.						
Prepare for injection. 16. Remove the needle cap by pulling it straight off, and expel any air bubbles from the syringe.						
17. Pull the skin down or to one side (Z-track technique) with non-dominant hand.						
Administer the injection. 18. Deltoid: quickly insert the needle with a dart like motion at a 90° angle muscle. Ventro-gluteal: quickly insert the needle using a dart-like motion and steady pressure at a 90° angle to the iliac crest in the middle of the V .						
19. Aspirate by pulling back on the plunger and observe for blood.						
20. If blood appears, remove needle and discard in a sharps container.						

21. Inject medication slowly if there is no blood present. about 10 sec/ml.						
22. Wait 10 seconds after the medication had been injected, then smoothly withdraw the needle at the same angle of insertion.						
23. Apply gentle pressure at the site with a dry, sterile 2 × 2 gauze; do not massage the injection site. Swab using gentle pressure.						
24. Discard the needle and syringe in a sharps container; do not recap the needle.						
25. Position client for comfort; encourage client receiving ventro-gluteal injections to perform leg exercises (flexion and extension).						
26. Remove gloves, wash hands.						
27. Record on the MAR the dosage, route, site, and time.						

Evaluated by: _____



COLLEGE OF NURSING AND ALLIED HEALTH SCIENCE

NAME OF STUDENT: _____ DATE: _____

SUBCUTANEOUS INJECTION
Performance Checklist

Equipment:

- Medication administration report (MAR)
- Medication as prescribed
- Sterile 3-ml syringe and long bevel, 20 to 22 gauge
- Alcohol swab
- to 2-inch needle
- Nonsterile gloves
- Sterile 2X2 gauze pad

PROCEDURE:

STEPS	1	2	3	4	5	Rationale
1. Check with the client and the chart for any known allergies.						
2. Wash hands.						
3. Follow the five rights.						
4. Prepare the medication from an ampule or vial;						
6. Add 0.1 to 0.2 ml of air to the syringe.						
7. Take medication to the client's room & place on a clean surface.						
8. Check the client's identification armband.						
9. Explain the procedure to the client.						
10. Place the client in an appropriate position to expose the site. • Abdomen: supine position.						
11. Wash hands and don non-sterile gloves.						

12. Select and clean the site (encircling the umbilicus.)						
13. Assess the client’s skin for redness, scarring, breaks in the skin, and palpate for lumps or nodules						
14. Select an appropriate site using appropriate anatomic landmarks.						
15. Cleanse the site with an alcohol wipe using a firm circular motion; cleanse from inside to outside; allow alcohol to dry.						
Prepare for injection. 16. Remove the needle cap by pulling it straight off, and expel any air bubbles from the syringe.						
17. Stretch the skin to side with non-dominant hand.						
18. Quickly insert the needle with a dart like motion at a 45° angle muscle.						
19. Aspirate by pulling back on the plunger, and observe for blood.						
20. If blood appears, remove needle and discard in a sharps container.						
21. Inject medication slowly if there is no blood present. about 10 sec/ml.						
22. Wait 10 seconds after the medication had been injected, then smoothly withdraw the needle at the same angle of insertion with a dry cotton ball applied at the site.						
23. Apply gentle pressure at the site with a dry, sterile 2 × 2 gauze; do not massage the injection site. Swab using gentle pressure.						
24. Discard the needle and syringe in a sharps container; do not recap the needle.						
25. Position client for comfort.						

26. Remove gloves, wash hands.							
27. Record on the MAR the dosage, route, site, and time.							

Evaluated by: _____



NAME OF STUDENT: _____ **DATE:** _____

INTRADERMAL INJECTION
Performance Checklist

Equipment:

- Medication administration record (MAR)
- Medication
- Medicine ticket
- Sterile water for injection
- Sterile tuberculin syringe and short bevel, 25-27 gauge, 3/8- to 1/2-inch needle.
- Disposable gloves
- Alcohol swab
- sterile 2 × 2 gauze pad

PROCEDURE:

STEPS	1	2	3	4	5	Rationale
1. Check with the client and the chart for any known allergies.						
2. Wash hands.						
3. Follow the five rights.						
4. Prepare the medication from an ampule or vial. Withdraw 0.9 cc sterile water for injection and add 0.1 cc from the ordered medicine. Compare to medicine ticket.						
5. Place the prepared medication with the medicine ticket on a tray and bring to the client's room and place on a clean surface.						
6. Ask the Client to state his/her name and compare to the client's identification armband.						
7. Explain the procedure to the client.						
8. Place the client in a comfortable position; Provide Privacy						

9. Wash hands and don non-sterile gloves.						
10. Select and clean the site.						
11. Assess the client’s skin for bruises, redness, or broken tissue.						
12. Select an appropriate site using appropriate anatomic landmarks.						
13. Cleanse the site with an alcohol wipe using a firm circular motion; cleanse from inside to outside; allow alcohol to dry.						
Prepare the syringe for injection. 14. Remove the needle guard.						
15. Check the amount of solution in the syringe.						
16. Inject the medication.						
17. Hold the syringe in dominant hand.						
18. With non-dominant hand, grasp the client’s dorsal forearm and gently pull the skin taut on ventral forearm.						
19. Administer the medication slowly; observe the development of a bleb (large flaccid vesicle that resembles a mosquito bite). If none appears, withdraw the needle slightly.						
20. Withdraw the needle.						
21. Pat area gently with a dry 2 × 2 sterile gauze pad/ dry cotton ball.						
22. Remove gloves, dispose into appropriate receptacle, and wash hands.						

23. Draw a circle around the perimeter of the bleb with a ballpoint pen and write the due time. Instruct the Client not to scratch the site.						
24. Observe for signs of an allergic reaction. Inform the Physician of the due time for reading and document the result.						
25. Document medication and site of injection on the MAR and medicine ticket.						

Evaluated by: _____