



NAME OF STUDENT: _____ DATE: _____

WITHDRAWING FROM A VIAL
Performance Checklist

Equipment:

- Medication administration record (MAR)
- Vial of medication
- Sterile syringe and needle
- Sterile needle
- Alcohol swab

PROCEDURE:

STEPS	1	2	3	4	5	Rationale
1. Do handwashing.						
Prepare the vial. 2. Open the alcohol wipe.						
3. For a new vial, remove metal cap from vial of medicine and cleanse the rubber top of the vial. For used vial, cleanse the rubber top of the vial.						
Prepare syringe. 4. Choose a syringe of appropriate size to accommodate the volume of medication to be withdrawn.						
5. Grasp needle and turn barrel of syringe to the right.						
6. Remove the needle cap and pull back on plunger to fill syringe with an amount of air equal to amount of solution to be withdrawn from the vial.						
7. Insert the needle into the center of the upright vial and inject air into the vial.						
9. Invert vial; keep the vial at eye level and the						

needle's bevel below the fluid level, and remove the exact amount of medicine while touching only the syringe barrel and plunger tip						
10. Expel air from the syringe while needle remains within the inverted vial by tapping the side of the syringe with your finger.						
11. Check the amount of medicine in the syringe.						
12. Turn vial upright and remove the needle.						
13. Replace the needle cap. Open the sterile package of the new needle. Remove used needle, and dispose in the sharps container.						
14. Attach the new needle to the syringe by turning the barrel to the right.						
15. Compare the medication in the syringe with the prescribed dosage.						
16. Wash hands.						

- Level of Performance:
- 5 - Excellent (E)
 - 4 - Very Satisfactory (VS)
 - 3 - Satisfactory (S)
 - 2 - Poor (P)
 - 1 - Needs Improvement (NI)

Evaluated by: _____