

Assignment 1

Alen Mehmedbegovic

January 13th 2024

1/ NAND gate pretty much opposite of AND gate

1i/ NOT as NAND & fanout

X	NOT(X)
0	1
1	0

Want this with NAND & fanout

$$\Rightarrow \text{NOT}(X) \equiv \text{NAND}(\text{fanout}(X))$$

X	NAND(fanout(X))
0	1
1	0

1ii/ WTS: From $\{ \text{NAND}, \text{fanout} \}$, we can construct $\{ \text{AND}, \text{OR}, \text{NOT}, \text{fanouts} \}$

From 1i/ we showed that $\text{NOT}(X) \equiv \text{NAND}(\text{fanout}(X))$. Now we can use NOT in our implementation to mean $\text{NAND}(\text{fanout}(X))$.

So NOT is covered ✓ And fanout is part of original set ✓

Now let's make OR.

X	Y	OR(X,Y)
0	0	0
0	1	1
1	0	1
1	1	1

$$\Rightarrow$$

X	Y	NAND(NOT(X), NOT(Y))
0	0	0
0	1	1
1	0	1
1	1	1

$$\Rightarrow \text{OR}(X,Y) \equiv \text{NAND}(\text{NOT}(X), \text{NOT}(Y))$$

Finally $\text{AND}(X,Y) \equiv \text{NAND}(\text{NOT}(X), \text{NOT}(Y))$ ✓ $\Rightarrow \{ \text{NAND}, \text{NOT} \}$ is universal

$$2) \text{ let } |0\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, |1\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, |2\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

$$2.i) |\psi\rangle = \begin{pmatrix} \frac{1}{\sqrt{3}} \\ \frac{i}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} \end{pmatrix}, |\phi\rangle = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} \end{pmatrix}$$

$$2.ii) \langle \psi | \psi \rangle = \left(\frac{1}{\sqrt{3}} \quad \frac{-i}{\sqrt{3}} \quad \frac{1}{\sqrt{3}} \right) \begin{pmatrix} \frac{1}{\sqrt{3}} \\ \frac{i}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} \end{pmatrix} = \frac{1}{3} + \frac{1}{3} + \frac{1}{3} = 1$$

$$\langle \phi | \phi \rangle = \left(0 \quad \frac{1}{\sqrt{2}} \quad \frac{i}{\sqrt{2}} \right) \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} \end{pmatrix} = 0 + \frac{1}{2} + \frac{1}{2} = 1$$

$$\langle \psi | \phi \rangle = \left(\frac{1}{\sqrt{3}} \quad \frac{-i}{\sqrt{3}} \quad \frac{1}{\sqrt{3}} \right) \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} \end{pmatrix} = \frac{-i}{\sqrt{6}} - \frac{i}{\sqrt{6}} = \frac{-2i}{\sqrt{6}} = -\frac{\sqrt{2}}{3} i$$

$$|\psi\rangle\langle\phi| = \begin{pmatrix} \frac{1}{\sqrt{3}} \\ \frac{i}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} \end{pmatrix} \left(0 \quad \frac{1}{\sqrt{2}} \quad \frac{i}{\sqrt{2}} \right) = \begin{pmatrix} 0 & \frac{1}{\sqrt{6}} & \frac{i}{\sqrt{6}} \\ 0 & \frac{i}{\sqrt{6}} & -\frac{1}{\sqrt{6}} \\ 0 & -\frac{1}{\sqrt{6}} & -\frac{i}{\sqrt{6}} \end{pmatrix}$$

$$|\psi\rangle \otimes |\phi\rangle = \begin{pmatrix} \frac{1}{\sqrt{3}} \\ \frac{i}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} \end{pmatrix} \otimes \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} \end{pmatrix}$$

2.ii)

$$|\psi\rangle \otimes |\phi\rangle =$$

$$\begin{pmatrix} 0 \\ \frac{1}{\sqrt{6}} \\ \frac{i}{\sqrt{6}} \\ 0 \\ \frac{1}{\sqrt{6}} \\ 0 \\ -\frac{1}{\sqrt{6}} \\ \frac{i}{\sqrt{6}} \end{pmatrix}$$

$$2.iii) |\psi\rangle + |\phi\rangle = \begin{pmatrix} \frac{1}{\sqrt{3}} \\ \frac{i}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} \end{pmatrix} + \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{3}} \\ \frac{i}{\sqrt{3}} + \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{3}} + \frac{i}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{3}} \\ \frac{\sqrt{2}i + \sqrt{3}}{\sqrt{6}} \\ \frac{-\sqrt{2} - \sqrt{3}i}{\sqrt{6}} \end{pmatrix}$$

$$||\psi\rangle + |\phi\rangle|| = \sqrt{\left|\frac{1}{\sqrt{3}}\right|^2 + \left|\frac{\sqrt{2}i + \sqrt{3}}{\sqrt{6}}\right|^2 + \left|\frac{-\sqrt{2} - \sqrt{3}i}{\sqrt{6}}\right|^2}$$

$$= \sqrt{\frac{1}{3} + \frac{-2 + 2\sqrt{6}i + 3}{6} + \frac{2 + 2\sqrt{6}i - 3}{6}}$$

$$= \sqrt{\frac{2 + 4\sqrt{6}i - 1}{6}}$$

$$= \sqrt{\frac{1 + 2\sqrt{3}i}{6}}$$

2iii) $\| |\psi\rangle + |\phi\rangle \| = \frac{\sqrt{1+2\sqrt{3}i}}{\sqrt{3}} \neq 1$, it's not a unit vector

To make it a unit vector, divide each component with its magnitude

$$\frac{|\psi\rangle + |\phi\rangle}{\| |\psi\rangle + |\phi\rangle \|} = \begin{pmatrix} \frac{\sqrt{1+2\sqrt{3}i}}{3} \\ \frac{(\sqrt{2}i + \sqrt{3}) \cdot \sqrt{1+2\sqrt{3}i}}{\sqrt{18}} \\ \frac{(-\sqrt{2} - \sqrt{3}i) \cdot \sqrt{1+2\sqrt{3}i}}{\sqrt{18}} \end{pmatrix} \quad \checkmark$$

3i) $|\psi_3\rangle = \begin{pmatrix} \frac{1}{\sqrt{2}} \\ e^{i\theta} \\ \frac{1}{\sqrt{2}} \end{pmatrix} \quad \theta \in \mathbb{R}$

3ii) $|0\rangle \rightarrow \left| \frac{1}{\sqrt{2}} \right|^2 = \frac{1}{2} \quad |1\rangle \rightarrow \left| \frac{e^{i\theta}}{\sqrt{2}} \right|^2 = \frac{|e^{2i\theta}|}{2}$

3iii) $\frac{1}{\sqrt{2}} \begin{vmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{vmatrix} \begin{pmatrix} \frac{1}{\sqrt{2}} \\ e^{i\theta} \\ \frac{1}{\sqrt{2}} \end{pmatrix} = \begin{vmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{vmatrix} \begin{pmatrix} \frac{1}{\sqrt{2}} \\ e^{i\theta} \\ \frac{1}{\sqrt{2}} \end{pmatrix} = \begin{vmatrix} \frac{1}{2} + \frac{e^{i\theta}}{2} \\ \frac{1}{2} - \frac{e^{i\theta}}{2} \end{vmatrix} = \begin{pmatrix} \frac{1+e^{i\theta}}{2} \\ \frac{1-e^{i\theta}}{2} \end{pmatrix}$

$|0\rangle \rightarrow \left| \frac{1+e^{i\theta}}{2} \right|^2 = \frac{|1+2e^{i\theta}+e^{2i\theta}|}{4} = \frac{1}{4} + \frac{|2e^{i\theta}+e^{2i\theta}|}{4}$

$|1\rangle \rightarrow \left| \frac{1-e^{i\theta}}{2} \right|^2 = \frac{|1-2e^{i\theta}+e^{2i\theta}|}{4} = \frac{1}{4} + \frac{|-2e^{i\theta}+e^{2i\theta}|}{4}$

$$4ii) Y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, Y|+\rangle = |+\rangle. \text{ let } |+\rangle = \begin{pmatrix} a \\ b \end{pmatrix}$$

$$\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} -ib \\ ia \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}, a = -ib \text{ \& } b = ia$$

$$\Rightarrow b = i \text{ \& } a = 1 \text{ as } a = -ib, 1 = -i \cdot i \text{ \& } b = ia, i = i \cdot 1 \text{ both true}$$

$$\rightarrow |+\rangle = \begin{pmatrix} 1 \\ i \end{pmatrix} \checkmark$$

$$\text{Now let } |-\rangle = \begin{pmatrix} a \\ b \end{pmatrix}$$

$$\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = - \begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} -ib \\ ia \end{pmatrix} = \begin{pmatrix} -a \\ -b \end{pmatrix}, a = ib \text{ \& } b = -ia$$

$$\Rightarrow b = 1 \text{ \& } a = i \text{ as } a = ib, i = i \cdot 1 \text{ \& } b = -i \cdot a, 1 = -i \cdot i \text{ both true}$$

$$\rightarrow |-\rangle = \begin{pmatrix} i \\ 1 \end{pmatrix} \checkmark$$

$$4iii) U = \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix}, UU^\dagger = \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix} \begin{pmatrix} 1 & -i \\ -i & 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} \neq I$$

$$UU^\dagger \neq I \Rightarrow \text{not unitary}$$

$$iii) \begin{pmatrix} 1 & -i \\ -i & 1 \end{pmatrix} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix} = \begin{pmatrix} 1 & -i \\ -i & 1 \end{pmatrix} \begin{pmatrix} 1 & -i \\ i & -1 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix}$$