
Adipogenesis


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.41 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Allograft rejection


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.37 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Androgen response


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.37 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Angiogenesis


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.86 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Apical junction


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.86 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Apical surface


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.79 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Apoptosis


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.49 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Bile acid metabolism


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.95 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Cholesterol homeostasis


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.15 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Coagulation


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Complement


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.62 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

DNA repair


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

E2F targets


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Epithelial mesenchymal transition


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.83 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Estrogen response early


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.94 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Estrogen response late


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.93 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Fatty acid metabolism


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.60 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

G2M checkpoint


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Glycolysis


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.68 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Hedgehog signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.69 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Heme metabolism


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.50 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Hypoxia


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.83 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Il2 stat5 signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

IL6 JAK STAT3 signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.53 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Inflammatory response


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.61 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Interferon alpha response


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.45 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Interferon gamma response


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.52 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

KRAS signaling DN


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.72 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

KRAS signaling UP


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.94 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Mitotic spindle


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

MTORC1 signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.34 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

MYC targets v1


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.50 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

MYC targets v2


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.28 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Myogenesis


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.91 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Notch signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.21 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Oxidative phosphorylation


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.45 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

P53 pathway


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.46 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Pancreas beta cells


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 1.03 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Peroxisome


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.43 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

PI3K AKT mTOR signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.34 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Protein secretion


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Reactive oxygen species pathway


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Spermatogenesis


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.55 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

TGF beta signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.58 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

TNFα signaling via NFκB


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.57 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Unfolded protein response


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.43 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

UV response DN


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

UV response UP


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.66 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

WNT beta catenin signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.31 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Xenobiotic metabolism

202 children (PROTECT) · 610 adults (exploratory four-study summary)



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.79 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.