
Adipogenesis


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 102/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.41 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Allograft rejection


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 151/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.37 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Androgen response


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 42/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.37 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Angiogenesis


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 243/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.86 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Apical junction


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 131/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.86 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Apical surface


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 237/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.79 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Apoptosis


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 88/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.49 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Bile acid metabolism


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 225/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.95 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Cholesterol homeostasis


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 151/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.15 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Coagulation


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 126/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Complement


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 95/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.62 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

DNA repair


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 58/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

E2F targets


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 48/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Epithelial mesenchymal transition


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 256/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.83 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Estrogen response early


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 245/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.94 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Estrogen response late


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 240/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.93 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Fatty acid metabolism


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 151/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.60 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

G2M checkpoint


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 55/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Glycolysis


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 192/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.68 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Hedgehog signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 199/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.69 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Heme metabolism


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 155/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.50 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Hypoxia


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 135/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.83 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Il2 stat5 signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 150/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

IL6 JAK STAT3 signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 173/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.53 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Inflammatory response


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 226/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.61 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Interferon alpha response


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 80/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.45 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Interferon gamma response


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 127/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.52 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

KRAS signaling DN


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 153/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.72 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

KRAS signaling UP


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 171/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.94 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Mitotic spindle


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 93/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

MTORC1 signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 113/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.34 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

MYC targets v1


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 16/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.50 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

MYC targets v2


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 20/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.28 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Myogenesis


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 180/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.91 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Notch signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 191/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.21 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Oxidative phosphorylation


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 92/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.45 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

P53 pathway


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 109/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.46 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Pancreas beta cells


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 194/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 1.03 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Peroxisome


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 126/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.43 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

PI3K AKT mTOR signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 143/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.34 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Protein secretion


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 107/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Reactive oxygen species pathway


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 80/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Spermatogenesis


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 233/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.55 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

TGF beta signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 95/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.58 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

TNFα signaling via NFκB


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 174/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.57 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Unfolded protein response


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 35/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.43 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

UV response DN


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 159/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

UV response UP


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 155/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.66 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

WNT beta catenin signaling


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 131/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.31 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study




Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.

Xenobiotic metabolism


202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 143/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.79 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Exploratory age-bin evidence · separate fits within each study



Segments mark whole age bins; 60+ continues beyond the axis. Blank = no patients. Open circles do not establish equal scores.
Bin q-values use one separate FDR family across all pathways, studies and testable bins. They are not tests of the curves above.