
Adipogenesis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.41 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Allograft rejection

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.37 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Androgen response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.37 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Angiogenesis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.86 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Apical junction

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.86 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Apical surface

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.79 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Apoptosis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.49 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Bile acid metabolism

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.95 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Cholesterol homeostasis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.15 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Coagulation

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Complement

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.62 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

DNA repair

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

E2F targets

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Epithelial mesenchymal transition

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.83 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Estrogen response early

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.94 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Estrogen response late

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.93 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Fatty acid metabolism

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.60 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

G2M checkpoint

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Glycolysis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.68 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Hedgehog signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.69 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Heme metabolism

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.50 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Hypoxia

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.83 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Il2 stat5 signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

IL6 JAK STAT3 signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.53 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Inflammatory response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.61 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Interferon alpha response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.45 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Interferon gamma response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.52 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

KRAS signaling DN

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.72 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

KRAS signaling UP

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.94 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Mitotic spindle

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

MTORC1 signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.34 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

MYC targets v1

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.50 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

MYC targets v2

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.28 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Myogenesis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.91 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Notch signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.21 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Oxidative phosphorylation

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.45 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

P53 pathway

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.46 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Pancreas beta cells

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 1.03 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Peroxisome

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.43 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

PI3K AKT mTOR signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.34 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Protein secretion

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Reactive oxygen species pathway

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Spermatogenesis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.55 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

TGF beta signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.58 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

TNFα signaling via NFκB

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.57 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Unfolded protein response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.43 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

UV response DN

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

UV response UP

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.66 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

WNT beta catenin signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.31 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Xenobiotic metabolism

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Full patient distribution · every observed score is within the y limits



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.79 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.