
Adipogenesis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 102/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.41 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Allograft rejection

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 151/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.37 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Androgen response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 42/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.37 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Angiogenesis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 243/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.86 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Apical junction

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 131/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.86 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Apical surface

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 237/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.79 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Apoptosis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 88/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.49 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Bile acid metabolism

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 225/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.95 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Cholesterol homeostasis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 151/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.15 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Coagulation

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 126/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Complement

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 95/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.62 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

DNA repair

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 58/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

E2F targets

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 48/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Epithelial mesenchymal transition

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 256/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.83 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Estrogen response early

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 245/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.94 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Estrogen response late

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 240/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.93 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Fatty acid metabolism

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 151/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.60 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

G2M checkpoint

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 55/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Glycolysis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 192/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.68 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Hedgehog signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 199/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.69 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Heme metabolism

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 155/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.50 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Hypoxia

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 135/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.83 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Il2 stat5 signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 150/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

IL6 JAK STAT3 signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 173/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.53 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Inflammatory response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 226/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.61 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Interferon alpha response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 80/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.45 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Interferon gamma response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 127/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.52 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

KRAS signaling DN

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 153/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.72 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

KRAS signaling UP

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 171/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.94 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Mitotic spindle

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 93/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

MTORC1 signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 113/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.34 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

MYC targets v1

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 16/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.50 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

MYC targets v2

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 20/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.28 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Myogenesis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 180/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.91 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Notch signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 191/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.21 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Oxidative phosphorylation

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 92/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.45 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

P53 pathway

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 109/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.46 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Pancreas beta cells

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 194/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 1.03 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Peroxisome

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 126/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.43 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

PI3K AKT mTOR signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 143/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.34 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Protein secretion

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 107/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.36 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Reactive oxygen species pathway

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 80/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.44 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Spermatogenesis

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 233/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.55 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

TGF beta signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 95/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.58 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

TNFα signaling via NFκB

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 174/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.57 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Unfolded protein response

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 35/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.43 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

UV response DN

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 159/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.64 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

UV response UP

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 155/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.66 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

WNT beta catenin signaling

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 131/812 patient dots outside y limits; full distributions available in the explorer/PDF




Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.31 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.

Xenobiotic metabolism

202 children (PROTECT) · 610 adults (exploratory four-study summary)

Curve-focused view · 143/812 patient dots outside y limits; full distributions available in the explorer/PDF



Fixed study mix + shared adult age shapes; leaving one study out shifts group levels by up to 0.79 SD.
Dots: observed patient scores; lines: adjusted means. Whiskers: pointwise 95% model CIs. Grey gap: no participants.
Child and adult fits are separate; gaps stay unconnected. Cross-sectional age profiles, not within-person change.