TB–nonTB enrichment differences across supported ages

PROTECT
n = 202

ADTB
n = 359

GHL
n = 93

R2D2
n = 98

TBSQUARE
n = 60

Myogenesis
PI3K AKT mTOR signaling
Estrogen response late
DNA repair
Complement
Spermatogenesis
MYC targets v1
Fatty acid metabolism
G2M checkpoint
Protein secretion
Pancreas beta cells
E2F targets
Xenobiotic metabolism
Apical surface
Angiogenesis
Cholesterol homeostasis
MYC targets v2
Unfolded protein response
Heme metabolism
Hypoxia
Il2 stat5 signaling
Oxidative phosphorylation
UV response UP
Glycolysis
IL6 JAK STAT3 signaling
TGF beta signaling
Peroxisome
MTORC1 signaling
Mitotic spindle
Notch signaling
Apoptosis
Interferon alpha response
UV response DN
WNT beta catenin signaling
Androgen response
TNFα signaling via NFκB
Allograft rejection
KRAS signaling UP
Bile acid metabolism
Estrogen response early
Coagulation
Reactive oxygen species pathway
Hedgehog signaling
Apical junction
Interferon gamma response
Epithelial mesenchymal transition
Adipogenesis
P53 pathway
Inflammatory response
KRAS signaling DN

2 4 6 8 10 12 14
Age (years)

20 30 40 50 60 70 80
Age (years)

20 30 40 50 60
Age (years)

20 30 40 50 60 70 80
Age (years)

20 30 40 50 60
Age (years)

−1.5 −1.0 −0.5 0.0 0.5 1.0 1.5
TB − nonTB difference (pooled pathway-score SD)

All tested pathways shown; ordered by minimum interaction q, then p, across studies. Minimum age-interaction q = 0.986.
Blue: higher relative enrichment in TB. Orange: lower in TB. Grey: insufficient support. Color describes effect estimates; it does not mark significance.