
Adipogenesis across age


One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Age coverage: thick segments = supported comparisons; ticks = observed patients




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Allograft rejection across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed


Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Androgen response across age

One shared age axis · separate adjusted curves for each study




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Angiogenesis across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Apical junction across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Apical surface across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Apoptosis across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed


Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Bile acid metabolism across age


One shared age axis · separate adjusted curves for each study



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Cholesterol homeostasis across age

One shared age axis · separate adjusted curves for each study




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Coagulation across age


One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Age coverage: thick segments = supported comparisons; ticks = observed patients




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Complement across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed


Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

DNA repair across age

One shared age axis · separate adjusted curves for each study




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

E2F targets across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Epithelial mesenchymal transition across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Estrogen response early across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Estrogen response late across age


One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Age coverage: thick segments = supported comparisons; ticks = observed patients



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Fatty acid metabolism across age

One shared age axis · separate adjusted curves for each study




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

G2M checkpoint across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Glycolysis across age


One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Age coverage: thick segments = supported comparisons; ticks = observed patients




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Hedgehog signaling across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Heme metabolism across age


One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Age coverage: thick segments = supported comparisons; ticks = observed patients




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Hypoxia across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Il2 stat5 signaling across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed


Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

IL6 JAK STAT3 signaling across age

One shared age axis · separate adjusted curves for each study




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Inflammatory response across age


One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Age coverage: thick segments = supported comparisons; ticks = observed patients



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Interferon alpha response across age


One shared age axis · separate adjusted curves for each study



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Interferon gamma response across age

One shared age axis · separate adjusted curves for each study




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

KRAS signaling DN across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

KRAS signaling UP across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Mitotic spindle across age


One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Age coverage: thick segments = supported comparisons; ticks = observed patients




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

MTORC1 signaling across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

MYC targets v1 across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

MYC targets v2 across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Myogenesis across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Notch signaling across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Oxidative phosphorylation across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed


Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

P53 pathway across age

One shared age axis · separate adjusted curves for each study




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Pancreas beta cells across age


One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Age coverage: thick segments = supported comparisons; ticks = observed patients




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Peroxisome across age


One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Age coverage: thick segments = supported comparisons; ticks = observed patients




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

PI3K AKT mTOR signaling across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed


Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Protein secretion across age


One shared age axis · separate adjusted curves for each study



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Reactive oxygen species pathway across age


One shared age axis · separate adjusted curves for each study



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Spermatogenesis across age


One shared age axis · separate adjusted curves for each study



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

TGF beta signaling across age

One shared age axis · separate adjusted curves for each study




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

TNFα signaling via NFκB across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Unfolded protein response across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed


Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

UV response DN across age


One shared age axis · separate adjusted curves for each study



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

UV response UP across age

One shared age axis · separate adjusted curves for each study




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

WNT beta catenin signaling across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.




Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.

Xenobiotic metabolism across age

One shared age axis · separate adjusted curves for each study

PROTECT ADTB GHL R2D2 TBSQUARE TB: solid nonTB: dashed

Age-varying TB difference: 0/5 studies at FDR < 0.05. Visible bends alone do not establish an age effect.



Color = study. Solid = TB; dashed = nonTB. Pale ribbons = pointwise 95% confidence intervals.
Blank segments lack enough support for a TB–nonTB comparison. Each study retains its own clinical mix.
Different people at different ages: these curves do not follow patients as they grow older.