
Adipogenesis


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Allograft rejection


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Androgen response


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Angiogenesis


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Apical junction


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Apical surface


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Apoptosis


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Bile acid metabolism


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Cholesterol homeostasis


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Coagulation


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Complement


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

DNA repair


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

E2F targets


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Epithelial mesenchymal transition


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Estrogen response early


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Estrogen response late


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Fatty acid metabolism


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

G2M checkpoint


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Glycolysis


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Hedgehog signaling


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Heme metabolism


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Hypoxia


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Il2 stat5 signaling


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

IL6 JAK STAT3 signaling


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Inflammatory response


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Interferon alpha response


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Interferon gamma response


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

KRAS signaling DN


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

KRAS signaling UP


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Mitotic spindle


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

MTORC1 signaling


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

MYC targets v1


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

MYC targets v2


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Myogenesis


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Notch signaling


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Oxidative phosphorylation


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

P53 pathway


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Pancreas beta cells


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Peroxisome


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

PI3K AKT mTOR signaling


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Protein secretion


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Reactive oxygen species pathway


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Spermatogenesis


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

TGF beta signaling


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

TNFα signaling via NFκB


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Unfolded protein response


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

UV response DN


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

UV response UP


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

WNT beta catenin signaling


No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.

Xenobiotic metabolism

No study has an age-varying TB difference at FDR < 0.05.



Same age and score scales in every row. Whiskers: pointwise 95% intervals at selected ages.
Small ticks: patient ages. Gaps: insufficient support. Curves compare different people at different ages.