
Adipogenesis

202 children (PROTECT) · 610 adults · separate fits · 102/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.41 SD.

Allograft rejection

202 children (PROTECT) · 610 adults · separate fits · 151/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.37 SD.

Androgen response

202 children (PROTECT) · 610 adults · separate fits · 42/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.37 SD.

Angiogenesis

202 children (PROTECT) · 610 adults · separate fits · 243/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.86 SD.


Apical junction


202 children (PROTECT) · 610 adults · separate fits · 131/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.86 SD.


Apical surface
202 children (PROTECT) · 610 adults · separate fits · 237/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.25
1.00
0.75
0.50
0.25
0.00
−0.25
−0.50
−0.75
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.79 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.49 SD.

Bile acid metabolism

202 children (PROTECT) · 610 adults · separate fits · 225/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.95 SD.

Cholesterol homeostasis

202 children (PROTECT) · 610 adults · separate fits · 151/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.15 SD.

Coagulation

202 children (PROTECT) · 610 adults · separate fits · 126/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.


Complement

202 children (PROTECT) · 610 adults · separate fits · 95/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.62 SD.


DNA repair
202 children (PROTECT) · 610 adults · separate fits · 58/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.0
0.5
0.0
−0.5
−1.0
−1.5
−2.0
−2.5
−3.0
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.

E2F targets

202 children (PROTECT) · 610 adults · separate fits · 48/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.

Epithelial mesenchymal transition

202 children (PROTECT) · 610 adults · separate fits · 256/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.83 SD.

Estrogen response early

202 children (PROTECT) · 610 adults · separate fits · 245/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.94 SD.

Estrogen response late

202 children (PROTECT) · 610 adults · separate fits · 240/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.93 SD.


Fatty acid metabolism


202 children (PROTECT) · 610 adults · separate fits · 151/812 patient dots outside view





Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable


Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.60 SD.


G2M checkpoint
202 children (PROTECT) · 610 adults · separate fits · 55/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
−2.0
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.


Glycolysis
202 children (PROTECT) · 610 adults · separate fits · 192/812 patient dots outside view
TB
nonTB
Pathway score (SD)
0.5
0.0
−0.5
−1.0
−1.5
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.68 SD.



Hedgehog signaling
202 children (PROTECT) · 610 adults · separate fits · 199/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.5
1.0
0.5
0.0
−0.5
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.69 SD.


Heme metabolism
202 children (PROTECT) · 610 adults · separate fits · 155/812 patient dots outside view
TB
nonTB
Pathway score (SD)
0.5
0.0
−0.5
−1.0
−1.5
−2.0
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.50 SD.


Hypoxia
202 children (PROTECT) · 610 adults · separate fits · 135/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.0
0.5
0.0
−0.5
−1.0
−1.5
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.83 SD.


Il2 stat5 signaling

202 children (PROTECT) · 610 adults · separate fits · 150/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.53 SD.


Inflammatory response

202 children (PROTECT) · 610 adults · separate fits · 226/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.61 SD.


Interferon alpha response
202 children (PROTECT) · 610 adults · separate fits · 80/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.45 SD.


Interferon gamma response

202 children (PROTECT) · 610 adults · separate fits · 127/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.52 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable


Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.72 SD.



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable


Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.94 SD.


Mitotic spindle
202 children (PROTECT) · 610 adults · separate fits · 93/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1
0
−1
−2
−3
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.34 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.50 SD.



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.28 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.91 SD.

Notch signaling

202 children (PROTECT) · 610 adults · separate fits · 191/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.21 SD.


Oxidative phosphorylation

202 children (PROTECT) · 610 adults · separate fits · 92/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.45 SD.



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.46 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 1.03 SD.

Peroxisome

202 children (PROTECT) · 610 adults · separate fits · 126/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.43 SD.


PI3K AKT mTOR signaling

202 children (PROTECT) · 610 adults · separate fits · 143/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.34 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.

Reactive oxygen species pathway

202 children (PROTECT) · 610 adults · separate fits · 80/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.

Spermatogenesis

202 children (PROTECT) · 610 adults · separate fits · 233/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.55 SD.

TGF beta signaling

202 children (PROTECT) · 610 adults · separate fits · 95/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.58 SD.

TNFα signaling via NFκB

202 children (PROTECT) · 610 adults · separate fits · 174/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.57 SD.


Unfolded protein response

202 children (PROTECT) · 610 adults · separate fits · 35/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.43 SD.



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.66 SD.


WNT beta catenin signaling

202 children (PROTECT) · 610 adults · separate fits · 131/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.31 SD.


Xenobiotic metabolism
202 children (PROTECT) · 610 adults · separate fits · 143/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.0
0.5
0.0
−0.5
−1.0
−1.5
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.79 SD.