



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.41 SD.


Allograft rejection

202 children (PROTECT) · 610 adults · separate fits · 151/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.37 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.37 SD.

Angiogenesis

202 children (PROTECT) · 610 adults · separate fits · 243/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.86 SD.

Apical junction

202 children (PROTECT) · 610 adults · separate fits · 131/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.86 SD.


Apical surface

202 children (PROTECT) · 610 adults · separate fits · 237/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.79 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.49 SD.

Bile acid metabolism

202 children (PROTECT) · 610 adults · separate fits · 225/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.95 SD.


Cholesterol homeostasis


202 children (PROTECT) · 610 adults · separate fits · 151/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable


Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.15 SD.


Coagulation
202 children (PROTECT) · 610 adults · separate fits · 126/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.0
0.5
0.0
−0.5
−1.0
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.


Complement
202 children (PROTECT) · 610 adults · separate fits · 95/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.0
0.5
0.0
−0.5
−1.0
−1.5
−2.0
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.62 SD.


DNA repair

202 children (PROTECT) · 610 adults · separate fits · 58/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.

Epithelial mesenchymal transition

202 children (PROTECT) · 610 adults · separate fits · 256/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.83 SD.

Estrogen response early

202 children (PROTECT) · 610 adults · separate fits · 245/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.94 SD.


Estrogen response late


202 children (PROTECT) · 610 adults · separate fits · 240/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.93 SD.


Fatty acid metabolism
202 children (PROTECT) · 610 adults · separate fits · 151/812 patient dots outside view
TB
nonTB
Pathway score (SD)
0.5
0.0
−0.5
−1.0
−1.5
−2.0
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.60 SD.


G2M checkpoint
202 children (PROTECT) · 610 adults · separate fits · 55/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
−2.0
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.68 SD.


Hedgehog signaling

202 children (PROTECT) · 610 adults · separate fits · 199/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.69 SD.



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.50 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.83 SD.

Il2 stat5 signaling

202 children (PROTECT) · 610 adults · separate fits · 150/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.


IL6 JAK STAT3 signaling

202 children (PROTECT) · 610 adults · separate fits · 173/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.53 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.61 SD.

Interferon alpha response

202 children (PROTECT) · 610 adults · separate fits · 80/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.45 SD.

Interferon gamma response

202 children (PROTECT) · 610 adults · separate fits · 127/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.52 SD.


KRAS signaling DN

202 children (PROTECT) · 610 adults · separate fits · 153/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.72 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.94 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.34 SD.



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.50 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.28 SD.

Myogenesis

202 children (PROTECT) · 610 adults · separate fits · 180/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.91 SD.

Notch signaling

202 children (PROTECT) · 610 adults · separate fits · 191/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.21 SD.


Oxidative phosphorylation

202 children (PROTECT) · 610 adults · separate fits · 92/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.45 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.46 SD.


Pancreas beta cells

202 children (PROTECT) · 610 adults · separate fits · 194/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 1.03 SD.


Peroxisome
202 children (PROTECT) · 610 adults · separate fits · 126/812 patient dots outside view
TB
nonTB
Pathway score (SD)
0.5
0.0
−0.5
−1.0
−1.5
−2.0
−2.5
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.43 SD.


PI3K AKT mTOR signaling

202 children (PROTECT) · 610 adults · separate fits · 143/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.34 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.55 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.58 SD.

TNFα signaling via NFκB

202 children (PROTECT) · 610 adults · separate fits · 174/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.57 SD.

Unfolded protein response

202 children (PROTECT) · 610 adults · separate fits · 35/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.43 SD.

UV response DN

202 children (PROTECT) · 610 adults · separate fits · 159/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.

UV response UP

202 children (PROTECT) · 610 adults · separate fits · 155/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.66 SD.


WNT beta catenin signaling

202 children (PROTECT) · 610 adults · separate fits · 131/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.31 SD.



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.79 SD.