

Interferon gamma response

202 children (PROTECT) · 610 adults · separate fits · 127/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.52 SD.


Interferon alpha response
202 children (PROTECT) · 610 adults · separate fits · 80/812 patient dots outside view
TB
nonTB
Pathway score (SD)
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
Observed ages
PROTECT
ADTB
GHL
R2D2
TBSQUARE
Combined
0
10
20
30
40
50
60
70
80
85
Patient age (years)
Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable
Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.45 SD.


TNFα signaling via NFκB

202 children (PROTECT) · 610 adults · separate fits · 174/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.57 SD.




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.53 SD.

Inflammatory response

202 children (PROTECT) · 610 adults · separate fits · 226/812 patient dots outside view




Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.61 SD.

Complement

202 children (PROTECT) · 610 adults · separate fits · 95/812 patient dots outside view



Bin tests, separate FDR families: filled = q < 0.05 (color = higher group) · open = q ≥ 0.05 · × = unavailable

Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.62 SD.