



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.41 SD.

Allograft rejection


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.37 SD.

Androgen response


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.37 SD.

Angiogenesis


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.86 SD.

Apical junction


202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.86 SD.

Apical surface

202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.79 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.49 SD.

Bile acid metabolism

202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.95 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.15 SD.

Coagulation


202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.

Complement

202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.62 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.

E2F targets


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.

Epithelial mesenchymal transition


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.83 SD.

Estrogen response early


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.94 SD.

Estrogen response late


202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.93 SD.

Fatty acid metabolism

202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.60 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.

Glycolysis


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.68 SD.

Hedgehog signaling


202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.69 SD.

Heme metabolism

202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.50 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.83 SD.

Il2 stat5 signaling


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.

IL6 JAK STAT3 signaling


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.53 SD.

Inflammatory response


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.61 SD.

Interferon alpha response


202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.45 SD.

Interferon gamma response


202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.52 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.72 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.94 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.34 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.50 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.28 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.91 SD.

Notch signaling


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.21 SD.

Oxidative phosphorylation


202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.45 SD.

P53 pathway


202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.46 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 1.03 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.43 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.34 SD.

Protein secretion


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.55 SD.

TGF beta signaling


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.58 SD.

TNFα signaling via NFκB


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.57 SD.

Unfolded protein response


202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.43 SD.

UV response DN

202 children (PROTECT) · 610 adults · separate fits · full patient score range




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.66 SD.

WNT beta catenin signaling


202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.31 SD.

Xenobiotic metabolism

202 children (PROTECT) · 610 adults · separate fits · full patient score range



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.79 SD.