





Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.41 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.37 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.37 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.86 SD.

Apical junction

202 children (PROTECT) · 610 adults · separate fits · 131/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.86 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.79 SD.

Apoptosis


202 children (PROTECT) · 610 adults · separate fits · 88/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.49 SD.

Bile acid metabolism

202 children (PROTECT) · 610 adults · separate fits · 225/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.95 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.15 SD.

Coagulation


202 children (PROTECT) · 610 adults · separate fits · 126/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.62 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.

Epithelial mesenchymal transition


202 children (PROTECT) · 610 adults · separate fits · 256/812 patient dots outside view



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.83 SD.

Estrogen response early


202 children (PROTECT) · 610 adults · separate fits · 245/812 patient dots outside view



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.94 SD.

Estrogen response late


202 children (PROTECT) · 610 adults · separate fits · 240/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.93 SD.

Fatty acid metabolism


202 children (PROTECT) · 610 adults · separate fits · 151/812 patient dots outside view



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.60 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.68 SD.

Hedgehog signaling


202 children (PROTECT) · 610 adults · separate fits · 199/812 patient dots outside view



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.69 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.50 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.83 SD.

Il2 stat5 signaling


202 children (PROTECT) · 610 adults · separate fits · 150/812 patient dots outside view



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.

IL6 JAK STAT3 signaling


202 children (PROTECT) · 610 adults · separate fits · 173/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.53 SD.

Inflammatory response


202 children (PROTECT) · 610 adults · separate fits · 226/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.61 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.45 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.52 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.72 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.94 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.34 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.50 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.28 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.91 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.21 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.45 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.46 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 1.03 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.43 SD.

PI3K AKT mTOR signaling


202 children (PROTECT) · 610 adults · separate fits · 143/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.34 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.36 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.44 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.55 SD.

TGF beta signaling


202 children (PROTECT) · 610 adults · separate fits · 95/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.58 SD.

TNFα signaling via NFκB


202 children (PROTECT) · 610 adults · separate fits · 174/812 patient dots outside view




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.57 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.43 SD.




Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.64 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.66 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.31 SD.



Dots: observed patients; lines/whiskers: adjusted means and pointwise 95% CIs. Coverage: sampled years. Cross-sectional.
Exploratory adult summary: fixed study mix and shared age shapes. Maximum study-omission shift: 0.79 SD.