

Classification of waste materials in very-high-resolution satellite imagery

State of the art and thesis proposal

Alessandro Potenza · M.Sc. Computer Science and Engineering, AI · PERIVALLON

Outline


1. Problem and task: what has to be recognised, in which images, over which materials (slides 3-7).
2. Literature search: how the search was run, what it returned, what was kept and why (slides 8-10).
3. State of the art: the relevant works, from site-level detection down to the direct predecessor, and what is missing (slides 11-21).
4. Proposed work: approach, available imagery, asbestos pilot, evaluation (slides 22-25).

Context

Illegal waste dumping is an environmental crime with direct public-health consequences. Agencies (ARPA) have limited inspection capacity, and priority depends on what is dumped: rubble, plastics and asbestos-cement imply very different hazards.

Detecting dump sites from images is mature. Recognising the material is not: the reference survey (50 works, 1987-2023, almost all RGB) marks it as an open problem.

One thesis in the group has addressed material classification directly (Alari 2024). This work starts from there.



Task definition

Objective: given a VHR satellite image of a suspected dump site, output the set of waste materials present.


Task: multi-label classification at image level.

Input: VHR optical imagery, GSD 0.2 to 1.3 m. Aerial RGB for the baseline (AerialWaste); satellite VNIR for the multispectral arm, up to 8 bands. SWIR excluded: not in the planned acquisitions, and its 3.7 m GSD does not match the task. Sentinel-2 excluded: too coarse (10-20 m).

Technique: classification, not object detection or segmentation. Annotations are image-level (Alari 2024: 11,477 multi-label); waste piles have no stable shape for boxes; no segmentation masks exist for these datasets.

Research question: does VNIR input improve waste-material classification over RGB, and for which materials?

Task scheme




One image in, a set of material labels out. The same scheme runs with RGB or VNIR input; only the first layer of the network changes.

The material taxonomy



Rubble



Bulky items



Wood




Scrap




Plastic



Vehicles



Tires



Big Bags



Closed Containers



Unknown material

Ten-category grouping used in the group predecessor; the full annotation set counts 13 categories (adds foundry waste, sludge, tanks as separate classes).


Materials: which are considered and why

Material	Target	Spectral analysis	Reason
Rubble, plastic, wood, tires	yes	yes	frequent and confusable in RGB
Asbestos-cement	yes	yes, dedicated pilot	public regional ground truth (WFS)
Vehicles, tanks, containers, scrap, bulky, big bags	yes	no	shape-based; RGB sufficient in literature
Sludge, foundry waste	yes	no	few labels; visually ambiguous at this GSD

All 13 categories remain classification targets, for continuity with the group dataset. The RGB-versus-VNIR analysis concentrates on the subset where colour is ambiguous and labels are sufficient.

Literature search: how it was run

Scripted queries on the Scopus API (waste detection in remote sensing; asbestos roof mapping), deduplication, manual screening with explicit criteria, snowballing from the Fraternali 2024 survey and the Alari 2024 references.



What the search returned

Topic	Works at task scale	Notes
Waste sites, no material distinction	Torres 2023, Gibellini 2025, Sun 2023, CascadeDumpNet 2024, CWLD 2024, Disaitek 2024	mature line; answers where, not what
Multi-material waste classification	Alari 2024 (group thesis)	the only direct precedent
Asbestos-cement	Saba 2026, Bonifazi 2026, Abbasi 2024, Cilia 2015	roofs only, never inside a waste taxonomy
Tanks	Ramachandran 2024, YOLOv7-OT 2024	mature object detection
Vehicles, containers	vehicle DA 2020, truck-and-container 2025	VHR object detection
Scrap	ELV Hybrid-YOLOv5 2025	close-range infrared, not satellite
Rubble / inert	CWLD 2024; debris volume UAV 2022	C&D segmentation; UAV photogrammetry
Plastic, wood, tires, bulky, big bags, sludge, foundry	none found in scope	only as classes in AerialWaste / Alari

What was kept, what was excluded

Excluded group	Examples	Reason
Sentinel-2 and marine debris	MARIDA 2022, Tisza 2023	10-20 m: pixels too mixed; different domain
Spaceborne hyperspectral	Shepherd 2025 (EnMAP), EMIT 2025	material evidence, but 30-60 m GSD
SWIR-based VHR	Aguilar 2021, Guo and Li 2020, Zhou 2021, Aguilar 2025	SWIR not in our acquisitions; 3.7 m GSD
Laboratory / spectral libraries	Vitek 2025, SpectralWaste 2024, Knaeps 2020	not Earth observation; kept as band evidence
EO foundation models	DOFA 2024, AnySat 2025, Prithvi-EO-2 2024, SpectralGPT 2024, +6	pretrained at 10-30 m; sub-metre transfer unproven

Kept: works on terrestrial waste or roof materials at task-compatible GSD, plus the reference spectral library. The excluded papers stay in the annotated library as context.

Related work: site-level waste detection

Work (year)	Input / GSD	Task	Method	Result
Gibellini 2025	aerial RGB, 20 cm	classification	Swin-T, RSP pretraining	F1 92.0; cross-region -5.1
AW ensemble 2025 *	aerial RGB	classification	CNN + transformer ensemble	binary F1 92.4
Disaitek 2024	Pléiades Neo, 30 cm	detection	operational service	~95% on sites >2 m2 (vendor)
CascadeDumpNet 2024	Pléiades, 50 cm	object detection	two-stage CNN + AutoML	mAP 84.6, cross-city transfer
Sun 2023	VHR RGB, 0.3-1 m	object detection	BCA-Net (Faster R-CNN)	~2,500 dumpsites, 28 cities
CWLD 2024	GF-2 + GE, 0.5-0.8 m	segmentation	DeepLabV3+ variant	F1 88.9, construction waste
AerialWaste, Torres 2023	aerial RGB, 20-50 cm	dataset + cls.	ResNet-FPN baseline	F1 80.7; 22 material tags

AerialWaste: the reference dataset

10,434 locations from ARPA Lombardia records, 487 municipalities; three sources: AGEA orthophotos (20 cm), WorldView-3 (30 cm), Google Earth (50 cm).

Binary site labels for detection, plus 22 material tags (type of visible object + storage mode) annotated by experts.

Material tags cover about 72% of positives, but the dataset was published for detection: material tags are metadata, not a benchmark.

This is the label base both Gibellini 2025 and Alari 2024 build on.



Gibellini 2025: the site-level baseline

Binary waste / no-waste classification on AerialWaste; Swin-T with remote-sensing pretraining, two-step fine-tuning.

F1 92.02 in domain. Cross-region generalisation drops 5.1 points on average (Greece 85.4, Serbia 83.8, Romania 91.5).

Saliency maps confirm the model looks at waste heaps, but the output is presence only: no material information.

This architecture and training recipe is the starting point of the proposal.



Related work: objects and close range

Work (year)	Input	Task	Result
Ramachandran 2024	VHR satellite	object detection	tanks, P 0.96 / R 0.97, >169k mapped
YOLOv7-OT 2024 *	VHR satellite	object detection	tanks, precision 95.9
Truck-and-container 2025 *	VHR satellite	object detection	container size and status
Vehicle DA 2020 *	VHR 30-50 cm	object detection	+10% with domain adaptation
ELV Hybrid-YOLOv5 2025	close-range IR	object detection	scrap metals, mAP 84.2
UAV solid waste 2024 *	UAV RGB	segmentation	OA >94, generic waste piles
C&D debris UAV 2022 *	UAV + photogrammetry	segmentation	IoU 0.90 + volume estimation
fCLIPSeg 2025	aerial RGB	segmentation	debris, Dice 0.70, event-agnostic
Plastic UAV-SWIR 2026 *	UAV hyperspectral SWIR	segmentation	plastic waste, cross-domain
Plastic UAV-IoT 2025 *	UAV RGB	object detection	plastic waste, edge deployment

The asbestos line, in detail

Saba 2026: WorldView-3, VNIR only, 32 classifiers compared; best Macro-F1 97.6 per-pixel. Red Edge and NIR carry the discrimination.

Bonifazi 2026: WorldView-3 multi-temporal workflow; building-level decisions from pixel classification; tracks roof removals between acquisitions.

Abbasi 2024: aerial RGB with temporal features, OA ~96: shape and time can substitute spectra at fine GSD.

Cilia 2015: airborne hyperspectral, PA 89 / UA 86, plus a weathering index from red/NIR: the degradation angle.

Shepherd 2025: spaceborne hyperspectral (EnMAP, 30 m), field-calibrated library, 86% match on exhaustive ground truth: the signature survives from orbit; the limit is resolution.

Asbestos slate on drone RGB (2023): partially recognisable by shape and texture alone.

Why it matters here: public ground truth (Lombardy WFS, 10,903 roofs) and VNIR evidence make asbestos the natural pilot.



Related work: material-level classification

Work (year)	Input / GSD	Task	Result
Alari 2024 (PoliMi)	satellite RGB	multi-label cls.	wF1 69.2 (5 cat.), 59.4 (10 cat.)
Saba 2026 *	WV-3 VNIR, 1.24 m	pixel cls.	asbestos, Macro-F1 97.6
Bonifazi 2026	WV-3 VNIR+SWIR	pixel cls.	asbestos roofs, removal tracking
Abbasi 2024 *	aerial RGB	OBIA cls.	asbestos, OA ~96 (shape + time)
Cilia 2015	airborne HSI, 3 m	pixel cls.	asbestos, PA 89 / UA 86

One multi-material predecessor; everything else addresses a single material, asbestos, on roofs. No work measures RGB versus VNIR on waste materials. The predecessor is examined next.

Alari 2024: the direct predecessor

The anchor reference of this thesis: the only work that frames waste-material recognition as multi-label classification, done in the same group, on the same imagery base.

Dataset: 11,477 multi-label annotations over 13 categories; 3,190 positive and 7,190 negative images, built on AerialWaste imagery and ARPA records.

Models: ResNet-50 and Swin backbones with FPN; three classification-head designs; weighted binary cross-entropy for label imbalance; different pretraining sources compared.

The taxonomy on slide 6 and the target materials on slide 7 come directly from this work.

Headline results: weighted F1 69.21 on five categories, 59.42 on ten. Input is RGB only; the spectral dimension is untouched.


Alari 2024: results by category

Growing the taxonomy from 5 to 10 categories costs 9.8 weighted-F1 points (69.2 to 59.4).

Five of ten classes stay below F1 50. The thesis attributes this to few annotations (tires, big bags), high intra-class variance (wood) and inter-class similarity in RGB.

The classes that hold are shape- or texture-distinctive (rubble, bulky items, containers); the ones that fail are those where RGB appearance is ambiguous.

Per-category behaviour, not the aggregate score, is where the margin is. This is the starting point of the proposal.



The in-scope state of the art at a glance

Work	Input	Task	Material info
Fraternali 2024 (survey)	-	survey, 50 works	declares the material gap
AerialWaste 2023	aerial RGB	dataset	22 tags, metadata only
Gibellini 2025	aerial RGB	classification	none (presence)
Alari 2024	satellite RGB	multi-label cls.	13 categories, wF1 59-69
AW ensemble 2025 *	aerial RGB	classification	none (binary)
Sun 2023 / CascadeDumpNet 2024	VHR RGB	object detection	none (sites)
Disaitek 2024 (vendor)	Pléiades Neo	detection	coarse type qualification
CWLD 2024	GF-2 + GE	segmentation	one material (C&D)
Saba / Bonifazi / Abbasi / Cilia	WV-3, aerial, HSI	pixel cls.	one material (asbestos)
Tanks / vehicles / scrap works	VHR, close range	object detection	shape, not composition

Where RGB falls short, and what VNIR adds


Different materials share the same colour at 0.3-1.3 m: plastic sheets, asbestos-cement and concrete all appear grey. The separation lies beyond 700 nm (spectral libraries: Kokaly 2017, Knaeps 2020).

In the group predecessor, moving from 5 to 10 material categories costs 9.8 F1 points (69.2 to 59.4, Alari 2024). Finer material distinctions are the hard part.

Red Edge and NIR separate vegetation, bare soil and weathered surfaces; in Saba 2026 they drive asbestos discrimination.

On 10 construction-and-demolition materials, RGB gives 0.87 accuracy; RGB plus two bands at 650-750 and 850-1000 nm gives 0.96, matching the full spectrum (Vitek 2025, laboratory study).

Whether this helps waste materials at task GSD, and which ones, has not been measured. It is the object of this thesis.




What is missing in the literature

1. Multi-material classification has one direct precedent, with ample margin: wF1 59-69 (Alari 2024).
2. No work measures the added value of VNIR bands over RGB for waste materials at very high resolution.
3. Results are reported as aggregate scores; per-material behaviour is not analysed.
4. Generalisation across regions is rarely evaluated. At site level it costs 5 F1 points (Gibellini 2025).
5. Asbestos is studied on roofs, in isolation, and never inside a waste-material taxonomy.


Proposed work: approach

Multi-label image classification, continuing the group line (Gibellini 2025, Alari 2024). Same taxonomy, input extended from RGB to VNIR.

Band ablation with everything else fixed: same architecture, same splits, only the input changes.



Available imagery



Panchromatic: 0.31 m (WorldView-3), 0.30 m (Pléiades Neo). Access: commercial, or free quota via ESA proposal.

This is the band budget of the study: no SWIR, no Sentinel-2. RGB baseline runs on AerialWaste (20-50 cm).

First step: the asbestos pilot

Asbestos is the entry material, for the reasons seen in the analysis (slide 15): a single material, public pixel-accurate labels, direct regulatory value.

1. Extract roof polygons from the Lombardy WFS registry (10,903 mapped asbestos-cement roofs) and pair them with the available imagery.
2. Build matched RGB and VNIR inputs for the same roofs; add negative roofs from regional building footprints.
3. Train the same classifier on both inputs; compare F1, precision, recall on held-out areas.
4. **Decision point: the measured VNIR delta on a single, well-labelled material tells whether the full multi-label extension is worth the acquisition cost.**

Proposed work: evaluation

Per-material F1 alongside weighted and macro averages:
aggregates hide exactly the classes this thesis targets.

Delta versus the RGB baseline for each band configuration, with
confidence intervals over repeated runs.

Generalisation: train and test on disjoint geographic areas, besides
the standard split.

Reference points: wF1 69.2 / 59.4 (Alari 2024); Macro-F1 97.6
(Saba 2026, per-pixel) as upper reference for the pilot, not directly
comparable.

If VNIR does not help a material, that is a documented negative
result with practical value for sensor choice.

	standard split	cross-region split
RGB	baseline F1	baseline F1
RGB + NIR	$\Delta F1$ per material	$\Delta F1$ per material
full VNIR	$\Delta F1$ per material	$\Delta F1$ per material

References (1/3): waste detection

- Alari 2024. Fighting environmental crime with deep learning: classifying waste materials from illegal landfills in satellite imagery. M.Sc. thesis, PoliMi, 10589/230633.
- Fraternali et al. 2024. Solid waste detection, monitoring and mapping in remote sensing images: a survey. Waste Management / arXiv:2402.09066.
- Gibellini et al. 2025. A deep learning pipeline for solid waste detection in remote sensing images. Waste Management Bulletin.
- Torres, Fraternali 2023. AerialWaste: a dataset for illegal landfill discovery in aerial images. Scientific Data 10:63.
- Sharmily et al. 2025. AerialWaste ensembles (preprint). · Zhang, Ma 2024. CascadeDumpNet. Remote Sensing of Environment 313.
- Sun et al. 2023. Revealing influencing factors on global waste distribution. Nature Communications 14:1444.
- CWLD 2024. Construction waste landfill dataset. Scientific Data. · Disaitek 2024, vendor report (Airbus).
- Tisza et al. 2023. Waste detection and change analysis from multispectral satellite imagery (Sentinel-2). · MARIDA 2022. Marine debris benchmark. PLOS ONE.

References (2/3): materials and spectra

- Saba et al. 2026. Asbestos-cement roofs from WorldView-3 VNIR. J. Hazardous Materials. · Bonifazi et al. 2026. Geomatics 6(3):41.
- Abbasi et al. 2024. Multi-temporal asbestos change detection. RSASE. · Cilia et al. 2015. ISPRS IJGI 4(2).
- Shepherd et al. 2025. Asbestos-cement rooftops with EnMAP hyperspectral data. Scientific Reports.
- Asbestos slate from drone imagery, 2023 (training-data study).
- Vitek et al. 2025. Critical wavelengths for construction and demolition waste materials. Resources, Conservation and Recycling.
- Kokaly et al. 2017. USGS Spectral Library v7 (splib07a). USGS DS 1035. · Knaeps et al. 2020. Hyperspectral plastics library. ESSD.
- EMIT 2025. Global-scale plastic detection from space. Geophys. Res. Lett. · Zhou et al. 2021. Plastic classifier. Remote Sensing of Environment.
- Guo, Li 2020. Normalized difference plastic index on WorldView-3. ISPRS JPRS. · Aguilar et al. 2025. Urban macroplastics with WorldView-3. · Aguilar et al. 2021. WV-3 VNIR/SWIR ablation. Remote Sensing.
- SpectralWaste 2024. Multimodal spectral sorting dataset. · Plastic UAV-SWIR segmentation, 2026. Remote Sensing. · Plastic UAV-IoT detection, 2025. J. Hazardous Materials Advances.

References (3/3): objects, platforms, excluded backbones

Ramachandran et al. 2024. Deep learning to map well pads and storage tanks. Nature Communications.

YOLOv7-OT 2024. Storage tank detection. Remote Sensing. · Truck-and-container detection from satellite imagery, 2025 (preprint).

Vehicle detection with domain adaptation in VHR imagery, 2020. · ELV Hybrid-YOLOv5 2025. Non-ferrous metals in end-of-life vehicles.

UAV solid-waste segmentation, 2024. · C&D debris volume from UAV photogrammetry, 2022. Drones. · AI-powered drones in waste management: systematic review, 2025. Drones.

fCLIPSeg 2025. Event-agnostic debris segmentation.

DOFA 2024 (Xiong et al.). · AnySat 2025. · Prithvi-EO-2.0 2024 (Szwarcman et al.). · SpectralGPT 2024. TPAMI.

SatMAE 2022 (Cong et al.). NeurIPS. · SSL4EO-S12 2023 (Wang et al.). · SoftCon 2024 (Wang et al.).

Corley et al. 2024. Revisiting pre-trained remote sensing benchmarks. · Foundation models for remote sensing: a survey, 2024. · DEFLECT 2025 (Thoreau et al.). ICCV.