

Linear Algebra for Data Science

Alejandro Moreno Becerra

September 28, 2026

Contents

Preface	iii
1 Systems of linear equations	1
1.0 Introduction	1
1.1 Solving systems of linear equations	1
2 Systems of linear equations	3
2.0 Introduction	3
2.1 Solving systems of linear equations	3

Preface

The purpose of this notes is being the reference for our Linear Algebra course this semester. This means they will have at least all content that will go into the final exam, but probably a little more. For any errors detected, please contact me at alejandro.morenobecerra@ceu.es.

1

Systems of linear equations

1.0 INTRODUCTION

Systems of linear equations represent the very first sets of equations that we learn to solve. As we go deeper into their understanding, we begin to unleash the power of linear algebra, only paired by its beauty.

1.1 SOLVING SYSTEMS OF LINEAR EQUATIONS

Gauss, Jordan

A linear equation is an expression of the form $a_1x_1 + \dots + a_nx_n = b$, where the coefficients a_i and the independent term b are scalars, that is, elements of the field $\mathbb{R}$ or $\mathbb{C}$, that will be specified beforehand, and the x_i are indeterminates or variables.

The equation $0 = 0$ is called *trivial* and the equation $0 = b$ with $b \neq 0$ is called *inconsistent*. Any other equation is called *consistent*.

Definition 1.1.2. A **vector** is an ordered tuple of scalars (numbers) of the field. Any of the scalars of a vector v is called a *coordinate* and if the vector has n coordinates that are, for example, real numbers, we say that $v \in \mathbb{R}^n$. We will write vectors horizontally like

$$v = \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix}.$$

Nevertheless, we will usually write $v = (v_1, \dots, v_n)^T$ to save space, meaning the same vector written horizontally like the one before. Later on, we will see that this v^T operation has a name and meaning.

A **solution** to a linear system like (S) is a vector $(c_1, \dots, c_n)$ such that

If we look from a solutions perspective, we can classify equations in three types:

1. The equation $0 = 0$ which is always true, and we call **trivial**.
2. The equation $0 = b$ with $b \neq 0$, which is always false and we call **inconsistent**.
3. Any other equation has one solution, and we call them **consistent**.

In the same manner, we can classify linear systems by their solutions:

1. If every vector is a solution of (S), we call it **trivial**. Note that this is

- only the case when all equations are trivial.
2. If no vector is a solution of (S) , we call it **inconsistent**. This can be the case without any inconsistent equations.
 3. In any other case (the system has solutions, but not every vector is a solution), we call (S) **consistent**:
 - If there is only one solution, we call (S) **determined**.
 - If there is more than one solution, we call (S) **underdetermined**.

Key idea. This is an optional framed box. Use `textblock` for an unframed block or `shadedbox` for a pale gray background.

Key idea. Shaded box

EXERCISES FOR SECTION 1.1

- 1.1.1** Prove that $T(-x) = -T(x)$ for every linear map T .
-

2

Systems of linear equations

2.0 INTRODUCTION

Systems of linear equations represent the very first sets of equations that we learn to solve. As we go deeper into their understanding, we begin to unleash the power of linear algebra, only paired by its beauty.

2.1 SOLVING SYSTEMS OF LINEAR EQUATIONS

Gauss, Jordan

A linear equation is an expression of the form $a_1x_1 + \dots + a_nx_n = b$, where the coefficients a_i and the independent term b are scalars, that is, elements of the **field** $\mathbb{R}$ or $\mathbb{C}$, that will be specified beforehand, and the x_i are indeterminates or variables.

The equation $0 = 0$ is called *trivial* and the equation $0 = b$ with $b \neq 0$ is called *inconsistent*. Any other equation is called *consistent*.

Definition 2.1.2. A **vector** is an ordered tuple of scalars (numbers) of the field. Any of the scalars of a vector v is called a *coordinate* and if the vector has n coordinates that are, for example, real numbers, we say that $v \in \mathbb{R}^n$. We will write vectors horizontally like

$$v = \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix}.$$

Nevertheless, we will usually write $v = (v_1, \dots, v_n)^T$ to save space, meaning the same vector written horizontally like the one before. Later on, we will see that this v^T operation has a name and meaning.

A **solution** to a linear system like (S) is a vector $(c_1, \dots, c_n)$ such that

If we look from a solutions perspective, we can classify equations in three types:

1. The equation $0 = 0$ which is always true, and we call **trivial**.
2. The equation $0 = b$ with $b \neq 0$, which is always false and we call **inconsistent**.
3. Any other equation has one solution, and we call them **consistent**.

In the same manner, we can classify linear systems by their solutions:

1. If every vector is a solution of (S), we call it **trivial**. Note that this is only the case when all equations are trivial.
2. If no vector is a solution of (S), we call it **inconsistent**. This can be the

case without any inconsistent equations.

3. In any other case (the system has solutions, but not every vector is a solution), we call (S) **consistent**:

If there is only one solution, we call (S) **determined**.

If there is more than one solution, we call (S) **underdetermined**.

Key idea. This is an optional framed box. Use `textblock` for an unframed block or `shadedbox` for a pale gray background.

Key idea. Shaded box

EXERCISES FOR SECTION 2.1

- 2.1.1** Prove that $T(-x) = -T(x)$ for every linear map T .
-

