

A naive model for the vertical motion of an air bubble in water

Alessandro Gentilini (5th January 2021)

Introduction

I attended courses in free and SCUBA diving and they explained us the basic gas laws, during a lesson I was asking myself: what is the motion of an air bubble generated at some depth in the sea?

A naive model

The motion equation in one dimension is

$$m \frac{d^2 y}{dt^2} = F$$

where t is time, y is the depth in the water and F sums all the relevant forces applied to the bubble. $y = 0$ is the water level and $y < 0$ is below the water.

F is the sum of just two component:

$$F = -m g + A$$

where m is the mass of the bubble, g is the gravity acceleration and A is the Archimede's force. The Archimede's force is the product of the density ρ of the water, the volume V of the bubble and the the gravity acceleration: $A = \rho V g$.

But the volume of the air bubble depends on the pressure p according to the perfect gas law:

$p V = n R T$ where n is the number of moles of air, R is the gas constant and T is the temperature.

The pressure p at depth y is due to the water column (so called hydrostatic pressure): $p = \rho g |y|$ (it's valid only when $y < 0$).

The number of moles n is related to the molar mass of the air M and the bubble mass m through $n = \frac{m}{M}$.

So

$$V = \frac{n R T}{p} = \frac{m}{M} \cdot \frac{R T}{\rho g |y|}$$

and

$$F = -m g + \rho V g$$

Substituting V in F I get

$$F = -m g + \rho g \cdot \frac{m}{M} \cdot \frac{R T}{\rho g |y|}$$

and I can simplify ρg (so this model is valid for any fluid? Uhmmm... first doubt about my naive model) to get

$$F = -m g + \frac{m}{M} \cdot \frac{R T}{|y|}$$

Now the sum of forces must be balanced by the product of the mass of the bubble and its acceleration:

$$m \frac{d^2 y}{dt^2} = -m g + \frac{m}{M} \cdot \frac{R T}{|y|}$$

I can simplify the mass of the bubble (uhm... second doubt about this model):

$$\frac{d^2 y}{dt^2} = -g + \frac{R T}{M} \cdot \frac{1}{|y|}$$

So let's try to solve it symbolically!

Solution of the differential equation

```
In[1]:= DSolve [y''[t] == -g + (R * T) / (M * Abs[y[t]]), y[t], t]
```

```
Out[1]:= Solve [ ∫1y[t]  $\frac{1}{\sqrt{c_1 + 2 \int_1^{K[2]} \frac{R T - g M \text{Abs}[K[1]]}{M \text{Abs}[K[1]]} dK[1]}}$  dK[2]2 == (t + c2)2, y[t]
```

OK, it seems it cannot be solved... so I look for a numerical solution; first of all I set some values (in SI units):

```
In[2]:= R := 8.31
        T := 293
        M := 28.97 / 1000
        g := 9.8
        h := -10
```

And then solve it numerically:

```
In[7]:= ysol = NDSolveValue [{y''[t] == -g + (R * T) / (M * Abs[y[t]]), y[0] == h, y'[0] == 0}, y, {t, 0, 10}]
```


```
NDSolveValue : At t == 0.04325234866474409`, step size is effectively zero; singularity or stiff system suspected.
```

```
Out[7]:= InterpolatingFunction [ { +  Domain : {{0., 0.0433 }}
Output : scalar }
```

I would say the problem is the *singularity* (even because I do not know what a *stiff system* is ☺).

This is the plot of the depth y vs time t :


```
In[8]:= Plot[ysol[t], {t, 0, 0.04}, AxesLabel -> {t, y}]
```



And this is the plot of the bubble radius multiplied by a constant k :

```
In[9]:= kradius[t_] := CubeRoot[-1/ysol[t]]
```


```
Plot[kradius[t], {t, 0, 0.04}, AxesLabel -> {t, k radius}]
```



Now I would like to draw the bubble in ten equispaced time instants (copied from <https://mathematica.stackexchange.com/a/207586>)

```
In[11]:= bubble[t_] := Circle[{0, ysol[t]}, kradius[t]]
```

```
In[12]:= Graphics[{{bubble /@ Array[# &, 10, {0, 0.04}]}, ImageSize -> 50]
```



Out[12]=




An animation

An animation (copied from <https://mathematica.stackexchange.com/a/17205>):

In[13]:= Manipulate [
 Graphics [{Circle [{0, ysol[t]}, kradius [t]]}, PlotRange -> {{-1, 1}, {-11, 0}}, {t, 0, 0.04}]

Out[13]=



What's wrong with this model?

The main problem I can see, based on my limited experience in observing bubbles, is that it neglects the “friction” of the water acting on the moving bubble: I remember that I saw bubbles starting as sphere but then deforming into mushroom-like shapes.

Then I have doubts about the mass of my bubble.

I think this model will make physicists laughing (and numerical analysts too)... but I am not able to grasp the 1917 formulation by Lord Rayleigh and my Mathematica trial license is soon expiring! If I had had more time I would have tried to solve the Lord Rayleigh's equation.

Further readings

Lord Rayleigh O.M. F.R.S. (1917) VIII. *On the pressure developed in a liquid during the collapse of a spherical cavity* , The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science, 34:200, 94-98, DOI: 10.1080/14786440808635681