

# Social Networks & Recommendation Systems

## XI. Agent-based models.

---

Grzegorz Siudem

Warsaw University of Technology



**European  
Funds**  
Knowledge Education Development

**Warsaw University  
of Technology**

**European Union**  
European Social Fund



MSc program in Data Science has been developed  
as a part of task 10 of the project  
„NERW PW. Science - Education - Development - Cooperation”  
co-funded by European Union from European Social Fund.

# Lecture

---

# Epidemics – introduction

## SIR model (Kermack, McKendrick, 1932)

- We assume a constant size of the population over time.

# Epidemics – introduction

## SIR model (Kermack, McKendrick, 1932)

- We assume a constant size of the population over time.
- Each individual is in one of three states:

# Epidemics – introduction

## SIR model (Kermack, McKendrick, 1932)

- We assume a constant size of the population over time.
- Each individual is in one of three states:
  - S – susceptible,
  - I – infected,
  - R – recovered/ removed.

# Epidemics – introduction

## SIR model (Kermack, McKendrick, 1932)

- We assume a constant size of the population over time.
- Each individual is in one of three states:
  - S – susceptible,
  - I – infected,
  - R – recovered/ removed.
- We only allow the following transitions:  $S \rightarrow I \rightarrow R$ .

# Epidemics – introduction

## SIR model (Kermack, McKendrick, 1932)

- We assume a constant size of the population over time.
- Each individual is in one of three states:
  - S – susceptible,
  - I – infected,
  - R – recovered/ removed.
- We only allow the following transitions:  $S \rightarrow I \rightarrow R$ .
- Classically we assume that every agent is in contact with each other, and then the variables describe the system in which  $\beta, \gamma > 0$

$$\begin{cases} \frac{dS}{dt} &= -\beta SI, \\ \frac{dI}{dt} &= \beta SI - \gamma I, \\ \frac{dR}{dt} &= \gamma I. \end{cases}$$



## Epidemics – other variants of the model

- *SIRS* model (e.g. computer viruses)

$$S \rightarrow I \rightarrow R \rightarrow S.$$

# Epidemics – other variants of the model

- *SIRS* model (e.g. computer viruses)

$$S \rightarrow I \rightarrow R \rightarrow S.$$

- *SEIR* model (e.g. rubella, sexually transmitted diseases)

$$S \rightarrow E \rightarrow I \rightarrow R.$$

# Epidemics – other variants of the model

- *SIRS* model (e.g. computer viruses)

$$S \rightarrow I \rightarrow R \rightarrow S.$$

- *SEIR* model (e.g. rubella, sexually transmitted diseases)

$$S \rightarrow E \rightarrow I \rightarrow R.$$

- *SIS* model

$$S \rightarrow I \rightarrow S.$$

## SIS model on the graphs: notation

- $I(t)$  number of infected nodes,

# SIS model on the graphs: notation

- $I(t)$  number of infected nodes,
- $S(t)$  number of susceptible nodes, wherein

$$S(t) + I(t) = N.$$

# SIS model on the graphs: notation

- $I(t)$  number of infected nodes,
- $S(t)$  number of susceptible nodes, wherein

$$S(t) + I(t) = N.$$

- $\beta$  probability that in a single time step a healthy individual will become infected from a sick neighbor.

# SIS model on the graphs: notation

- $I(t)$  number of infected nodes,
- $S(t)$  number of susceptible nodes, wherein

$$S(t) + I(t) = N.$$

- $\beta$  probability that in a single time step a healthy individual will become infected from a sick neighbor.
- $\gamma$  the probability that the sick agent will recover in a single time step.

# SIS model on the graphs: notation

- $I(t)$  number of infected nodes,
- $S(t)$  number of susceptible nodes, wherein

$$S(t) + I(t) = N.$$

- $\beta$  probability that in a single time step a healthy individual will become infected from a sick neighbor.
- $\gamma$  the probability that the sick agent will recover in a single time step.
- Model can be described, in the mean-field approach, as follows

$$\frac{dI(t)}{dt} = \left[ \beta \left( \langle k \rangle \frac{I(t)}{N} \right) \right] S(t) - \gamma I(t).$$



Let's change variables

$$\frac{di(t)}{dt} = i(t) [\beta \langle k \rangle s(t) - \gamma],$$

where  $i(t) = I(t)/N$  and  $s(t) = S(t)/N$

Let's change variables

$$\frac{di(t)}{dt} = i(t) [\beta \langle k \rangle s(t) - \gamma],$$

where  $i(t) = I(t)/N$  and  $s(t) = S(t)/N$

Let us ask two questions:

- Under what conditions does an epidemic burst?
- What happens to the number of infected people in the large time limit?

## SIS model on the graphs: steady state

$$\frac{di(t)}{dt} = i(t) [\beta \langle k \rangle s(t) - \gamma],$$

Under what conditions does an epidemic burst?

- Let us compute  $di(t)/dt|_{t=0}$ .

## SIS model on the graphs: steady state

$$\frac{di(t)}{dt} = i(t) [\beta \langle k \rangle s(t) - \gamma],$$

Under what conditions does an epidemic burst?

- Let us compute  $di(t)/dt|_{t=0}$ .
- Received epidemic threshold

$$\lambda_c = \frac{1}{\langle k \rangle s(0)},$$

should be compared with the model parameters

$$\lambda = \frac{\beta}{\gamma}.$$

# SIS model on the graphs: steady state

$$\frac{di(t)}{dt} = i(t) [\beta \langle k \rangle s(t) - \gamma],$$

Under what conditions does an epidemic burst?

- Let us compute  $di(t)/dt|_{t=0}$ .
- Received epidemic threshold

$$\lambda_c = \frac{1}{\langle k \rangle s(0)},$$

should be compared with the model parameters

$$\lambda = \frac{\beta}{\gamma}.$$

Conclusions?

## SIS model on the graphs: epidemic threshold

$$\frac{di(t)}{dt} = i(t) [\beta \langle k \rangle s(t) - \gamma],$$

What happens to the number of infected people in the large time limit?

- We are looking for stable solutions:

$$\left. \frac{di(t)}{dt} \right|_{t \rightarrow \infty} = 0.$$

# SIS model on the graphs: epidemic threshold

$$\frac{di(t)}{dt} = i(t) [\beta \langle k \rangle s(t) - \gamma],$$

What happens to the number of infected people in the large time limit?

- We are looking for stable solutions:

$$\frac{di(t)}{dt} \Big|_{t \rightarrow \infty} = 0.$$

- We get two solutions

$$i_{\infty} = 0, \quad i_{\infty} = 1 - \frac{\gamma}{\beta \langle k \rangle}.$$

# SIS model on the graphs: epidemic threshold

$$\frac{di(t)}{dt} = i(t) [\beta \langle k \rangle s(t) - \gamma],$$

What happens to the number of infected people in the large time limit?

- We are looking for stable solutions:

$$\frac{di(t)}{dt} \Big|_{t \rightarrow \infty} = 0.$$

- We get two solutions

$$i_{\infty} = 0, \quad i_{\infty} = 1 - \frac{\gamma}{\beta \langle k \rangle}.$$

Conclusions?



# SIS model on scale-free networks

The analysis needs to be a bit more subtle

- $i_k(t)$ ,  $l_k(t)$  number (ratio) of infected nodes with degree  $k$ ,

The analysis needs to be a bit more subtle

- $i_k(t)$ ,  $I_k(t)$  number (ratio) of infected nodes with degree  $k$ ,
- $s_k(t)$ ,  $S_k(t)$  number (ratio) of susceptible nodes with degree  $k$ ,

# SIS model on scale-free networks

The analysis needs to be a bit more subtle

- $i_k(t)$ ,  $I_k(t)$  number (ratio) of infected nodes with degree  $k$ ,
- $s_k(t)$ ,  $S_k(t)$  number (ratio) of susceptible nodes with degree  $k$ ,
- $Q_I$  probability that any random network's edge is connected to infected vertex (for ER graphs  $Q_I(t) = I(t)/N$ ).

# SIS model on scale-free networks

The analysis needs to be a bit more subtle

- $i_k(t)$ ,  $I_k(t)$  number (ratio) of infected nodes with degree  $k$ ,
- $s_k(t)$ ,  $S_k(t)$  number (ratio) of susceptible nodes with degree  $k$ ,
- $Q_I$  probability that any random network's edge is connected to infected vertex (for ER graphs  $Q_I(t) = I(t)/N$ ).
- Furthermore we assume

$$I_k(t) + S_k(t) = N_k = N\mathcal{P}(k).$$

# SIS model on scale-free networks

The analysis needs to be a bit more subtle

- $i_k(t)$ ,  $I_k(t)$  number (ratio) of infected nodes with degree  $k$ ,
- $s_k(t)$ ,  $S_k(t)$  number (ratio) of susceptible nodes with degree  $k$ ,
- $Q_I$  probability that any random network's edge is connected to infected vertex (for ER graphs  $Q_I(t) = I(t)/N$ ).
- Furthermore we assume

$$I_k(t) + S_k(t) = N_k = N\mathcal{P}(k).$$

Then the equation takes the form

$$\frac{di_k(t)}{dt} = \beta k Q_I s_k(t) - \gamma i_k(t).$$

## SIS model on scale-free networks – asymptotic

$$\frac{di_k(t)}{dt} = \beta k Q_I s_k(t) - \gamma i_k(t).$$

We are looking for the stable asymptotic solutions

$$\left. \frac{di_k(t)}{dt} \right|_{t \rightarrow \infty} = 0 \Rightarrow i_k = \frac{\lambda k Q_I}{1 + \lambda k Q_I}.$$

$$\frac{di_k(t)}{dt} = \beta k Q_I s_k(t) - \gamma i_k(t).$$

We are looking for the stable asymptotic solutions

$$\left. \frac{di_k(t)}{dt} \right|_{t \rightarrow \infty} = 0 \Rightarrow i_k = \frac{\lambda k Q_I}{1 + \lambda k Q_I}.$$

Analysis of the solution leads to

$$\lambda_c = \frac{\langle k \rangle}{\langle k^2 \rangle}.$$

# SIS model on scale-free networks – asymptotic

$$\frac{di_k(t)}{dt} = \beta k Q_I s_k(t) - \gamma i_k(t).$$

We are looking for the stable asymptotic solutions

$$\left. \frac{di_k(t)}{dt} \right|_{t \rightarrow \infty} = 0 \Rightarrow i_k = \frac{\lambda k Q_I}{1 + \lambda k Q_I}.$$

Analysis of the solution leads to

$$\lambda_c = \frac{\langle k \rangle}{\langle k^2 \rangle}.$$

Which is very bad news...

Why?



## BA algorithm modification

- We are looking for citations vector of scientists with  $C$  citations and  $N$  publications.

## BA algorithm modification

- We are looking for citations vector of scientists with  $C$  citations and  $N$  publications.
- At each time step, the agent publishes works that are cited according to the rich get richer.

## BA algorithm modification

- We are looking for citations vector of scientists with  $C$  citations and  $N$  publications.
- At each time step, the agent publishes works that are cited according to the rich get richer.
- Is it consistent with empirical data? (visualization)

- In every graph's node we put variable  $\sigma_i = \pm 1$  (opinion, preferences, etc.).

- In every graph's node we put variable  $\sigma_i = \pm 1$  (opinion, preferences, etc.).
- At every time step random vertex change its opinion according neighbors' opinions

$$m_i = \sum_{j \in \mathcal{N}(i)} \sigma_j,$$

due to the formula

$$S_i := S(m_i) = \text{sgn}(m_i).$$

- In every graph's node we put variable  $\sigma_i = \pm 1$  (opinion, preferences, etc.).
- At every time step random vertex change its opinion according neighbors' opinions

$$m_i = \sum_{j \in \mathcal{N}(i)} \sigma_j,$$

due to the formula

$$S_i := S(m_i) = \text{sgn}(m_i).$$

## Illustration

Thank you for your attention!



**European  
Funds**  
Knowledge Education Development

**Warsaw University  
of Technology**

**European Union**  
European Social Fund



MSc program in Data Science has been developed  
as a part of task 10 of the project  
„NERW PW. Science - Education - Development - Cooperation”  
co-funded by European Union from European Social Fund.